

# F3 IAL Model Answers Kprime 2

June 2016

1. The curve  $C$  has equation

$$y = 9 \cosh x + 3 \sinh x + 7x$$

Use differentiation to find the exact  $x$  coordinate of the stationary point of  $C$ , giving your answer as a natural logarithm. (6)

$$1. \frac{dy}{dx} = 9 \sinh x + 3 \cosh x + 7$$

$$\frac{dy}{dx} = 0 \Rightarrow 9 \sinh x + 3 \cosh x + 7 = 0$$

$$\therefore \frac{9e^x - 9e^{-x} + 3e^x + 3e^{-x}}{2} = -7$$

$$\therefore 12e^x - 6e^{-x} = -14$$

$$\textcircled{xe^x} \Rightarrow 12e^{2x} + 14e^x - 6 = 0$$
$$6e^{2x} + 7e^x - 3 = 0$$

$$(3e^x - 1)(2e^x + 3) = 0$$

$$e^x \neq -\frac{3}{2}$$

$$\Rightarrow e^x = \frac{1}{3}$$

$$\therefore x = \ln\left(\frac{1}{3}\right)$$

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2. An ellipse has equation

$$\frac{x^2}{25} + \frac{y^2}{4} = 1$$

The point  $P$  lies on the ellipse and has coordinates  $(5 \cos \theta, 2 \sin \theta)$ ,  $0 < \theta < \frac{\pi}{2}$

The line  $L$  is a normal to the ellipse at the point  $P$ .

(a) Show that an equation for  $L$  is

$$5x \sin \theta - 2y \cos \theta = 21 \sin \theta \cos \theta \quad (5)$$

Given that the line  $L$  crosses the  $y$ -axis at the point  $Q$  and that  $M$  is the midpoint of  $PQ$ ,

(b) find the exact area of triangle  $OPM$ , where  $O$  is the origin, giving your answer as a multiple of  $\sin 2\theta$

(6)

$$2(a) \text{ @ } P, \quad \frac{\partial y}{\partial x} = \frac{\partial y / \partial \theta}{\partial x / \partial \theta} = \frac{2 \cos \theta}{-5 \sin \theta}$$

$$\therefore \text{gradient of normal} = \frac{5 \sin \theta}{2 \cos \theta}$$

$$y - y_1 = m(x - x_1)$$

$$\Rightarrow y - 2 \sin \theta = \frac{5 \sin \theta}{2 \cos \theta} (x - 5 \cos \theta)$$

$$(x - 5 \cos \theta) \Rightarrow 2y \cos \theta - 4 \sin \theta \cos \theta = 5x \sin \theta - 25 \sin \theta \cos \theta$$

$$\therefore 5x \sin \theta - 2y \cos \theta = 25 \sin \theta \cos \theta - 4 \sin \theta \cos \theta$$

$$\therefore 5x \sin \theta - 2y \cos \theta = 21 \sin \theta \cos \theta$$

as required.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

(b) @ Q


$$x=0 \Rightarrow -2y \cos \theta = 21 \sin \theta \cos \theta$$

$$\therefore y_Q = -\frac{21 \sin \theta}{2}$$

$$Q: \left(0, -\frac{21}{2} \sin \theta\right) \quad P: (5 \cos \theta, 2 \sin \theta)$$

$\therefore M$  has coordinates  $x_M = \frac{5 \cos \theta}{2}$

$$y_M = \frac{2 \sin \theta - \frac{21}{2} \sin \theta}{2} = -\frac{17 \sin \theta}{4}$$



$$\text{Area} = \left| \frac{1}{2} \left( \frac{21 \cos \theta}{5} \right) (2 \sin \theta) \right| + \left| \frac{1}{2} \left( \frac{21 \cos \theta}{5} \right) \left( -\frac{17 \sin \theta}{4} \right) \right|$$

$$= \frac{21}{5} \sin \theta \cos \theta + \frac{357}{40} \sin \theta \cos \theta = \frac{105}{8} \sin \theta \cos \theta$$

$$\text{Area} = \frac{105}{16} \sin 2\theta$$



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

3. Without using a calculator, find

(a)  $\int_{-2}^1 \frac{1}{x^2 + 4x + 13} dx$ , giving your answer as a multiple of  $\pi$ , (5)

(b)  $\int_{-1}^4 \frac{1}{\sqrt{4x^2 - 12x + 34}} dx$ , giving your answer in the form  $p \ln(q + r\sqrt{2})$ ,

where  $p$ ,  $q$  and  $r$  are rational numbers to be found. (7)

3.(a).

$$\int_{-2}^1 \frac{1}{x^2 + 4x + 13} dx = \int_{-2}^1 \frac{1}{(x+2)^2 + 9} dx$$

$$= \left[ \frac{1}{3} \arctan\left(\frac{x+2}{3}\right) \right]_{-2}^1$$

$$= \frac{1}{3} \arctan(1) - \frac{1}{3} \arctan(0)$$

$$= \frac{1}{3} \arctan(1) = \frac{\pi}{12}$$

(b)  $4x^2 - 12x + 34 = 4\left(x^2 - 3x + \frac{17}{2}\right)$

$$= 4\left[\left(x - \frac{3}{2}\right)^2 + \frac{25}{4}\right]$$



Question 3 continued

$$= \int_{-1}^4 \frac{1}{\sqrt{4x^2 - 12x + 34}} dx = \int_{-1}^4 \frac{1}{\sqrt{4\left(x - \frac{3}{2}\right)^2 + \frac{25}{4}}} dx$$

$$= \frac{1}{2} \int_{-1}^4 \frac{1}{\sqrt{\left(x - \frac{3}{2}\right)^2 + \frac{25}{4}}} dx$$

$$= \frac{1}{2} \left[ \operatorname{arsinh} \frac{x - \frac{3}{2}}{5/2} \right]_{-1}^4 = \frac{1}{2} \left[ \operatorname{arsinh} \frac{2x - 3}{5} \right]_{-1}^4$$

$$= \frac{1}{2} \operatorname{arsinh}(1) - \frac{1}{2} \operatorname{arsinh}(-1)$$

$$= \frac{1}{2} \ln(1 + \sqrt{2}) - \frac{1}{2} \ln(-1 + \sqrt{2})$$

$$= \frac{1}{2} \ln(3 + 2\sqrt{2})$$

4.

$$M = \begin{pmatrix} 1 & k & 0 \\ -1 & 1 & 1 \\ 1 & k & 3 \end{pmatrix}, \text{ where } k \text{ is a constant}$$

(a) Find  $M^{-1}$  in terms of  $k$ .

(5)

Hence, given that  $k = 0$

(b) find the matrix  $N$  such that

$$MN = \begin{pmatrix} 3 & 5 & 6 \\ 4 & -1 & 1 \\ 3 & 2 & -3 \end{pmatrix}$$

(4)

$$4(a). \det(M) = \begin{vmatrix} 1 & 1 \\ k & 3 \end{vmatrix} - k \begin{vmatrix} -1 & 1 \\ 1 & 3 \end{vmatrix}$$

$$= 3 - k - k(-4) = \underline{3 + 3k}$$

$$\begin{pmatrix} 3-k & -4 & -k-1 \\ 3k & 3 & 0 \\ k & 1 & k+1 \end{pmatrix}$$

$$\text{apply } \begin{pmatrix} + & - & + \\ - & + & - \\ + & - & + \end{pmatrix}^T \Rightarrow \begin{pmatrix} 3-k & 4 & -k-1 \\ -3k & 3 & 0 \\ k & -1 & k+1 \end{pmatrix}^T$$

$$= \begin{pmatrix} 3-k & -3k & k \\ 4 & 3 & -1 \\ -k-1 & 0 & k+1 \end{pmatrix}$$



$$\therefore M^{-1} = \frac{1}{3k+3} \begin{pmatrix} 3-k & -3k & k \\ 4 & 3 & -1 \\ -k-1 & 0 & k+1 \end{pmatrix}$$

$$(b) k=0 \Rightarrow M^{-1} = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 \\ 4 & 3 & -1 \\ -1 & 0 & 1 \end{pmatrix}$$

$$M^{-1}MN = N$$

$$\Rightarrow N = \frac{1}{3} \begin{pmatrix} 3 & 0 & 0 \\ 4 & 3 & -1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 5 & 6 \\ 4 & -1 & 1 \\ 3 & 2 & -3 \end{pmatrix}$$

$$= \frac{1}{3} \begin{pmatrix} 9 & 15 & 18 \\ 21 & 15 & 30 \\ 0 & -3 & -1 \end{pmatrix}$$

$$N = \begin{pmatrix} 3 & 5 & 6 \\ 7 & 5 & 10 \\ 0 & -1 & -3 \end{pmatrix}$$

(Total 12 marks)

Q3



P 4 6 6 8 4 A 0 1 1 3 2

11

Turn over

5. Given that  $y = \operatorname{artanh}(\cos x)$

(a) show that

$$\frac{dy}{dx} = -\operatorname{cosec} x \quad (2)$$

(b) Hence find the exact value of

$$\int_0^{\frac{\pi}{6}} \cos x \operatorname{artanh}(\cos x) dx$$

giving your answer in the form  $a \ln(b + c\sqrt{3}) + d\pi$ , where  $a$ ,  $b$ ,  $c$  and  $d$  are rational numbers to be found. (5)

5. (a).  $y = \operatorname{ar} \tanh(\cos x)$

$$\therefore \tanh y = \cos x$$

$$\therefore \frac{dy}{dx} \operatorname{sech}^2 y = -\sin x$$

$$\operatorname{sech}^2 y = 1 - \tanh^2 y$$

$$\therefore \frac{dy}{dx} = \frac{-\sin x}{1 - \tanh^2 y}$$

$$\therefore \frac{dy}{dx} = \frac{-\sin x}{1 - \cos^2 x} = \frac{-\sin x}{\sin^2 x}$$

$$= \frac{-1}{\sin x} = -\operatorname{cosec} x$$

as required.





(b) Let  $u = \operatorname{arctanh}(\cos x)$      $u' = -\operatorname{cosec} x$

Let  $v' = \cos x$      $v = \sin x$

$$\therefore \int_0^{\pi/6} \cos x \operatorname{arctanh}(\cos x) dx$$

$$= \left[ \sin x \operatorname{arctanh}(\cos x) \right]_0^{\pi/6} + \int_0^{\pi/6} 1 dx$$

$$= \frac{1}{2} \operatorname{arctanh}\left(\frac{\sqrt{3}}{2}\right) + [x]_0^{\pi/6}$$

$$= \frac{1}{4} \ln(7+4\sqrt{3}) + \frac{\pi}{6}$$

---

---

6. The coordinates of the points  $A$ ,  $B$  and  $C$  relative to a fixed origin  $O$  are  $(1, 2, 3)$ ,  $(-1, 3, 4)$  and  $(2, 1, 6)$  respectively. The plane  $\Pi$  contains the points  $A$ ,  $B$  and  $C$ .

(a) Find a cartesian equation of the plane  $\Pi$ .

(5)

The point  $D$  has coordinates  $(k, 4, 14)$  where  $k$  is a positive constant.

Given that the volume of the tetrahedron  $ABCD$  is 6 cubic units,

(b) find the value of  $k$ .

(4)

$$\text{(a) } \vec{AB} = \begin{pmatrix} -1 \\ 3 \\ 4 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}$$

$$\vec{AC} = \begin{pmatrix} 2 \\ 1 \\ 6 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$$

$$\therefore \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix} \times \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} = \begin{matrix} -2 & 1 & 3 \\ 1 & -1 & 3 \\ -1 & 3 & -1 \end{matrix} \begin{matrix} \times \\ \times \\ \times \end{matrix} \begin{matrix} 1 & -1 & 3 \\ -1 & 3 & -1 \\ 1 & -1 & 3 \end{matrix}$$

$$\vec{n} = \begin{pmatrix} 4 \\ 7 \\ 1 \end{pmatrix}$$

$$\therefore \vec{r} \cdot \vec{n} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 7 \\ 1 \end{pmatrix} = 21$$

$$\therefore \underline{4x + 7y + z = 21}$$



$$(b) \frac{1}{6} \left| \vec{AD} \cdot (\vec{AB} \times \vec{AC}) \right| = 6$$

$$\therefore \left| \vec{AD} \cdot (\vec{AB} \times \vec{AC}) \right| = 36$$

$$\vec{AD} = \begin{pmatrix} k \\ 4 \\ 14 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} k-1 \\ 2 \\ 11 \end{pmatrix}$$

$$\begin{pmatrix} k-1 \\ 2 \\ 11 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ 7 \\ 1 \end{pmatrix} = 36$$

$$\Rightarrow 4k - 4 + 14 + 11 = 36$$

$$\therefore 4k = 15$$

$$\Rightarrow k = \frac{15}{4}$$

7. The curve  $C$  has parametric equations

$$x = 3t^4, \quad y = 4t^3, \quad 0 \leq t \leq 1$$

The curve  $C$  is rotated through  $2\pi$  radians about the  $x$ -axis. The area of the curved surface generated is  $S$ .

(a) Show that

$$S = k\pi \int_0^1 t^5 (t^2 + 1)^{\frac{1}{2}} dt$$

where  $k$  is a constant to be found.

(4)

(b) Use the substitution  $u^2 = t^2 + 1$  to find the value of  $S$ , giving your answer in the form  $p\pi(11\sqrt{2} - 4)$  where  $p$  is a rational number to be found.

(7)

$$\text{7(a). } S = 2\pi \int_0^1 y \sqrt{\left(\frac{\partial x}{\partial t}\right)^2 + \left(\frac{\partial y}{\partial t}\right)^2} dt$$

$$= 2\pi \int_0^1 4t^3 \sqrt{(12t^3)^2 + (12t^2)^2} dt$$

$$= 2\pi \int_0^1 4t^3 \sqrt{144t^6 + 144t^4} dt$$

$$= 2\pi \int_0^1 4t^3 \sqrt{144t^4(t^2 + 1)} dt$$

$$= 2\pi \int_0^1 48t^5 (t^2 + 1)^{1/2} dt$$

$$= \underset{k=96}{96\pi} \int_0^1 t^5 (t^2 + 1)^{1/2} dt$$

as required.



Question 7 continued

$$(b) \quad u^2 = t^2 + 1$$

$$\therefore 2u \frac{du}{dt} = 2t \Rightarrow \frac{du}{dt} = \frac{t}{u}$$

$$\therefore dt = \frac{u}{t} du = \frac{u}{u^2-1} du$$

$$\left. \begin{array}{l} t=1 \Rightarrow u^2=2 \Rightarrow u=\sqrt{2} \\ t=0 \Rightarrow u^2=1 \Rightarrow u=1 \end{array} \right\} \int \text{ becomes } \int$$

$$t^5 = (u^2-1)^{5/2}$$

$$\therefore S = 96\pi \int_0^1 t^5 (t^2+1)^{1/2} dt$$

$$= 96\pi \int_1^{\sqrt{2}} (u^2-1)^{5/2} u \cdot \frac{u}{u^2-1} du$$

$$= 96\pi \int_1^{\sqrt{2}} u^2 (u^2-1)^2 du$$

$$= 96\pi \int_1^{\sqrt{2}} u^2 (u^4 - 2u^2 + 1) du$$

$$= 96\pi \int_1^{\sqrt{2}} (u^6 - 2u^4 + u^2) du$$



Question 7 continued

$$= 96\pi \left[ \frac{1}{7}u^7 - \frac{2}{5}u^5 + \frac{1}{3}u^3 \right]_1^{\sqrt{2}}$$

$$= 96\pi \left[ u^3 \left( \frac{1}{7}u^4 - \frac{2}{5}u^2 + \frac{1}{3} \right) \right]_1^{\sqrt{2}}$$

$$= 96\pi \left[ 2\sqrt{2} \left( \frac{4}{7} - \frac{4}{5} + \frac{1}{3} \right) - \frac{8}{105} \right]$$

$$= 96\pi \left( \frac{22}{105}\sqrt{2} - \frac{8}{105} \right)$$

$$= 96\pi \times 2 \times \left( \frac{11\sqrt{2} - 4}{105} \right)$$

$$= \frac{64}{35}\pi (11\sqrt{2} - 4) \quad p = \frac{64}{35}$$

blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8.

$$I_n = \int_0^{\ln 2} \tanh^{2n} x \, dx, \quad n \geq 0$$

(a) Show that, for  $n \geq 1$

$$I_n = I_{n-1} - \frac{1}{2n-1} \left(\frac{3}{5}\right)^{2n-1} \quad (5)$$

(b) Hence show that

$$\int_0^{\ln 2} \tanh^4 x \, dx = p + \ln 2$$

where  $p$  is a rational number to be found.

(5)

8(a). ~~11~~

$$I_n = \int_0^{\ln 2} \tanh^{2n} x \, dx = \int_0^{\ln 2} \tanh^{2n-2} x \tanh^2 x \, dx$$

$$= \int_0^{\ln 2} \tanh^{2n-2} x (1 - \operatorname{sech}^2 x) \, dx$$

$$= \int_0^{\ln 2} \tanh^{2n-2} x - \operatorname{sech}^2 x \tanh^{2n-2} x \, dx$$


$$= \int_0^{\ln 2} \tanh^{2n-2} x \, dx - \int_0^{\ln 2} \operatorname{sech}^2 x (\tanh x)^{2n-2} \, dx$$

$$= \int_0^{\ln 2} \tanh x^{2(n-1)} \, dx - \left[ \frac{(\tanh x)^{2n-1}}{2n-1} \right]_0^{\ln 2}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

$$= I_{n-1} - \left[ \frac{(\tanh(\ln 2))^{2n-1}}{2n-1} - \frac{\tanh(b)^{2n-1}}{2n-1} \right]$$

$$= I_{n-1} - \left[ \frac{\left(\frac{3}{5}\right)^{2n-1}}{2n-1} - 0 \right]$$

 $\Rightarrow$ 

$$I_n = I_{n-1} - \frac{\left(\frac{3}{5}\right)^{2n-1}}{2n-1} \quad \text{as required.}$$

$$(b) \quad I_2 = I_1 - \frac{1}{3} \left(\frac{3}{5}\right)^3$$

$$I_1 = I_0 - \frac{1}{1} \left(\frac{3}{5}\right)$$

$$\therefore I_1 = I_0 - \frac{3}{5}$$

$$I_0 = \int_0^{\ln 2} \tanh^0 x dx = \int_0^{\ln 2} 1 dx = \underline{\underline{\ln 2}}$$





Question 8 continued

$$\therefore I_1 = \ln 2 - \frac{3}{5}$$

$$I_2 = \ln 2 - \frac{3}{5} - \frac{1}{3} \left( \frac{3}{5} \right)^3$$

$$= \frac{-84}{125} + \ln 2$$