

FP3 June 11 M.A. Perime 2

1. The curve C has equation $y = 2x^3$, $0 \leq x \leq 2$.

The curve C is rotated through 2π radians about the x -axis.

Using calculus, find the area of the surface generated, giving your answer to 3 significant figures.

(5)

$$y = 2x^3 \Rightarrow \frac{dy}{dx} = 6x^2$$

$$\text{Area} = 2\pi \int_0^2 y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$\therefore \text{Area} = 2\pi \int_0^2 2x^3 \sqrt{1 + 36x^4} dx$$

$$= 4\pi \int_0^2 x^3 (1 + 36x^4)^{1/2} dx$$

$$\int f'(x) f(x)^n dx = \frac{f(x)^{n+1}}{n+1}$$

$$= 4\pi = \frac{\pi}{36} \int_0^2 144x^3 (1 + 36x^4)^{1/2} dx$$

$$= \frac{\pi}{36} \left[\frac{(1 + 36x^4)^{3/2}}{3/2} \right]_0^2 = \frac{\pi}{36} \left[\frac{577^{3/2}}{3/2} - \frac{1}{3/2} \right]$$

$$= 806 \text{ units}^2$$

$$= \frac{\pi}{36} \left(\frac{73^{3/2}}{3/2} - \frac{1}{3/2} \right) \quad (3sf)$$

$$= 36.2 \text{ units}^2$$

2. (a) Given that $y = x \arcsin x$, $0 \leq x \leq 1$, find

(i) an expression for $\frac{dy}{dx}$,

(ii) the exact value of $\frac{dy}{dx}$ when $x = \frac{1}{2}$.

(3)

(b) Given that $y = \arctan(3e^{2x})$, show that

$$\frac{dy}{dx} = \frac{3}{5 \cosh 2x + 4 \sinh 2x}$$

(5)

2(a) $y = x \arcsin x$

(i) Product rule:

$$\frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}} + \arcsin x$$

(ii) $\left(\frac{dy}{dx}\right)_{x=1/2} = \frac{1/2}{\sqrt{1-1/4}} + \arcsin \frac{1}{2}$

$$= \frac{\sqrt{3}}{3} + \frac{\pi}{6} = \frac{2\sqrt{3} + \pi}{6}$$

(b) $y = \arctan(3e^{2x})$

~~use chain rule~~

~~$\frac{dy}{dx} = 3e^{2x}$~~

$\tan y = 3e^{2x}$

$$y = \arctan(3e^{2x})$$

$$\therefore \tan y = 3e^{2x}$$

$$\frac{s^2 + c^2}{c^2} = \frac{1}{c^2}$$

$$\therefore \frac{dy}{dx} \sec^2 y = 6e^{2x}$$

$$1 + t^2 = \sec^2$$

$$\therefore \frac{dy}{dx} (\tan^2 y + 1) = 6e^{2x}$$

$$\therefore \frac{dy}{dx} = \frac{6e^{2x}}{\tan^2 y + 1} = \frac{6e^{2x}}{9e^{4x} + 1}$$

$$\text{Let } t = \frac{dy}{dx} = \frac{6e^{2x}}{9e^{4x} + 1} \times \frac{\frac{1}{2e^{2x}}}{\frac{1}{2e^{2x}}}$$

$$= \frac{3}{\frac{9}{2}e^{2x} + \frac{1}{2}e^{-2x}} = \text{LHS}$$

$$\text{Let } \frac{dy}{dx} = \frac{3}{\frac{9}{2}e^{2x} + \frac{1}{2}e^{-2x}} = \text{LHS}$$

$$\text{RHS} = \frac{3}{5 \cosh 2x + 4 \sinh 2x} = \frac{3}{5 \cdot \frac{e^{2x} + e^{-2x}}{2} + 4 \cdot \frac{e^{2x} - e^{-2x}}{2}}$$

$$= \frac{3}{\frac{5e^{2x} + 5e^{-2x} + 4e^{2x} - 4e^{-2x}}{2}}$$

$$= \frac{3}{\frac{9e^{2x} + e^{-2x}}{2}}$$

$$= \frac{3}{\frac{9}{2}e^{2x} + \frac{1}{2}e^{-2x}}$$

= LHS

$$\therefore \frac{dy}{dx} = \frac{3}{5 \cosh 2x + 4 \sinh 2x}$$

as required

3. Show that

(a) $\int_5^8 \frac{1}{x^2 - 10x + 34} dx = k\pi$, giving the value of the fraction k , (5)

(b) $\int_5^8 \frac{1}{\sqrt{(x^2 - 10x + 34)}} dx = \ln(A + \sqrt{n})$, giving the values of the integers A and n . (4)

3(a) $\int_5^8 \frac{1}{x^2 - 10x + 34} dx = \int_5^8 \frac{1}{(x-5)^2 + 9}$

$= \left[\frac{1}{3} \arctan \frac{x-5}{3} \right]_5^8$

$= \frac{1}{3} \arctan 1 - \frac{1}{3} \arctan 0$

$= \frac{1}{12} \pi \quad k = \frac{1}{12}$

(b) $\int_5^8 \frac{1}{\sqrt{(x-5)^2 + 9}} dx = \left[\operatorname{arsinh} \frac{x-5}{3} \right]_5^8$

$= \operatorname{arsinh} 1 - \operatorname{arsinh} 0$

$= \ln(1 + \sqrt{2}) - \ln(0 + \sqrt{1})$

$= \ln(1 + \sqrt{2}) \quad A = 1 \quad n = 2$

4.

$$I_n = \int_1^e x^2 (\ln x)^n dx, \quad n \geq 0$$

(a) Prove that, for $n \geq 1$,

$$I_n = \frac{e^3}{3} - \frac{n}{3} I_{n-1} \quad (4)$$

(b) Find the exact value of I_3 . (4)

$$(a). \quad I_n = \int_1^e x^2 (\ln x)^n dx$$

$$\text{Let } u = (\ln x)^n \quad u' = \frac{n(\ln x)^{n-1}}{x}$$

$$v' = x^2 \quad v = \frac{1}{3} x^3$$

$$\therefore I_n = \left[\frac{1}{3} x^3 (\ln x)^n \right]_1^e - \frac{1}{3} n \int_1^e \frac{x^2 (\ln x)^{n-1}}{x} dx$$

$$= \frac{1}{3} e^3 - \frac{1}{3} n \int_1^e \frac{x^2 (\ln x)^{n-1}}{x} dx$$

$$= \frac{e^3}{3} - \frac{n}{3} \int_1^e x^2 (\ln x)^{n-1} dx$$

$$= \frac{e^3}{3} - \frac{n}{3} I_{n-1}$$

as required.

Question 4 continued

$$(b) \quad I_3 = \frac{e^3}{3} - I_2$$

$$= \frac{e^3}{3} - \left(\frac{e^3}{3} - \frac{2}{3} I_1 \right)$$

~~$$= \frac{e^3}{3} - \frac{2}{3} I_1$$~~

$$= \frac{2}{3} I_1 = \frac{2}{3} \int_1^e x^2 \ln x \, dx$$

$$\begin{array}{l} \underline{u=2} \\ \left. \begin{array}{l} u = \ln x \quad u' = \frac{1}{x} \\ v' = x^2 \quad v = \frac{1}{3} x^3 \end{array} \right\} \end{array}$$

~~$$= \frac{2}{3} \int$$~~

$$= \frac{2}{3} \left(\left[\frac{x^3}{3} \ln x \right]_1^e - \frac{1}{3} \int_1^e x^2 \, dx \right)$$


~~$$= \frac{2}{3} \left(\frac{e^3}{3} - \frac{1}{3} \left[\frac{1}{3} x^3 \right]_1^e \right)$$~~

$$= \frac{2}{3} \left(\frac{e^3}{3} - \frac{1}{3} \left(\frac{e^3}{3} - \frac{1}{3} \right) \right)$$

$$= \frac{2}{3} \left(\frac{2}{9} e^3 + \frac{1}{9} \right) = \frac{4}{27} e^3 + \frac{2}{27}$$

$$= \frac{4e^3 + 2}{27}$$

5. The curve C_1 has equation $y = 3\sinh 2x$, and the curve C_2 has equation $y = 13 - 3e^{2x}$.
- (a) Sketch the graph of the curves C_1 and C_2 on one set of axes, giving the equation of any asymptote and the coordinates of points where the curves cross the axes. (4)
- (b) Solve the equation $3\sinh 2x = 13 - 3e^{2x}$, giving your answer in the form $\frac{1}{2} \ln k$, where k is an integer. (5)



Question 5 continued

$$b) 3 \sinh 2x = 13 - 3e^{2x}$$

$$\frac{3e^{2x} - 3e^{-2x}}{2} = 13 - 3e^{2x}$$

$$\therefore 3e^{2x} - 3e^{-2x} = 26 - 6e^{2x}$$

$$\therefore 3e^{4x} - 3 = 26e^{2x} - 6e^{4x}$$

$$\therefore 9e^{4x} - 26e^{2x} - 3 = 0$$

$$\therefore (e^{2x} - 3)(9e^{2x} + 1) = 0$$

$$\therefore e^{2x} = 3 \Rightarrow 2x = \ln 3 \Rightarrow x = \frac{1}{2} \ln 3$$

~~$$e^{2x} = -\frac{1}{9} \Rightarrow 2x = \ln$$~~

6. The plane P has equation

$$r = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix}$$

(a) Find a vector perpendicular to the plane P .

(2)

The line l passes through the point $A(1, 3, 3)$ and meets P at $(3, 1, 2)$.

The acute angle between the plane P and the line l is α .

(b) Find α to the nearest degree.

(4)

(c) Find the perpendicular distance from A to the plane P .


(4)

6(a).

$$n = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix} \times \begin{pmatrix} 3 \\ 2 \\ 2 \end{pmatrix} = \begin{vmatrix} i & j & k \\ 0 & 2 & -1 \\ 3 & 2 & 2 \end{vmatrix} = \begin{pmatrix} 0 \cdot 2 - (-1) \cdot 2 \\ -1 \cdot 2 - 0 \cdot 3 \\ 0 \cdot 2 - 3 \cdot 2 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \\ -6 \end{pmatrix}$$

$$n = \begin{pmatrix} 6 \\ -3 \\ -6 \end{pmatrix}$$

(b)



$$n \cdot \begin{pmatrix} 6 \\ -3 \\ -6 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ -3 \\ -6 \end{pmatrix}$$

$$= 18 - 3 - 12$$

$$n \cdot n = 3$$

$$AP = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix} - \begin{pmatrix} 1 \\ 3 \\ 3 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \\ -1 \end{pmatrix}$$

Question 6 continued

$$\cos \theta = \frac{\begin{pmatrix} 2 \\ -2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ -3 \\ -6 \end{pmatrix}}{\sqrt{2^2+2^2+1} + \sqrt{6^2+3^2+6^2}} = \frac{24}{3 \times 9}$$




$$= \frac{8}{9}$$

$$\Rightarrow \theta = 27.266\dots$$

$\alpha = 90 - \theta$

~~$\alpha = 90 - 27^\circ$~~ (~~nearest degree~~)

$\alpha = 63^\circ$



(C)

$$|A| = \sqrt{2^2+2^2+1^2} = 3$$

$\sin \alpha = \frac{8}{9}$

$\therefore \sin \alpha = \frac{d}{3}$

$\cos \alpha =$

~~$d = 3 \sin \alpha = 3 \times \frac{8}{9} = \frac{8}{3}$~~

$d = 3 \sin \alpha = 3 \times \frac{8}{9} = \frac{8}{3}$



7. The matrix M is given by

$$M = \begin{pmatrix} k & -1 & 1 \\ 1 & 0 & -1 \\ 3 & -2 & 1 \end{pmatrix}, \quad k \neq 1$$

(a) Show that $\det M = 2 - 2k$.

(2)

(b) Find M^{-1} , in terms of k .

(5)

The straight line l_1 is mapped onto the straight line l_2 by the transformation represented

by the matrix $\begin{pmatrix} 2 & -1 & 1 \\ 1 & 0 & -1 \\ 3 & -2 & 1 \end{pmatrix}$.

The equation of l_2 is $(r-a) \times b = 0$, where $a = 4i + j + 7k$ and $b = 4i + j + 3k$.

(c) Find a vector equation for the line l_1 .

(5)

~~7(a) $\det(M) = k(0-2) + (1+3) + (-2)$~~

~~$= -2k + 4 - 2 = 2 - 2k$~~

as required

7(a) $\det(M) = k \begin{vmatrix} 0 & -1 \\ -2 & 1 \end{vmatrix} - (-1) \begin{vmatrix} 1 & -1 \\ 3 & 1 \end{vmatrix}$

$+ 1 \begin{vmatrix} 1 & 0 \\ 3 & -2 \end{vmatrix}$

$= k(0-2) + (1+3) + 1(-2)$

$= -2k + 4 - 2 = 2 - 2k$

Question 7 continued

$$(b) C = \begin{pmatrix} +(-2) & -(4) & +(-2) \\ -(1) & +(k-3) & -(3-2k) \\ +1 & -(-k-1) & +(1) \end{pmatrix}$$

$$= \begin{pmatrix} -2 & -4 & -2 \\ -1 & k-3 & 2k-3 \\ 1 & k+1 & 1 \end{pmatrix}$$

$$\therefore C^T = \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$$

$$\therefore M^{-1} = \frac{1}{2-2k} \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$$

$$(c) k=2$$

$$M l_1 = l_2 \quad M^{-1} = \frac{-1}{2} \begin{pmatrix} -2 & -1 & 1 \\ -4 & -1 & 3 \\ -2 & 1 & 1 \end{pmatrix}$$

$$\cancel{M} l_1 = M^{-1} l_2$$

$$= \frac{1}{2} \begin{pmatrix} 2 & 1 & -1 \\ 4 & 1 & -3 \\ 2 & -1 & -1 \end{pmatrix}$$



$$L \times \frac{1}{2} = a \times \frac{1}{2}$$



$$L = \begin{pmatrix} 4 \\ 1 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 + 4\lambda \\ 1 + \lambda \\ 7 + 3\lambda \end{pmatrix}$$

$$\therefore L_1 = \frac{1}{2} \begin{pmatrix} 2 & 1 & -1 \\ 4 & 1 & -3 \\ 2 & -1 & -1 \end{pmatrix} \begin{pmatrix} 4 + 4\lambda \\ 1 + \lambda \\ 7 + 3\lambda \end{pmatrix}$$

(3x3) (3x1)

$$= \frac{1}{2} \begin{pmatrix} 8 + 8\lambda + 1 + \lambda - 7 - 3\lambda \\ 16 + 16\lambda + 1 + \lambda - 21 - 9\lambda \\ 8 + 8\lambda - 1 - \lambda - 7 - 3\lambda \end{pmatrix}$$

$$= \frac{1}{2} \begin{pmatrix} 2 + 6\lambda \\ 8\lambda - 4 \\ 4\lambda \end{pmatrix} = \begin{pmatrix} 1 + 3\lambda \\ 4\lambda - 2 \\ 2\lambda \end{pmatrix}$$

$$\therefore L = \begin{pmatrix} 1 \\ -2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ 4 \\ 2 \end{pmatrix}$$

8. The hyperbola H has equation

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

- (a) Use calculus to show that the equation of the tangent to H at the point $(a \cosh \theta, b \sinh \theta)$ may be written in the form

$$xb \cosh \theta - ya \sinh \theta = ab \quad (4)$$

The line l_1 is the tangent to H at the point $(a \cosh \theta, b \sinh \theta)$, $\theta \neq 0$.
Given that l_1 meets the x -axis at the point P ,

- (b) find, in terms of a and θ , the coordinates of P . (2)

The line l_2 is the tangent to H at the point $(a, 0)$.
Given that l_1 and l_2 meet at the point Q ,

- (c) find, in terms of a , b and θ , the coordinates of Q . (2)

- (d) Show that, as θ varies, the locus of the mid-point of PQ has equation

$$x(4y^2 + b^2) = ab^2 \quad (6)$$

8(a). $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ differentiate:

$$\therefore \frac{2}{a^2}x - \frac{2}{b^2}y \frac{dy}{dx} = 0$$

$$\therefore \frac{2}{a^2}x = \frac{2}{b^2}y \frac{dy}{dx}$$

$$\therefore \frac{b^2}{a^2} \frac{x}{y} = \frac{dy}{dx}$$

$$\therefore \left(\frac{dy}{dx} \right) \frac{a \cosh \theta}{b \sinh \theta} = \frac{b^2}{a^2} \cdot \frac{a \cosh \theta}{b \sinh \theta}$$



Question 8 continued

$$= \frac{b}{a} \coth \theta = \left(\frac{\partial y}{\partial x} \right) \frac{a \cosh \theta}{b \sinh \theta}$$

$$\therefore y - b \sinh \theta = \frac{b}{a} \coth \theta (x - a \cosh \theta)$$

$$\therefore y - b \sinh \theta = \frac{b}{a} \coth \theta x - b \frac{\cosh^2 \theta}{\sinh \theta}$$

$$x(a \sinh \theta): \quad y a \sinh \theta = a b \sinh^2 \theta = x b \cosh \theta - a b \cosh^2 \theta$$

$$\therefore \cancel{y a \sinh \theta} - x b \cosh \theta$$

$$\therefore x b \cosh \theta - y a \sinh \theta = a b \cosh^2 \theta - a b \sinh^2 \theta$$

$$\therefore \cancel{x b \cosh \theta} - y a \sinh \theta = a b (\cosh^2 \theta - \sinh^2 \theta)$$

$$\therefore x b \cosh \theta - y a \sinh \theta = a b$$

$$\textcircled{b} \text{ at } l, \quad y=0 \Rightarrow b \cosh \theta = a$$


$$(b) \quad y=0 \Rightarrow x b \cosh \theta = ab$$

$$\Rightarrow x = \frac{a}{\cosh \theta}$$

$$\therefore P \left(\frac{a}{\cosh \theta}, 0 \right)$$

(c)

$$\frac{a^2}{a^2} + \frac{y^2}{b^2} = 1$$



l_2 has eqn: $x = a$

$$x = a \Rightarrow ab \cosh \theta - y \sinh \theta = ab$$

$$\therefore ab(\cosh \theta - 1) = y \sinh \theta$$

$$\therefore y = \frac{b(\cosh \theta - 1)}{\sinh \theta}$$

$$\therefore Q \left(a, \frac{b(\cosh \theta - 1)}{\sinh \theta} \right)$$

(d) Let midpoint = M

$$M: \left(\frac{a + \frac{a}{\cosh \theta}}{2}, \frac{b(\cosh \theta - 1)}{2 \sinh \theta} \right)$$

$$\therefore X = \frac{a \cosh \theta + a}{2 \cosh \theta} \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$$

$$\text{LHS} = x(4y^2 + b^2) = \frac{a \cosh \theta + a}{2 \cosh \theta} \left(4 \cdot \frac{b^2(\cosh \theta - 1)^2}{4 \sinh^2 \theta} + b^2 \right)$$

$$= \frac{a(a \cosh \theta + a)}{2 \cosh \theta} \left(\frac{b^2(\cosh \theta - 1)^2}{\sinh^2 \theta} + b^2 \right)$$

$$= \frac{a b^2 (\cosh \theta + 1)(\cosh \theta - 1)^2}{2 \cosh \theta \sinh^2 \theta} + \frac{a \cosh \theta + a}{2 \cosh \theta} b^2$$

$$\cosh^2 \theta - 1 = \sinh^2 \theta$$

$$= \frac{a b^2 (\cancel{\cosh^2 \theta - 1})(\cosh \theta - 1)}{2 \cosh \theta \sinh^2 \theta} + b^2 \frac{a \cosh \theta + a}{2 \cosh \theta}$$

$$= \frac{a b^2 (\cosh \theta - 1) + b^2 a \cosh \theta + b^2 a}{2 \cosh \theta}$$



$$= \frac{2ab^2 \cosh \theta}{2 \cosh \theta}$$

$$= ab^2 = \text{RHS}$$

// as required

$$\Rightarrow x(4y^2 + b^2) = ab^2$$

//