



**ADVANCED GCE UNIT
MATHEMATICS**

4727/01

Further Pure Mathematics 3
THURSDAY 25 JANUARY 2007

Morning

Time: 1 hour 30 minutes

Additional Materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.

ADVICE TO CANDIDATES

- Read each question carefully and make sure you know what you have to do before starting your answer.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **4** printed pages.

- 1 (i) Show that the set of numbers $\{3, 5, 7\}$, under multiplication modulo 8, does not form a group. [2]
- (ii) The set of numbers $\{3, 5, 7, a\}$, under multiplication modulo 8, forms a group. Write down the value of a . [1]
- (iii) State, justifying your answer, whether or not the group in part (ii) is isomorphic to the multiplicative group $\{e, r, r^2, r^3\}$, where e is the identity and $r^4 = e$. [2]

- 2 Find the equation of the line of intersection of the planes with equations

$$\mathbf{r} \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 4 \quad \text{and} \quad \mathbf{r} \cdot (\mathbf{i} + 5\mathbf{j} + 4\mathbf{k}) = 6,$$

giving your answer in the form $\mathbf{r} = \mathbf{a} + t\mathbf{b}$. [5]

- 3 (i) Solve the equation $z^2 - 6z + 36 = 0$, and give your answers in the form $r(\cos \theta \pm i \sin \theta)$, where $r > 0$ and $0 \leq \theta \leq \pi$. [4]
- (ii) Given that Z is either of the roots found in part (i), deduce the exact value of Z^{-3} . [3]

- 4 The variables x and y are related by the differential equation

$$\frac{dy}{dx} = \frac{x^2 - y^2}{xy}. \quad (\text{A})$$

- (i) Use the substitution $y = xz$, where z is a function of x , to obtain the differential equation

$$x \frac{dz}{dx} = \frac{1 - 2z^2}{z}. \quad [3]$$

- (ii) Hence show by integration that the general solution of the differential equation (A) may be expressed in the form $x^2(x^2 - 2y^2) = k$, where k is a constant. [6]

- 5 A multiplicative group G of order 9 has distinct elements p and q , both of which have order 3. The group is commutative, the identity element is e , and it is given that $q \neq p^2$.

- (i) Write down the elements of a proper subgroup of G

(a) which does not contain q , [1]

(b) which does not contain p . [1]

- (ii) Find the order of each of the elements pq and pq^2 , justifying your answers. [3]

(iii) State the possible order(s) of proper subgroups of G . [1]

(iv) Find two proper subgroups of G which are distinct from those in part (i), simplifying the elements. [4]

6 The variables x and y satisfy the differential equation

$$\frac{dy}{dx} + 3y = 2x + 1.$$

Find

(i) the complementary function, [1]

(ii) the general solution. [5]

In a particular case, it is given that $\frac{dy}{dx} = 0$ when $x = 0$.

(iii) Find the solution of the differential equation in this case. [3]

(iv) Write down the function to which y approximates when x is large and positive. [1]

7 The position vectors of the points A, B, C, D, G are given by

$$\mathbf{a} = 6\mathbf{i} + 4\mathbf{j} + 8\mathbf{k}, \quad \mathbf{b} = 2\mathbf{i} + \mathbf{j} + 3\mathbf{k}, \quad \mathbf{c} = \mathbf{i} + 5\mathbf{j} + 4\mathbf{k}, \quad \mathbf{d} = 3\mathbf{i} + 6\mathbf{j} + 5\mathbf{k}, \quad \mathbf{g} = 3\mathbf{i} + 4\mathbf{j} + 5\mathbf{k}$$

respectively.

(i) The line through A and G meets the plane BCD at M . Write down the vector equation of the line through A and G and hence show that the position vector of M is $2\mathbf{i} + 4\mathbf{j} + 4\mathbf{k}$. [6]

(ii) Find the value of the ratio $AG : AM$. [1]

(iii) Find the position vector of the point P on the line through C and G , such that $\overrightarrow{CP} = \frac{4}{3}\overrightarrow{CG}$. [2]

(iv) Verify that P lies in the plane ABD . [4]

8 (i) Use de Moivre's theorem to find an expression for $\tan 4\theta$ in terms of $\tan \theta$. [4]

(ii) Deduce that $\cot 4\theta = \frac{\cot^4 \theta - 6 \cot^2 \theta + 1}{4 \cot^3 \theta - 4 \cot \theta}$. [1]

(iii) Hence show that one of the roots of the equation $x^2 - 6x + 1 = 0$ is $\cot^2(\frac{1}{8}\pi)$. [3]

(iv) Hence find the value of $\operatorname{cosec}^2(\frac{1}{8}\pi) + \operatorname{cosec}^2(\frac{3}{8}\pi)$, justifying your answer. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.