

# 4727 Further Pure Mathematics 3

<p><b>1 (a) (i)</b> e.g. <math>ap \neq pa \Rightarrow</math> not commutative</p>	<p>B1 <b>1</b></p>	<p>For correct reason and conclusion</p>
<p><b>(ii)</b> 3</p>	<p>B1 <b>1</b></p>	<p>For correct number</p>
<p><b>(iii)</b> <math>e, a, b</math></p>	<p>B1 <b>1</b></p>	<p>For correct elements</p>
<p><b>(b)</b> <math>c^3</math> has order 2  <math>c^4</math> has order 3  <math>c^5</math> has order 6</p>	<p>B1                      B1                      B1 <b>3</b>  <span style="border: 1px solid black; padding: 2px;">6</span></p>	<p>For correct order                      For correct order                      For correct order</p>
<p><b>2</b> <math>m^2 - 8m + 16 = 0</math>  <math>\Rightarrow m = 4</math>  <math>\Rightarrow</math> CF (<math>y =</math>) <math>(A + Bx)e^{4x}</math>                      For PI try <math>y = px + q</math>  <math>\Rightarrow -8p + 16(px + q) = 4x</math>  <math>\Rightarrow p = \frac{1}{4} \quad q = \frac{1}{8}</math>  <math>\Rightarrow</math> GS <math>y = (A + Bx)e^{4x} + \frac{1}{4}x + \frac{1}{8}</math></p>	<p>M1                      A1                      A1√                      M1                        A1 A1                      B1√ <b>7</b>  <span style="border: 1px solid black; padding: 2px;">7</span></p>	<p>For stating and attempting to solve auxiliary eqn                      For correct solution                      For CF of correct form. f.t. from <math>m</math>                      For using linear expression for PI                        For correct coefficients                      For GS = CF + PI. Requires <span style="border: 1px solid black; padding: 2px;">y =</span>. f.t. from CF and PI with 2 arbitrary constants in CF and none in PI</p>
<p><b>3 (i)</b> line segment <math>OA</math></p>	<p>B1                      B1 <b>2</b></p>	<p>For stating line through <math>O</math> OR <math>A</math>                      For correct description <b>AEF</b></p>
<p><b>(ii)</b> <math>(\mathbf{r} - \mathbf{a}) \times (\mathbf{r} - \mathbf{b}) = \vec{AP} \times \vec{BP}</math>  <math>=  \mathbf{AP}   \mathbf{BP}  \sin \pi \cdot \hat{\mathbf{n}} = \mathbf{0}</math></p>	<p>B1                        B1 <b>2</b></p>	<p>For identifying <math>\mathbf{r} - \mathbf{a}</math> with <math>\vec{AP}</math> and <math>\mathbf{r} - \mathbf{b}</math> with <math>\vec{BP}</math>                      Allow direction errors                      For using <math>\times</math> of 2 parallel vectors = <math>\mathbf{0}</math>                      OR <math>\sin \pi = 0</math> or <math>\sin 0 = 0</math>                      in an appropriate vector expression</p>
<p><b>(iii)</b> line through <math>O</math>                      parallel to <math>AB</math></p>	<p>B1                      B1                      B1 <b>3</b>    <span style="border: 1px solid black; padding: 2px;">7</span></p>	<p>For stating line                      For stating through <math>O</math>                      For stating correct direction    <b>SR</b> For <math>\vec{AB}</math> or <math>\vec{BA}</math> allow B1 B0 B1</p>
<p><b>4</b> <math>(C + iS =) \int_0^{\frac{1}{2}\pi} e^{2x} (\cos 3x + i \sin 3x) (dx)</math>  <math>\cos 3x + i \sin 3x = e^{3ix}</math>  <math>\int_0^{\frac{1}{2}\pi} e^{(2+3i)x} (dx) = \frac{1}{2+3i} \left[ e^{(2+3i)x} \right]_0^{\frac{1}{2}\pi}</math>  <math>= \frac{2-3i}{4+9} \left( e^{(2+3i)\frac{1}{2}\pi} - e^0 \right) = \frac{2-3i}{13} (-ie^\pi - 1)</math>  <math>= \left\{ \frac{1}{13} (-2 - 3e^\pi + i(3 - 2e^\pi)) \right\}</math>  <math>C = -\frac{1}{13} (2 + 3e^\pi)</math>  <math>S = \frac{1}{13} (3 - 2e^\pi)</math></p>	<p>B1                      M1*                      A1                      A1                      M1                      (dep*)                      M1                      (dep*)                      A1                      A1    <span style="border: 1px solid black; padding: 2px;">8</span></p>	<p>For using de Moivre, seen or implied                      For writing as a single integral in exp form                      For correct integration (ignore limits)                      For substituting limits correctly (unsimplified)                      (may be earned at any stage)                      For multiplying by complex conjugate of <math>2+3i</math>                      For equating real and/or imaginary parts                      For correct expression <b>AG</b>                      For correct expression</p>

<p>5 (i) IF <math>e^{\int \frac{1}{x} dx} = e^{\ln x} = x</math>  OR <math>x \frac{dy}{dx} + y = x \sin 2x</math>  <math>\Rightarrow \frac{d}{dx}(xy) = x \sin 2x</math>  <math>\Rightarrow xy = \int x \sin 2x (dx)</math>  <math>xy = -\frac{1}{2} x \cos 2x + \frac{1}{2} \int \cos 2x (dx)</math>  <math>xy = -\frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x (+c)</math>  <math>\Rightarrow y = -\frac{1}{2} \cos 2x + \frac{1}{4x} \sin 2x + \frac{c}{x}</math></p>	<p>M1 A1 M1 A1 M1 A1 6</p>	<p>For correct process for finding integrating factor  OR for multiplying equation through by <math>x</math>  For writing DE in this form (may be implied)  For integration by parts the correct way round  For 1st term correct  For their 1st term and attempt at integration of <math>\frac{\cos}{\sin} kx</math>  For correct expression for <math>y</math></p>
<p>(ii) <math>(\frac{1}{4}\pi, \frac{2}{\pi}) \Rightarrow \frac{2}{\pi} = \frac{1}{\pi} + \frac{4c}{\pi} \Rightarrow c = \frac{1}{4}</math>  <math>\Rightarrow y = -\frac{1}{2} \cos 2x + \frac{1}{4x} \sin 2x + \frac{1}{4x}</math></p>	<p>M1 A1 2</p>	<p>For substituting <math>(\frac{1}{4}\pi, \frac{2}{\pi})</math> in solution  For correct solution. Requires <math>\boxed{y =}</math>.</p>
<p>(iii) <math>(y \approx) -\frac{1}{2} \cos 2x</math></p>	<p>B1√ 1 <b>9</b></p>	<p>For correct function <b>AEF</b> f.t. from (ii)</p>
<p>6 (i)  METHOD 1  State <math>B = (-1, -7, 2) + t(1, 2, -2)</math>  On plane <math>\Rightarrow (-1+t) + 2(-7+2t) - 2(2-2t) = -1</math>  <math>\Rightarrow t = 2 \Rightarrow B = (1, -3, -2)</math>  <math>AB = \sqrt{2^2 + 4^2 + 4^2}</math> OR <math>2\sqrt{1^2 + 2^2 + 2^2} = 6</math></p>	<p>M1 M1 M1 A1 A1 5</p>	<p><b>Either coordinates or vectors may be used</b>  Methods 1 and 2 may be combined, for a maximum of 5 marks  For using vector normal to plane  For substituting parametric form into plane  For solving a linear equation in <math>t</math>  For correct coordinates  For correct length of <math>AB</math></p>
<p>METHOD 2  <math>AB = \frac{ -1-14-4+1 }{\sqrt{1^2+2^2+2^2}} = 6</math>  OR <math>AB = \mathbf{AC} \cdot \frac{\mathbf{AB}}{\sqrt{1^2+2^2+2^2}} = \frac{[6, 7, 1] \cdot [1, 2, -2]}{\sqrt{1^2+2^2+2^2}} = 6</math>  <math>B = (-1, -7, 2) \pm 6 \frac{(1, 2, -2)}{\sqrt{1^2+2^2+2^2}}</math>  <math>B = (-1, -7, 2) \pm (2, 4, -4)</math>  <math>B = (1, -3, -2)</math></p>	<p>M1 A1 M1 B1 A1</p>	<p>For using a correct distance formula  For correct length of <math>AB</math>  For using <math>B = A + \text{length of } AB \times \text{unit normal}</math>  For checking whether + or - is needed (substitute into plane equation)  For correct coordinates (allow even if B0)</p>
<p>(ii) Find vector product of any two of <math>\pm[6, 7, 1], \pm[6, -3, 0], \pm(0, 10, 1)</math>  Obtain <math>k[1, 2, -20]</math>  <math>\theta = \cos^{-1} \frac{ [1, 2, -2] \cdot [1, 2, -20] }{\sqrt{1^2+2^2+2^2} \sqrt{1^2+2^2+20^2}}</math>  <math>\theta = \cos^{-1} \frac{45}{\sqrt{9} \sqrt{405}} = 41.8^\circ (41.810\dots^\circ, 0.72972\dots)</math></p>	<p>M1 A1 M1* M1 (dep*) A1√ A1 6 <b>11</b></p>	<p>For finding vector product of two relevant vectors  For correct vector <math>\mathbf{n}</math>  For using scalar product of two normal vectors  For stating both moduli in denominator  For correct scalar product. f.t. from <math>\mathbf{n}</math>  For correct angle</p>



<p><b>8 (i)</b> Group A: <math>e = 6</math>                  Group B: <math>e = 1</math>                  Group C: <math>e = 2^0</math> OR 1                  Group D: <math>e = 1</math></p>	<p>B1                  B1  <b>2</b></p>	<p>For any two correct identities                  For two other correct identities  <b>AEF</b> for D, but not “<math>m = n</math>”</p>
<p><b>(ii)</b> EITHER OR</p> <p>A   2 4 6 8                  2   4 8 2 6 orders of elements                  4 8 6 4 2 1, 2, 4, 4                  6 2 4 6 8 OR cyclic group                  8   6 2 8 4</p> <p>B   1 5 7 11                  1   1 5 7 11 orders of elements                  5 5 1 11 7 1, 2, 2, 2                  7 7 11 1 5 OR non-cyclic group                  11 11 7 5 1 OR Klein group</p> <p>C   <math>2^0</math> <math>2^1</math> <math>2^2</math> <math>2^3</math>  <math>2^0</math>   <math>2^0</math> <math>2^1</math> <math>2^2</math> <math>2^3</math> orders of elements  <math>2^1</math>   <math>2^1</math> <math>2^2</math> <math>2^3</math> <math>2^0</math> 1, 2, 4, 4  <math>2^2</math>   <math>2^2</math> <math>2^3</math> <math>2^0</math> <math>2^1</math> OR cyclic group  <math>2^3</math>   <math>2^3</math> <math>2^0</math> <math>2^1</math> <math>2^2</math></p> <p><math>A \not\cong B</math>  <math>B \not\cong C</math>  <math>A \cong C</math></p>	<p>B1*                  B1*                    B1                  (dep*)                  B1                  (dep*)                  B1                  (dep*)  <b>5</b></p>	<p>For showing group table                  OR sufficient details of orders of elements                  OR stating cyclic / non-cyclic / Klein group                  (as appropriate)</p> <p>for one of groups A, B, C                  for another of groups A, B, C</p> <p>For stating non-isomorphic } with sufficient detail                  For stating non-isomorphic } relating to the first 2 marks                  For stating isomorphic }</p>
<p><b>(iii)</b> <math>\frac{1+2m}{1+2n} \times \frac{1+2p}{1+2q} = \frac{1+2m+2p+4mp}{1+2n+2q+4nq}</math></p> <p><math>= \frac{1+2(m+p+2mp)}{1+2(n+q+2nq)} \equiv \frac{1+2r}{1+2s}</math></p>	<p>M1*                  M1                  (dep*)                  A1                  A1 <b>4</b></p>	<p>For considering product of 2 distinct elements of this form                  For multiplying out                  For simplifying to form shown                  For identifying as correct form, so closed</p> <p><b>SR</b> <math>\frac{\text{odd}}{\text{odd}} \times \frac{\text{odd}}{\text{odd}} = \frac{\text{odd}}{\text{odd}}</math> earns full credit  <b>SR</b> If clearly attempting to prove commutativity, allow at most M1</p>
<p><b>(iv)</b> Closure not satisfied                  Identity and inverse not satisfied</p>	<p>B1                  B1 <b>2</b>    <b>13</b></p>	<p>For stating closure                  For stating identity and inverse  <b>SR</b> If associativity is stated as not satisfied, then award at most B1 B0 OR B0 B1</p>