

OCR Further Maths AS-level

Additional Pure

Formula Sheet

Provided in formula book

Not provided in formula book

This work by [PMT Education](https://www.pmt.education) is licensed under [CC BY-NC-ND 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)



Sequences and Series

Behaviour of Sequences

Periodic	Terms of the sequence repeat regularly. The number of repeated terms is called the period.	$S = \{u_1, u_2, u_3, \dots, u_{n-1}, u_n, u_1, u_2, \dots\}$ Periodic with period n
Oscillating	Periodic with two terms.	$S = \{u_1, u_2, u_1, u_2, \dots\}$
Convergent	Terms of the sequence get closer to a limiting value.	$S = (u_n)$ $\lim_{n \rightarrow \infty} u_n = k$
Divergent	Sequence is not convergent, and the sum of the sequence is not finite.	$S = (u_n)$ $\lim_{n \rightarrow \infty} u_n$ does not exist $\sum_n u_n$ is undefined
Monotonic Increasing (or Decreasing)	Each term in the sequence is greater/less than or equal to the previous term	$S = (u_n)$ $u_n \geq u_{n-1}$ – monotonic increasing $u_n \leq u_{n-1}$ – monotonic decreasing

Fibonacci and Related Numbers

Fibonacci Recurrence Relation	$u_{n+2} = u_{n+1} + u_n, u_1 = 1, u_2 = 1$ Begins 1, 1, 2, 3, 5, 8, ...
Golden Ratio	Golden Ratio $\phi = \frac{1+\sqrt{5}}{2}$ is the limit of the ratio of consecutive terms in the Fibonacci sequence.
Lucas Recurrence Relation	$u_{n+2} = u_{n+1} + u_n, u_1 = 1, u_2 = 3$ Begins 1, 3, 4, 7, 11, 18, ...



Solving Recurrence Relations

1 st order linear recurrence relations with constant coefficients	$u_{n+1} = ku_n + f(n)$
Homogeneous 1 st order linear recurrence relation	$f(n) = 0$ so, of the form $u_{n+1} = ku_n$
Complementary function	Solution to homogenous version of the recurrence relation. 1 st order linear will have the form $u_n = A \times r^n$
Particular solution	Any solution of the recurrence relation.
General solution	Sum of the complementary function and the particular solution.
Recurrence system	Consists of a recurrence relation, initial conditions, and the range of the variable n .

Number Theory

Divisibility Tests

Divisible by 2	Last digit divisible by 2.
Divisible by 3	Sum of digits divisible by 3.
Divisible by 4	Number formed by final 2 digits divisible by 4.
Divisible by 5	Final digit is 0 or 5.
Divisible by 8	Number formed by final 3 digits divisible by 8.
Divisible by 9	Sum of digits divisible by 9.
Divisible by 11	Result of adding and subtracting digits in alternating order beginning at leftmost digit is divisible by 11.

Division Algorithm

If a is divided by b , where $0 < b < a$, then there is a unique quotient q and residue/remainder r (with $r < b$) such that $a = bq + r$. If $r = 0$, then $b|a$.



Finite (Modular) Arithmetic

If $a = nq + r$ then $a \equiv r \pmod{n}$

Rules	If $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$ then:
$a + c \equiv b + d \pmod{n}$	$a - c \equiv b - d \pmod{n}$
$ac \equiv bd \pmod{n}$	$a^k \equiv b^k \pmod{n}$

Linear Congruences

Linear congruence	Equation of the form $ax \equiv b \pmod{n}$.
Condition for a solution	$d b$ where d is the highest common factor of a and n . So if n is prime then $ax \equiv b \pmod{n}$ will have a solution as $\text{hcf}(a, n) = 1$ and $1 b$ for all integers b .
Solutions	$x_1 + \frac{n}{d} \times r$ where x_1 is a solution found by inspection and $r = 0, 1, \dots, d - 1$.

Prime Numbers

Prime number	An integer greater than 1 with no divisors other than 1 and itself.
Composite number	An integer with at least one divisor other than 1 and itself.
Coprime (relatively prime)	Two or more integers are coprime if 1 is their only common factor.
Fundamental theorem of arithmetic	Every integer greater than 1 is either prime or the unique product of primes (ignoring rearrangements).



Useful results	For integers a, b, c :
If a and b are coprime and $a c$ and $b c$, then $ab c$.	If $a b$ and $c d$, then $ac bd$.
If $a b$ and $b c$, then $a c$.	If $a b$ and $a c$, then $a (bx + cy)$ where x, y are integers.
$hcf(a, b)$ can be found by finding the smallest integer that can be written as $bx + cy$.	If $hcf(a, b) = 1$, then a and b are coprime.

Euclid's Lemma

Euclid's Lemma	If a prime number p divides into the composite number $a_1 \times a_2 \times \dots \times a_n$ then p must divide into at least one of a_1 to a_n .
Result from Euclid's Lemma	If $a bc$, where a and b are coprime, then $a c$.



Groups

Binary operations

Binary operation	A process involving two members of a set.
Definitions	Consider elements a and b of a set S .
Closed	A set is closed under an operation $*$ if for all $a, b \in S$, $a * b \in S$.
Commutative	The operation $*$ is commutative if for all $a, b \in S$, $a * b = b * a$.
Associative	The operation $*$ is associative if for all $a, b \in S$, $(a * b) * c = a * (b * c)$.
Identity element e for the operation $*$	$e \in S$ satisfies: $a * e = e * a = a$ for all elements $a \in S$.
Inverse a^{-1} for element a with operation $*$	$a^{-1} \in S$ satisfies: $a * a^{-1} = a^{-1} * a = e$ where e is the identity element.
Self-inverse a^{-1}	$a^{-1} \in S$ satisfies: $a^{-1} = a$ so $a^2 = e$ where e is the identity element.

Definition of a Group

Conditions for a set to be a group under operation $*$
Closed
Associative
The set contains an identity element e
Every element of the set has an inverse

Abelian Group	If all elements of the group commute under the binary operation.
---------------	--

Orders and elements of groups

Order of a group, $ G $	The number of elements the group contains.
Order of an element	The smallest power an element is raised to that gives the identity element.



Subgroups

Subgroup	H is a subgroup of the group G if H is a subset of G and H is also a group under the same binary operation.
Trivial subgroup	The trivial subgroup is $\{e\}$ where e is the group identity element.
Proper subgroup	A subgroup of G which is not G itself.

Cyclic groups

Cyclic groups	Every element of the group G is of the form a^n , where $a \in G$ and $n \in \mathbb{Z}$. a is called the generator of the group and is not necessarily unique.
---------------	--

Properties of Cyclic Groups

Commutative

At least one element of the group must have order n

Properties of groups

Order of group is 1	Group is $\{e\}$.
Order of group is 2,3,4,5, or 7	Group is cyclic.
Order of group is 4	Group is cyclic where: at least one element has order 4 or group is Klein group.
Order of group is 6	Group is cyclic if one element has order 6, otherwise group forms a symmetric group.



Further Vectors

Vector Product

$$\mathbf{a} \times \mathbf{b} = |\mathbf{a}||\mathbf{b}| \sin \theta \hat{\mathbf{n}},$$

where \mathbf{a} , \mathbf{b} , $\hat{\mathbf{n}}$ in that order, form a right-hand triple.

Observations	
Magnitude	$ \mathbf{a} \times \mathbf{b} = \mathbf{a} \mathbf{b} \sin \theta $
Condition for parallel or co-linear vectors	$\mathbf{a} \times \mathbf{b} = \mathbf{0}$ given that $\mathbf{a} \neq \mathbf{0}$ or $\mathbf{b} \neq \mathbf{0}$
Not commutative	$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$
Distributive over addition	$\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}$
Linear	$\mathbf{a} \times \lambda \mathbf{b} = \lambda \mathbf{a} \times \mathbf{b} = \lambda(\mathbf{a} \times \mathbf{b})$
Equation of a straight line	$(\mathbf{r} - \mathbf{a}) \times \mathbf{d} = \mathbf{0}$

Area of triangle with sides \mathbf{a} , \mathbf{b} .	$\frac{1}{2} \mathbf{a} \times \mathbf{b} $
Area of parallelogram with sides \mathbf{a} , \mathbf{b}	$ \mathbf{a} \times \mathbf{b} $

Surfaces and Partial Differentiation

Partial Differentiation

Mixed derivative theorem $f_{xy} = f_{yx}$ for suitably well-defined continuous functions f .

Stationary Points

Stationary points of a function $f(x, y)$ occur when $f_x = f_y = 0$. There are three types of stationary points: maximum, minimum or saddle.

