

## Linear Transformations

### Questions

**Q1.**

$$\mathbf{A} = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix}$$

(a) Describe fully the single geometrical transformation  $U$  represented by the matrix  $\mathbf{A}$ . (3)

The transformation  $V$ , represented by the  $2 \times 2$  matrix  $\mathbf{B}$ , is a reflection in the line  $y = -x$

(b) Write down the matrix  $\mathbf{B}$ . (1)

Given that  $U$  followed by  $V$  is the transformation  $T$ , which is represented by the matrix  $\mathbf{C}$ ,

(c) find the matrix  $\mathbf{C}$ . (2)

(d) Show that there is a real number  $k$  for which the point  $(1, k)$  is invariant under  $T$ . (4)

**(Total for question = 10 marks)**

**Q2.**

(i)

$$\mathbf{A} = \begin{pmatrix} 2 & a \\ a - 4 & b \end{pmatrix}$$

where  $a$  and  $b$  are non-zero constants.

Given that the matrix  $\mathbf{A}$  is self-inverse,

(a) determine the value of  $b$  and the possible values for  $a$ . (5)

The matrix  $\mathbf{A}$  represents a linear transformation  $M$ .

Using the smaller value of  $a$  from part (a),

(b) show that the invariant points of the linear transformation  $M$  form a line, stating the equation of this line. (3)

(ii)

$$\mathbf{P} = \begin{pmatrix} p & 2p \\ -1 & 3p \end{pmatrix}$$

where  $p$  is a positive constant.

The matrix  $\mathbf{P}$  represents a linear transformation  $U$ .

The triangle  $T$  has vertices at the points with coordinates  $(1, 2)$ ,  $(3, 2)$  and  $(2, 5)$ .

The area of the image of  $T$  under the linear transformation  $U$  is 15

(a) Determine the value of  $p$ .

(4)

The transformation  $V$  consists of a stretch scale factor 3 parallel to the  $x$ -axis with the  $y$ -axis invariant followed by a stretch scale factor  $-2$  parallel to the  $y$ -axis with the  $x$ -axis invariant. The transformation  $V$  is represented by the matrix  $\mathbf{Q}$ .

(b) Write down the matrix  $\mathbf{Q}$ .

(2)

Given that  $U$  followed by  $V$  is the transformation  $W$ , which is represented by the matrix  $\mathbf{R}$ ,

(c) find the matrix  $\mathbf{R}$ .

(2)

**(Total for question = 16 marks)**

**Q3.**

The transformation  $P$  is an enlargement, centre the origin, with scale factor  $k$ , where  $k > 0$

The transformation  $Q$  is a rotation through angle  $\theta$  degrees anticlockwise about the origin.

The transformation  $P$  followed by the transformation  $Q$  is represented by the matrix

$$\mathbf{M} = \begin{pmatrix} -4 & -4\sqrt{3} \\ 4\sqrt{3} & -4 \end{pmatrix}$$

(a) Determine

(i) the value of  $k$ ,

(ii) the smallest value of  $\theta$

(4)

A square  $S$  has vertices at the points with coordinates  $(0, 0)$ ,  $(a, -a)$ ,  $(2a, 0)$  and  $(a, a)$  where  $a$  is a constant.

The square  $S$  is transformed to the square  $S'$  by the transformation represented by  $\mathbf{M}$ .

(b) Determine, in terms of  $a$ , the area of  $S'$

(2)

**(Total for question = 6 marks)**

**Q4.**

$$\mathbf{M} = \begin{pmatrix} 4 & -5 \\ 2 & -7 \end{pmatrix}$$

(a) Show that the matrix  $\mathbf{M}$  is non-singular.

(2)

The transformation  $T$  of the plane is represented by the matrix  $\mathbf{M}$ .

The triangle  $R$  is transformed to the triangle  $S$  by the transformation  $T$ .

Given that the area of  $S$  is 63 square units,

(b) find the area of  $R$ .

(2)

(c) Show that the line  $y = 2x$  is invariant under the transformation  $T$ .

(2)

**(Total for question = 6 marks)**

**Q5.**

$$\mathbf{A} = \begin{pmatrix} 4 & -2 \\ 5 & 3 \end{pmatrix}$$

The matrix  $\mathbf{A}$  represents the linear transformation  $M$ .

Prove that, for the linear transformation  $M$ , there are no invariant lines.

(5)

**(Total for question = 5 marks)**

Q6.

$$\left[ \begin{array}{l} \text{With respect to the right-hand rule, a rotation through } \theta^\circ \text{ anticlockwise about the} \\ \text{y-axis is represented by the matrix} \\ \\ \begin{pmatrix} \cos\theta & 0 & \sin\theta \\ 0 & 1 & 0 \\ -\sin\theta & 0 & \cos\theta \end{pmatrix} \end{array} \right]$$

The point  $P$  has coordinates  $(8, 3, 2)$

The point  $Q$  is the image of  $P$  under the transformation reflection in the plane  $y = 0$

(a) Write down the coordinates of  $Q$

(1)

The point  $R$  is the image of  $P$  under the transformation rotation through  $120^\circ$  anticlockwise about the  $y$ -axis, with respect to the **right-hand rule**.

(b) Determine the exact coordinates of  $R$

(2)

(c) Hence find  $|\vec{PR}|$  giving your answer as a simplified surd.

(2)

(d) Show that  $\vec{PR}$  and  $\vec{PQ}$  are perpendicular.

(1)

(e) Hence determine the exact area of triangle  $PQR$ , giving your answer as a surd in simplest form.

(2)

**(Total for question = 8 marks)**

**Q7.**

$$\mathbf{P} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \mathbf{Q} = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$$

(a) (i) Describe fully the single geometrical transformation  $P$  represented by the matrix  $\mathbf{P}$ .

(ii) Describe fully the single geometrical transformation  $Q$  represented by the matrix  $\mathbf{Q}$ .

(4)

The transformation  $P$  followed by the transformation  $Q$  is the transformation  $R$ , which is represented by the matrix  $\mathbf{R}$ .

(b) Determine  $\mathbf{R}$ .

(1)

(c) (i) Evaluate the determinant of  $\mathbf{R}$ .

(ii) Explain how the value obtained in (c)(i) relates to the transformation  $R$ .

(2)

**(Total for question = 7 marks)**

**Q8.**

$$\mathbf{A} = \begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix}, \quad \mathbf{P} = \begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix}$$

(a) Find  $\mathbf{A}^{-1}$

(2)

The transformation represented by the matrix  $\mathbf{B}$  followed by the transformation represented by the matrix  $\mathbf{A}$  is equivalent to the transformation represented by the matrix  $\mathbf{P}$ .

(b) Find  $\mathbf{B}$ , giving your answer in its simplest form.

(3)

**(Total for question = 5 marks)**

**Q9.**

(i)

$$\mathbf{A} = \begin{pmatrix} p & 2 \\ 3 & p \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} -5 & 4 \\ 6 & -5 \end{pmatrix}$$

where  $p$  is a constant.

(a) Find, in terms of  $p$ , the matrix  $\mathbf{AB}$

(2)

Given that

$$\mathbf{AB} + 2\mathbf{A} = k\mathbf{I}$$

where  $k$  is a constant and  $\mathbf{I}$  is the  $2 \times 2$  identity matrix,

(b) find the value of  $p$  and the value of  $k$ .

(4)

(ii)

$$\mathbf{M} = \begin{pmatrix} a & -9 \\ 1 & 2 \end{pmatrix}, \quad \text{where } a \text{ is a real constant}$$

Triangle  $T$  has an area of 15 square units.

Triangle  $T$  is transformed to the triangle  $T'$  by the transformation represented by the matrix  $\mathbf{M}$ .

Given that the area of triangle  $T'$  is 270 square units, find the possible values of  $a$ .

(5)

**(Total for question = 11 marks)****Q10.**

$$\mathbf{M} = \begin{pmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix}$$

(a) Show that  $\mathbf{M}$  is non-singular.

(2)

The hexagon  $R$  is transformed to the hexagon  $S$  by the transformation represented by the matrix  $\mathbf{M}$ .

Given that the area of hexagon  $R$  is 5 square units,

(b) find the area of hexagon  $S$ .

(1)

The matrix  $\mathbf{M}$  represents an enlargement, with centre  $(0, 0)$  and scale factor  $k$ , where  $k > 0$ , followed by a rotation anti-clockwise through an angle  $\theta$  about  $(0, 0)$ .

(c) Find the value of  $k$ .

(2)

(d) Find the value of  $\theta$ .

(2)

**(Total for question = 7 marks)**

**Mark Scheme – Linear Transformations**

Q1.

Question	Scheme	Marks	AOs
(a)	Rotation	B1	1.1b
	120 degrees (anticlockwise) or $\frac{2\pi}{3}$ radians (anticlockwise) Or 240 degrees clockwise or $\frac{4\pi}{3}$ radians clockwise	B1	2.5
	About (from) the origin. Allow (0, 0) or <i>O</i> for origin.	B1	1.2
		(3)	
(b)	$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$	B1	1.1b
		(1)	
(c)	$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} \dots & \dots \\ \dots & \dots \end{pmatrix}$	M1	1.1b
	$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix}$	A1ft	1.1b
		(2)	
(d)	$\begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} 1 \\ k \end{pmatrix} = \begin{pmatrix} 1 \\ k \end{pmatrix} = \dots$ or $\begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} = \dots$	M1	3.1a
	Note: $\begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} 1 \\ k \end{pmatrix} = \begin{pmatrix} -\frac{\sqrt{3}}{2} + \frac{1}{2}k \\ \frac{1}{2} + \frac{\sqrt{3}}{2}k \end{pmatrix} = \begin{pmatrix} 1 \\ k \end{pmatrix}$ can score M1 (for the matrix equation) but needs an equation to be “extracted” to score the next A1		
	$-\frac{\sqrt{3}}{2} + \frac{1}{2}k = 1$ or $\frac{1}{2} + \frac{\sqrt{3}}{2}k = k$ or $x = -\frac{\sqrt{3}}{2}x + \frac{1}{2}y$ or $y = \frac{1}{2}x + \frac{\sqrt{3}}{2}y$ (Note that candidates may then substitute $x = 1$ which is acceptable)	A1ft	1.1b
	$-\frac{\sqrt{3}}{2} + \frac{1}{2}k = 1$ or $x = -\frac{\sqrt{3}}{2}x + \frac{1}{2}y \Rightarrow k = 2 + \sqrt{3} \left( \text{or } \frac{1}{2 - \sqrt{3}} \right)$	A1	1.1b
	$\frac{1}{2} + \frac{\sqrt{3}}{2}k = k$ or $y = \frac{1}{2}x + \frac{\sqrt{3}}{2}y \Rightarrow k = 2 + \sqrt{3} \left( \text{or } \frac{1}{2 - \sqrt{3}} \right)$	B1	1.1b
	(4)		

(10 marks)



Notes
<p>(a)            B1: Identifies the transformation as a rotation            B1: Correct angle. Allow equivalents in degrees or radians.            B1: Identifies the origin as the centre of rotation            These marks can only be awarded as the elements of a <b>single transformation</b></p> <p>(b)            B1: Shows the correct matrix in the correct form</p> <p>(c)            M1: Multiplies the matrices in the correct order (evidence of multiplication can be taken from 3 correct or 3 correct ft elements)            A1ft: Correct matrix (follow through their matrix from part (b))            A correct matrix or a correct follow through matrix implies both marks.</p> <p>(d)            M1: Translates the problem into a matrix multiplication to obtain at least one equation in <math>k</math> or in <math>x</math> and <math>y</math>            A1ft: Obtains one correct equation (follow through their matrix from part (c))            A1: Correct value for <math>k</math> in any form            B1: Checks their answer by independently solving both equations <b>correctly</b> to obtain <math>2+\sqrt{3}</math> both times or substitutes <math>2+\sqrt{3}</math> into the other equation to confirm its validity</p>

Q2.

Question	Scheme	Marks	AOs	
(i) (a)	Multiplies the matrix $A$ by itself and sets equal to $I$ to form one equation in $a$ only and another equation involving both $a$ and $b$ . $\begin{pmatrix} 2 & a \\ a-4 & b \end{pmatrix} \begin{pmatrix} 2 & a \\ a-4 & b \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \Rightarrow 4+a(a-4)=1$ and either $2a+ab=0$ or $2(a-4)+b(a-4)=0$ or $a(a-4)+b^2=1$	M1	3.1a	
	Solves a 3TQ involving only the constant $a$ . This could come after a value of $b$ is found and this value substituted into an equation involving both $a$ and $b$ $a^2-4a+3=0 \Rightarrow (a-3)(a-1)=0 \Rightarrow a=...$	dM1	1.1b	
	$a=1, a=3$	A1	1.1b	
	Substitutes a value for $a$ into an equation involving both $a$ and $b$ and solves for $b$ . e.g. $2(1)+(1)b \Rightarrow b=...$ $2(1-4)b+(1-4)=0 \Rightarrow b=...$ $(1)(1-4)+b^2=1 \Rightarrow b=...$	Alternatively uses $2a+ab=0$ $a(2+b)=0$ As $a \neq 0$ $2+b=0 \Rightarrow b=...$	dM1	1.1b
	$b=-2$	A1	1.1b	
		(5)		

	<p style="text-align: center;"><b>Alternative (i) (a)</b></p> <p>Finds <math>A^{-1}</math> in terms of <math>a</math> and <math>b</math>, sets equal to <math>A</math> and attempts to find at least two different equations. Allow a single sign slip</p> $\frac{1}{2b-a(a-4)} \begin{pmatrix} b & -a \\ -(a-4) & 2 \end{pmatrix} = \begin{pmatrix} 2 & a \\ a-4 & b \end{pmatrix}$ <p>One equation from <math>\frac{b}{2b-a(a-4)} = 2, \frac{2}{2b-a(a-4)} = b</math></p> <p>One equation from <math>\frac{-a}{2b-a(a-4)} = a, \frac{-(a-4)}{2b-a(a-4)} = a-4</math></p>	M1	3.1a	
	<p>Uses their value of <math>b</math> and their value of the determinant to form and solve a 3TQ involving only the constant <math>a</math></p> $a^2 - 4a + 3 = 0$ $\Rightarrow (a-3)(a-1) = 0$ $\Rightarrow a = \dots$	<p>Eliminates <math>b</math> from their equations and solve a 3TQ involving only the constant <math>a</math></p> $a^2 - 4a + 3 = 0$ $\Rightarrow (a-3)(a-1) = 0$ $\Rightarrow a = \dots$	dM1	1.1b
	$a = 1, a = 3$		A1	1.1b
	$\frac{-a}{2b-a(a-4)} = a$ $\Rightarrow 2b-a(a-4) = -1 \Rightarrow \frac{b}{-1} = 2$ <p style="text-align: center;">Or</p> $\frac{-(a-4)}{2b-a(a-4)} = a-4$ $\Rightarrow 2b-a(a-4) = -1$ $\Rightarrow \frac{2}{-1} = b$	<p>Substitutes a value for <math>a</math> into an equation to find a value for <math>b</math></p>	dM1	1.1b
	$b = -2$		A1	1.1b

(b)	Uses their smallest value of <b>a</b> and their value for <b>b</b> to form two equations $\begin{pmatrix} 2 & 'a' \\ 'a-4' & 'b' \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow 2x + ay = x \text{ and } (a-4)x + by = y$	M1	3.1a
	$\begin{pmatrix} 2 & 1 \\ -3 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow 2x + y = x \text{ and } -3x - 2y = y$		
	$2x + y = x \Rightarrow x + y = 0 \text{ o.e. and } -3x - 2y = y \Rightarrow x + y = 0 \text{ o.e.}$	M1	1.1b
	$x + y = 0 \text{ o.e.}$	A1	2.1
		(3)	
(ii)(a)	Area of the triangle $T = 3$	B1	1.1b
	Complete method to find a value for $p$ . Need to see an attempt at the determinant and setting equal to 15 divided by their area of $T$ . The resulting 3TQ needs to be solved to find a value of $p$ . Determinant $3p \times p - (-1) \times 2p = \frac{15}{\text{'their area'}} \Rightarrow p = \dots$	M1	3.1a
	$3p^2 + 2p - 5 (= 0)$	A1	1.1b
	$p = 1 \text{ must reject } p = -\frac{5}{3}$	A1	1.1b
		(4)	
(b)	$\begin{pmatrix} 3 & 0 \\ 0 & -2 \end{pmatrix}$	B1 B1	1.1b 1.1b
		(2)	
(c)	(their matrix found in part (b)) $\begin{pmatrix} 'p' & 2'p' \\ -1 & 3'p' \end{pmatrix} = \begin{pmatrix} \dots & \dots \\ \dots & \dots \end{pmatrix}$	M1	1.1b
	$\begin{pmatrix} 3 & 0 \\ 0 & -2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix} = \begin{pmatrix} \dots & \dots \\ \dots & \dots \end{pmatrix}$		
	$\begin{pmatrix} 3 & 6 \\ 2 & -6 \end{pmatrix}$	A1ft	1.1b
		(2)	
<b>(16 marks)</b>			

Notes:
<p><b>(i)(a)</b>  <b>M1:</b> Forming two equations, one involving <math>a</math> only and one involving <math>a</math> and <math>b</math>  <b>dM1:</b> Dependent on previous mark, solves a 3TQ involving <math>a</math>  <b>A1:</b> Correct values for <math>a</math>  <b>dM1:</b> Dependent on first method mark Substitutes one of their values of <math>a</math> into an equation involving <math>a</math> and <math>b</math> and solve to find a value for <math>b</math>. Alternatively factorises either <math>2a + ab = 0</math> and uses <math>a = 0</math> to find a value for <math>b</math>.  <b>A1:</b> Correct value for <math>b</math></p>
<p><b>Alternative(i)(a)</b>  <b>M1:</b> Finds <math>A^{-1}</math> and sets equal to <math>A</math> and forms two different equations  <b>dM1:</b> Dependent on previous mark. Eliminates <math>b</math> from their equations and solves a 3TQ involving only the constant <math>a</math>. Alternatively if the value of <math>b</math> is found first substitutes their value for <math>b</math> into their determinant <math>= -1</math> to form and solve a 3TQ for <math>a</math>  <b>A1:</b> Correct value for <math>a</math>  <b>dM1:</b> Dependent on first method mark. Substitutes a value for <math>a</math> into an equation to find a value for <math>b</math>. Alternatively uses one equation to find the determinant <math>= -1</math> and uses this to find a value of <math>b</math>.  <b>A1:</b> Correct values for <math>b</math></p>
<p><b>(b)</b>  <b>M1:</b> Extracts simultaneous equations using their matrix <math>A</math> with their smaller value of <math>a</math>.  <b>M1:</b> Gathers terms from their two equations.  <b>A1:</b> Achieves the correct equations and deduces the correct line. Accept equivalent equations as long as both have been shown to be the same.</p>
<p><b>(ii)(a)</b>  <b>B1:</b> Area of the triangle <math>T = 3</math>  <b>M1:</b> Full method. Finds the determinant, sets equal to <math>15</math>/their area and solves the resulting 3TQ  <b>A1:</b> Correct quadratic  <b>A1:</b> <math>p = 1</math> only</p>
<p><b>(b)</b>  <b>B1</b> One correct row or column  <b>B1:</b> All correct</p>
<p><b>(c)</b>  <b>M1:</b> Multiplies the matrices <math>QP</math> in the correct order (if answer only then evidence can be taken from 3 correct or 3 correct ft elements)  <b>A1ft:</b> Correct matrix following through on their answer to part (b) and their value of <math>p</math> as long as it is a positive constant</p>

Q3.

Question	Scheme	Marks	AOs	
(a) Way 1	$\det \mathbf{M} = -4 \times -4 - 4\sqrt{3} \times -4\sqrt{3} = \dots \Rightarrow k = \sqrt{\det \mathbf{M}} = \dots$	M1	3.1a	
	$k = 8$	A1	1.1b	
	$\Rightarrow \mathbf{Q} = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \Rightarrow \cos \theta = -\frac{1}{2} \Rightarrow \theta = \dots$	M1	1.1b	
	$(\cos \theta < 0, \sin \theta > 0 \Rightarrow \text{Quadrant 2 so}) \quad \theta = 120^\circ$	A1	1.1b	
		(4)		
	Way 2	$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix} = k \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} = \begin{pmatrix} -4 & -4\sqrt{3} \\ 4\sqrt{3} & -4 \end{pmatrix}$	M1	3.1a
		Achieves both the equations $k \cos \theta = -4$ and $k \sin \theta = 4\sqrt{3}$	A1	1.1b
$\frac{k \sin \theta}{k \cos \theta} = \frac{4\sqrt{3}}{-4} \Rightarrow \tan \theta = -\sqrt{3} \Rightarrow \theta = \dots$		M1	1.1b	
$\theta = 120^\circ$ and $k = 8$		A1	1.1b	
		(4)		
(b)	Area of $S' = \text{area of } S \times k^2$ (The area of the square $S = 2a^2$ )	M1	1.1b	
	Area of $S' = 128a^2$	A1ft	2.2a	
		(2)		

(6 marks)

**Notes:****(a) Way 1**

**M1:** A full method to find  $k$  such as attempting the square root of the determinant of  $\mathbf{M}$ . It is immediately deducible so the method may be implied by  $k = 8$ .

**A1:**  $k = 8$

**M1:** A full method to find a value of  $\theta$  using their  $k$ , no need to justify quadrant. Only one equation needed for this mark. Allow if a radians answer is given. May be implied by a correct angle.

**A1:** Correct angle in degrees.

**Way 2**

**M1:** Multiplies the correct matrix representing transformation  $Q$  by the matrix representing transformation  $P$  and sets equal to matrix  $\mathbf{M}$ . Allow for the matrices either way round as the transformations commute. No need to see the identity matrix, just multiplying through by  $k$  is sufficient.

**A1:** Both correct equations. Note that if a correct value of  $k$  is found, this A is scored under Way 1.

**M1:** Solves their simultaneous equations to find a value for  $\theta$  (or  $k$ )

**A1:**  $\theta = 120^\circ$  and  $k = 8$

**(b)**

**M1:** Complete method to find the area of  $S'$ : 'their  $k^2 \times$  their  $2a^2$ '. Must be an attempt at the area of  $S$  but it need not be correct.

**A1ft:** Deduces the correct area for  $S'$ , follow through their value of  $k$

Q4.

Question	Scheme	Marks	AOs
(a)	$(\det(\mathbf{M}) \Rightarrow) (4)(-7) - (2)(-5)$	M1	1.1a
	$\mathbf{M}$ is non-singular because $\det(\mathbf{M}) = -18$ and so $\det(\mathbf{M}) \neq 0$	A1	2.4
		(2)	
(b)	$\text{Area } R = \frac{\text{Area } S}{(\pm) \det \mathbf{M} } = \dots$	M1	1.2
	$\text{Area}(R) = \frac{63}{ -18 } = \frac{7}{2}$ oe	A1ft	1.1b
		(2)	
(c)	$\begin{pmatrix} 4 & -5 \\ 2 & -7 \end{pmatrix} \begin{pmatrix} x \\ 2x \end{pmatrix} = \begin{pmatrix} 4x - 10x \\ 2x - 14x \end{pmatrix}$	M1	1.1b
	$= \begin{pmatrix} -6x \\ -12x \end{pmatrix}$ and so all points on $y = 2x$ map to points on $y = 2x$ , hence the line is invariant. OR $= -6 \begin{pmatrix} x \\ 2x \end{pmatrix}$ hence $y = 2x$ is invariant.	A1	2.1
		(2)	
<b>(6 marks)</b>			

Notes			
(a)	M1	An attempt to find $\det(\mathbf{M})$ . Just the calculation is sufficient. Site of $-18$ implies this mark, which may be embedded in an attempt at the inverse..	
	A1	$\det(\mathbf{M}) = -18$ and reference to zero, e.g. $-18 \neq 0$ and conclusion. The conclusion may precede finding the determinant (e.g. "Non-singular if $\det(\mathbf{M}) \neq 0$ , $\det(\mathbf{M}) = -18 \neq 0$ " is sufficient or accept "Non-singular if $\det(\mathbf{M}) \neq 0$ , $\det(\mathbf{M}) = -18$ , therefore non-singular" or some other indication of conclusion.) Need not mention " $\det(\mathbf{M})$ " to gain both marks here, a correct calculation, statement $-18 \neq 0$ , and conclusion hence $\mathbf{M}$ is non-singular can gain M1A1.	
(b)	M1	Recalls determinant is needed for area scale factor by dividing 63 by $\pm$ their determinant.	
	A1ft	$\frac{7}{2}$ or follow through $\frac{63}{ \text{their det} }$ . Must be positive and should be simplified to single fraction or exact decimal. (Allow if made positive following division by a negative determinant.)	
(c)	M1	Attempts the matrix multiplication shown or with equivalent, e.g. $\begin{pmatrix} 1 \\ 2 \end{pmatrix} y$ . May use $\begin{pmatrix} x \\ y \end{pmatrix}$ and substitute $y = 2x$ later and this is fine for the method.	
	A1	Correct multiplication and working leading to conclusion that the line is invariant. If the $-6$ is not extracted, they must make reference to image points being on line $y = 2x$ . If the $-6$ is extracted to show it is a multiple of $\begin{pmatrix} x \\ 2x \end{pmatrix}$ followed by a conclusion "invariant" as minimum.	

<b>Alt for (c)</b>	$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{1}{-18} \begin{pmatrix} -7 & 5 \\ -2 & 4 \end{pmatrix} \begin{pmatrix} x \\ 2x \end{pmatrix} = \frac{-1}{18} \begin{pmatrix} -7x+10x \\ -2x+8x \end{pmatrix}$	<b>M1</b>	1.1b
	$= \frac{-1}{18} \begin{pmatrix} 3x \\ 6x \end{pmatrix} = \frac{-1}{6} \begin{pmatrix} x \\ 2x \end{pmatrix} \Rightarrow b = 2a$ so points on line $y = 2x$ map to points on $y = 2x$ , hence it is invariant.	<b>A1</b>	2.1
Marks as per main scheme.			
<b>Alt 2</b>	(Since linear transformations map straight lines to straight lines...) E.g. $(1, 2)$ is on line $y = 2x$ , and $\begin{pmatrix} 4 & -5 \\ 2 & -7 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 4-10 \\ 2-14 \end{pmatrix}$	<b>M1</b>	1.1b
	$= \begin{pmatrix} -6 \\ -12 \end{pmatrix}$ , which is also on the line $y=2x$ , hence as $(0,0)$ and $(1,2)$ both map to points on $y = 2x$ (and transformation is linear) then $y=2x$ is invariant.	<b>A1</b>	2.1
<b>Notes</b>			
<b>M1</b>	Identifies a point on the line $y = 2x$ and finds its image under $T$ . If $(0,0)$ is used there must be a clear statement it is because this is on the line, but for other points accept with any line on $y = 2x$ without statement.		
<b>A1</b>	Shows the image and another point, which may be $(0,0)$ , on $y=2x$ both map to points on $y = 2x$ concludes line is invariant. Need not reference transformation being linear for either mark here.		
<b>Alt 3</b>	$\begin{pmatrix} 4 & -5 \\ 2 & -7 \end{pmatrix} \begin{pmatrix} x \\ mx+c \end{pmatrix} = \begin{pmatrix} X \\ mX+c \end{pmatrix} \Rightarrow \begin{aligned} 4x-5(mx+c) &= X \\ 2x-7(mx+c) &= mX+c \end{aligned}$ $\Rightarrow 2x-7(mx+c) = m(4x-5(mx+c))+c$ $\Rightarrow (5m^2-11m+2)x+(5m-8)c=0$ $\Rightarrow (5m-1)(m-2)=0 \Rightarrow m=...$ Or similar work with $c = 0$ throughout.	<b>M1</b>	2.1
	$(5m-8 \neq 0 \Rightarrow c=0)$ Hence $m = 2$ gives an invariant line (with $c = 0$ ), so $y = 2x$ is invariant.	<b>A1</b>	1.1b
<b>Notes</b>			
<b>M1</b>	Attempts to find the equation of a general invariant line, or general invariant line through the origin (so may have $c = 0$ throughout). To gain the method mark they must progress from finding the simultaneous equations to forming a quadratic in $m$ and solving to a value of $m$ .		
<b>A1</b>	Correct quadratic in $m$ found, with $m = 2$ as solution (ignore the other) and deduction that hence $y = 2x$ is an invariant line. Ignore errors in the $(5m-8)$ here as $c = 0$ is always a possible solution. No need to see $c = 0$ derived.		

Q5.

Question	Scheme	Marks	AOs	
	$\begin{pmatrix} 4 & -2 \\ 5 & 3 \end{pmatrix} \begin{pmatrix} x \\ mx+c \end{pmatrix} = \begin{pmatrix} X \\ mX+c \end{pmatrix}$ leading to an equation in $x$ , $m$ , $c$ and $X$	M1	3.1a	
	$4x - 2(mx+c) = X$ and $5x + 3(mx+c) = mX+c$	A1	1.1b	
	$5x + 3(mx+c) = m(4x - 2(mx+c)) + c$ leading to $5 + 3m = 4m - 2m^2$ <span style="float: right;"><math>(3c = -2mc + c)</math></span>	M1	2.1	
	$2m^2 - m + 5 = 0 \Rightarrow b^2 - 4ac =$ $(-1)^2 - 4(2)(5) = \dots$	Solves $3c = -2mc + c \Rightarrow m = \dots$	dM1	1.1b
	Correct expression for the discriminant = $\{-39\} < 0$ therefore there are no invariant lines.	$m = -1$ and shows a contradiction in $5 + 3m = 4m - 2m^2$ therefore there are no invariant lines.	A1	2.4
<b>Alternative</b>				
	$\begin{pmatrix} 4 & -2 \\ 5 & 3 \end{pmatrix} \begin{pmatrix} x \\ mx \end{pmatrix} = \begin{pmatrix} X \\ mX \end{pmatrix}$ leading to an equation in $x$ , $m$ and $X$	M1	3.1a	
	$4x - 2(mx) = X$ and $5x + 3(mx) = mX$	A1	1.1b	
	$5x + 3(mx) = m(4x - 2(mx))$ leading to $5 + 3m = 4m - 2m^2$	M1	2.1	
	$2m^2 - m + 5 = 0 \Rightarrow b^2 - 4ac = (-1)^2 - 4(2)(5) = \dots$		dM1	1.1b
	Correct expression for the discriminant = $\{-39\} < 0$ therefore there are no invariant lines that pass through the origin no invariant lines.		A1	2.4
		(5)		
(5 marks)				

**Notes:**

**M1:** Sets up a matrix equation in an attempt to find a fixed line and extract at least one equation.

**A1:** Correct equations.

**M1:** Eliminates  $X$  from the simultaneous equations and equates the coefficients of  $x$  leading to a quadratic equation in terms of  $m$ .

**dM1:** Dependent on the previous method, finds the value of the discriminant, this can be seen in an attempt to solve the quadratic using the formula.

Alternatively solves  $3c = -2mc + c$  and finds a value for  $m$

**Note:** If the quadratic equation in  $m$  is solved on a calculator and complex roots given this is M0 as they are not showing why there are no real roots.

**A1:** Correct expression for the discriminant, states  $< 0$  and draws the required conclusion.

Alternatively, correct value for  $m$ , shows a contradiction in  $5 + 3m = 4m - 2m^2$  and draws the required conclusion.

**Alternative**

**M1:** Sets up a matrix equation in an attempt to find a fixed line and extract at least one equation.

**A1:** Correct equations.

**M1:** Eliminates  $X$  from the simultaneous equations and equates the coefficients of  $x$  leading to a quadratic equation in terms of  $m$ .

**dM1:** Dependent on the previous method, finds the value of the discriminant.

**A1:** Correct expression for the discriminant, states  $< 0$  and draws the required conclusion.



Q6.

Question	Scheme	Marks	AOs
(a)	Coordinates of $Q$ are $(8, -3, 2)$	B1 (1)	2.2a
(b)	Coordinates of $R$ are $\begin{pmatrix} \cos 120^\circ & 0 & \sin 120^\circ \\ 0 & 1 & 0 \\ -\sin 120^\circ & 0 & \cos 120^\circ \end{pmatrix} \begin{pmatrix} 8 \\ 3 \\ 2 \end{pmatrix} = \dots$ or $\begin{pmatrix} -0.5 & 0 & \frac{\sqrt{3}}{2} \\ 0 & 1 & 0 \\ -\frac{\sqrt{3}}{2} & 0 & -0.5 \end{pmatrix} \begin{pmatrix} 8 \\ 3 \\ 2 \end{pmatrix} = \dots$	M1	1.1a
	So $R$ is $(-4 + \sqrt{3}, 3, -4\sqrt{3} - 1)$	A1 (2)	1.1b
(c)	Finds the distance $PR = \sqrt{(8 - (-4 + \sqrt{3}))^2 + (3 - 3)^2 + (2 - (-4\sqrt{3} - 1))^2}$ Alternatively finds their $\overline{PR}$ or their $\overline{RP}$ then applies length of a vector formula. $\sqrt{(12 - \sqrt{3})^2 + (3 + 4\sqrt{3})^2}$ or $\sqrt{(-12 + \sqrt{3})^2 + (-3 - 4\sqrt{3})^2}$ $= \sqrt{204} \quad (= 2\sqrt{51}) \text{ cso}$	M1 A1 (2)	2.1 1.1b
(d)	$\overline{PR} \cdot \overline{PQ} = (-12 + \sqrt{3}, 0, -3 - 4\sqrt{3}) \cdot (0, -6, 0) = 0$ hence perpendicular	B1ft (1)	1.1b
(e)	$PQ$ is perpendicular to $PR$ so Area = $\frac{1}{2} \times PQ \times PR$ $= \frac{1}{2} \times 6 \times \sqrt{204} = 6\sqrt{51} \text{ cso}$	M1 A1 (2)	1.1b 1.1b
<b>(8 marks)</b>			

Notes:
(a) B1: Coordinates of $Q$ correctly stated, accept as a column vector.
(b) M1: Correct attempt to find coordinates of $R$ using the given matrix with $\theta = 120$ . Must be multiplying in the correct way round. With no working two correct values or $(-2.27, 3, -7.93)$ implies this mark. A1: Correct exact coordinates as shown in scheme. Accept as a column vector. Cos 120 and sin 120 must have been evaluated.
(c) M1: Applies the distance formula with the coordinates of $P$ and their $R$ . Alternatively finds the vector $\overline{PR}$ or $\overline{RP}$ then applies length of a vector formula. A1: Correct answer following correct coordinates of $R$ , must be a surd but need not be fully simplified.

(d)

**B1ft:** Shows the dot product is zero between the vectors  $\overline{PR}$  and  $\overline{PQ}$  and draws the conclusion perpendicular. Accept with  $\pm$  vectors for each. Follow through as long as the vectors are of the

correct form, so  $\overline{PR} = \begin{pmatrix} a \\ 0 \\ b \end{pmatrix}$  and  $\overline{PQ} = \begin{pmatrix} 0 \\ c \\ 0 \end{pmatrix}$

Note They could state if vectors  $\overline{PR}$  and  $\overline{PQ}$  are perpendicular then  $\overline{PR} \cdot \overline{PQ} = 0$  then shows  $\overline{PR} \cdot \overline{PQ} = 0$  this is B1

(e)

**M1:** Correct method for the area of the triangle, follow through on their coordinates of  $R$  and  $Q$ . May see longer methods if they do not realise the triangle is right angled.

**A1:** For  $6\sqrt{51}$  cso following correct coordinates of  $R$

#### Alternative 1

**M1** Complete method to find the correct area

Finding all the lengths  $|PQ| = 6$ ,  $|PR| = \sqrt{240} = 4\sqrt{15}$ ,  $|QR| = \sqrt{204} = 2\sqrt{51}$

Use cosine rule to find an angle e.g.  $\cos PRQ = \frac{240 + 204 - 36}{2 \times \sqrt{240} \times \sqrt{204}} = \frac{\sqrt{85}}{10}$

leading to  $PRQ = 22.7\dots$  or  $\sin PRQ = \sqrt{1 - \left(\frac{\sqrt{85}}{10}\right)^2} = \dots \left\{ \frac{\sqrt{15}}{10} \right\}$

Uses the area of the triangle  $= \frac{1}{2} \times \sqrt{240} \times \sqrt{204} \times \frac{\sqrt{15}}{10}$  or  $= \frac{1}{2} \times \sqrt{240} \times \sqrt{204} \times \sin 22.8$

**A1:** For  $6\sqrt{51}$

#### Alternative 2

**M1:** Uses  $\frac{1}{2}|\mathbf{a} \times \mathbf{b}|$  to find the required area

e.g.  $QP = \begin{pmatrix} 0 \\ 6 \\ 0 \end{pmatrix}$   $RP = \begin{pmatrix} 12 - \sqrt{3} \\ 0 \\ 3 + 4\sqrt{3} \end{pmatrix}$  cross product

$$\begin{vmatrix} 0 & 6 & 0 \\ 12 - \sqrt{3} & 0 & 3 + 4\sqrt{3} \end{vmatrix} = -6(12 - \sqrt{3})\mathbf{i} + 6(3 + 4\sqrt{3})\mathbf{k}$$

$$\text{Area} = \frac{1}{2} \sqrt{\left(-6(12 - \sqrt{3})\right)^2 + \left(6(3 + 4\sqrt{3})\right)^2} = \frac{1}{2} \sqrt{7344}$$

**A1:** For  $6\sqrt{51}$

Q7.

Question	Scheme	Marks	AOs
(a)(i)	Rotation	B1	1.1b
	90 degrees anticlockwise about the origin	B1	1.1b
(ii)	Stretch	B1	1.1b
	Scale factor 3 parallel to the $y$ -axis	B1	1.1b
		(4)	
(b)	$\mathbf{QP} = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 3 & 0 \end{pmatrix}$	B1	1.1b
		(1)	
(c)(i)	$ \mathbf{R}  = 3$	B1ft	1.1b
(ii)	The area scale factor of the transformation	B1	2.4
		(2)	
(7 marks)			
Notes			
<p>(a)(i)            B1: Identifies the transformation as a rotation            B1: Correct angle (allow equivalents in degrees or radians), direction and centre the origin</p> <p>(ii)            B1: Identifies the transformation as a stretch            B1: Correct scale factor and parallel to/in/along the <math>y</math>-axis/<math>y</math> direction</p> <p>(b)            B1: Correct matrix</p> <p>(c)(i)            B1ft: Correct value for the determinant (follow through their <math>\mathbf{R}</math>)</p> <p>(ii)            B1: Correct explanation, must include area            Note: scale factor of the transformation is B0</p>			

## Q8.

Question Number	Scheme	Marks
(a)	$\mathbf{A} = \begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix}, \mathbf{P} = \begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix}$ $\mathbf{A}^{-1} = \frac{1}{10} \begin{pmatrix} 3 & 1 \\ -4 & 2 \end{pmatrix}$	M1 Either $\frac{1}{10}$ or $\begin{pmatrix} 3 & 1 \\ -4 & 2 \end{pmatrix}$ Correct matrix seen. A1 [2]
(b) Way 1	$\mathbf{P} = \mathbf{AB}$ $\Rightarrow \mathbf{A}^{-1}\mathbf{P} = \mathbf{A}^{-1}\mathbf{AB} \Rightarrow \mathbf{B} = \mathbf{A}^{-1}\mathbf{P}$ $\mathbf{B} = \frac{1}{10} \begin{pmatrix} 3 & 1 \\ -4 & 2 \end{pmatrix} \begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix}$ $= \begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}$	Multiplies <b>their</b> $\mathbf{A}^{-1}$ by $\mathbf{P}$ in correct order. This substituted statement is sufficient. M1 At least 2 elements correct or $k \begin{pmatrix} 20 & 10 \\ 10 & -40 \end{pmatrix}$ oe. A1 May be unsimplified Correct simplified matrix. A1 [3]
(b) Way 2	$\{\mathbf{P} = \mathbf{AB} \Rightarrow\}$ $\begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ $\begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix} = \begin{pmatrix} 2a-c & 2b-d \\ 4a+3c & 4b+3d \end{pmatrix}$ $\Rightarrow a=2, c=1, b=1, d=-4$ $\text{So, } \mathbf{B} = \begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}$	Attempt to multiply $\mathbf{A}$ by $\mathbf{B}$ in the correct order and puts equal to $\mathbf{P}$ M1 At least 2 elements are correct. A1 Correct matrix. A1 [3] 5

**Q9.**

Question Number	Scheme	Marks
(i)	$A = \begin{pmatrix} p & 2 \\ 3 & p \end{pmatrix}, B = \begin{pmatrix} -5 & 4 \\ 6 & -5 \end{pmatrix}, M = \begin{pmatrix} a & -9 \\ 1 & 2 \end{pmatrix}$ $p, a$ are constants.	
(a)	$\{AB\} = \begin{pmatrix} -5p+12 & 4p-10 \\ -15+6p & 12-5p \end{pmatrix}$ At least 2 elements are correct. Correct matrix.	M1 A1
(b)	$\{AB + 2A = kI\}$ $\begin{pmatrix} -5p+12 & 4p-10 \\ -15+6p & 12-5p \end{pmatrix} + 2 \begin{pmatrix} p & 2 \\ 3 & p \end{pmatrix} = k \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ $\begin{pmatrix} -3p+12 & 4p-6 \\ -9+6p & 12-3p \end{pmatrix} = \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$ "4p-10" + 4 = 0 or "-15+6p" + 6 = 0 or "-9+6p" = "4p-6" $\Rightarrow p = \frac{3}{2}$ $k = -5\left(\frac{3}{2}\right) + 12 + 2\left(\frac{3}{2}\right) \Rightarrow k = \dots$ $k = \frac{15}{2}$ If 'simultaneous equations' used, allocate marks as below. Forms an equation in $p$ $p = \frac{3}{2}$ o.e. Substitutes their $p = \frac{3}{2}$ into "their (-5p+12)" + 2p to find a value for $k$ or eliminates $p$ to find $k$ . $k = \frac{15}{2}$ oe	[2] M1 A1 M1 A1
(ii) Way 1	$\pm \frac{270}{15} \{=\pm 18\}$ $\det M = (a)(2) - (-9)(1)$ $\Rightarrow 2a+9 = 18$ or $2a+9 = -18$ $\Rightarrow a = 4.5$ or $a = -13.5$ Can be implied from calculations. Applies $ad - bc$ to $M$ . Require clear evidence of correct formula being used for M1 if errors seen. Equates their $\det A$ to either 18 or -18 At least one of either $a = 4.5$ or $a = -13.5$ Both $a = 4.5$ and $a = -13.5$	B1 M1 M1 A1 A1
(ii) Way 2	Consider vertices of triangle with area 15 units e.g. (0,0),(15,0) and (0,2) and attempting 2 values of $a$ . e.g. $\begin{pmatrix} a & -9 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 0 & 15 & 0 \\ 0 & 0 & 2 \end{pmatrix} = \begin{pmatrix} 0 & 15a & -18 \\ 0 & 15 & 4 \end{pmatrix}$ e.g. $\frac{1}{2} \begin{vmatrix} 0 & 15a & -18 & 0 \\ 0 & 15 & 4 & 0 \end{vmatrix} = 270$ $\Rightarrow a = 4.5$ or $a = -13.5$ Pre-multiplies their matrix by $M$ and obtains single matrix Equates their determinant to 270 and attempts to solve. At least one of either $a = 4.5$ or $a = -13.5$ Both $a = 4.5$ and $a = -13.5$	[5] B1 M1 M1 A1 A1
		[5] 11

## Q10.

Question	Scheme		Marks	AOs
(a)	$\det(\mathbf{M}) = (1)(1) - (\sqrt{3})(-\sqrt{3})$		M1	1.1a
	$\mathbf{M}$ is non-singular because $\det(\mathbf{M}) = 4$ and so $\det(\mathbf{M}) \neq 0$		A1	2.4
			(2)	
(b)	Area( $S$ ) = 4(5) = 20		B1ft	1.2
			(1)	
(c)	$k = \sqrt{(1)(1) - (\sqrt{3})(-\sqrt{3})}$		M1	1.1b
	= 2		A1ft	1.1b
			(2)	
(d)	$\cos \theta = \frac{1}{2}$ or $\sin \theta = \frac{\sqrt{3}}{2}$ or $\tan \theta = \sqrt{3}$		M1	1.1b
	$\theta = 60^\circ$ or $\frac{\pi}{3}$		A1	1.1b
			(2)	
(7 marks)				
<b>Question Notes</b>				
(a)	M1	An attempt to find $\det(\mathbf{M})$ .		
	A1	$\det(\mathbf{M}) = 4$ and reference to zero, e.g. $4 \neq 0$ and conclusion.		
(b)	B1ft	20 or a correct ft based on their answer to part (a).		
(c)	M1	$\sqrt{(\text{their } \det \mathbf{M})}$		
	A1ft	2		
(d)	M1	Either $\cos \theta = \frac{1}{(\text{their } k)}$ or $\sin \theta = \frac{\sqrt{3}}{(\text{their } k)}$ or $\tan \theta = \sqrt{3}$		
	A1	$\theta = 60^\circ$ or $\frac{\pi}{3}$ . Also accept any value satisfying $360n + 60^\circ$ , $n \in \mathbb{Z}$ , o.e.		