



**MECHANICS 1 (A) TEST PAPER 1 : ANSWERS AND MARK SCHEME**

1. (a)  $AB^2 = 12.25 + 144 = 156.25$        $AB = 12.5 \text{ m}$       M1 A1  
 (b)  $12.5 \div 5 = 2.5 \text{ ms}^{-1}$       (c)  $(0.7\mathbf{i} - 2.4\mathbf{j}) \text{ ms}^{-1}$       B1 B1; M1 A1      6
2. (a)  $0.8g = 2T \sin 30^\circ$        $T = 0.8g = 7.84 \text{ N}$       B1 M1 A1  
 (b)  $F = T \cos 30^\circ$ ,  $0.8g = T \sin 30^\circ$        $F = 0.8g\sqrt{3} = 13.6 \text{ N}$       B1 B1 M1 A1      7
3. (a)  $s = ut + \frac{1}{2}at^2$ :  $3u + 4.5a = 6$ ,  $9u + 40.5a = 39$   
 $21 = 27a$        $a = \frac{7}{9} \text{ ms}^{-2}$       (b)  $u = \frac{5}{6} \text{ ms}^{-1}$       M1 A1 A1  
 M1 A1; M1 A1      7
4. (a)  $F = Ma$ , so  $F = 3M$       M1 A1  
 (b)  $F - \mu Mg = 3M$        $F = M(3 + \mu g)$       M1 A1  
 (c)  $3 + \mu g = \frac{1}{2}g$        $\mu = \frac{1}{2} - \frac{3}{g} = 0.194$       M1 A1 A1      7
5. (a)  $F = ma$  for each:  $2.4g - T = 2.4a$ ,  $T - 1.8g = 1.8a$   
 Add:  $0.6g = 4.2a$        $a = \frac{1}{7}g = 1.4 \text{ ms}^{-2}$        $T = 20.2 \text{ N}$       M1 A1 A1  
 (b) Now  $1.8g - T = 1.8(0.7)$  so  $T = 16.38$ , and  $T - mg = m(0.7)$   
 $10.5m = 16.38$        $m = 1.56$       M1 A1      12
6. (a) (i)  $\frac{7}{3} \text{ ms}^{-2}$ ,  $-\frac{7}{4} \text{ ms}^{-2}$ ,  $0 \text{ ms}^{-2}$       B2 (-1 each error)  
 (ii)  $45 \times 15 + \frac{1}{2} \times 35^2 = 1287.5 \text{ m}$       M1 A1  
 (b) (i)   
 (ii)       B2 B2  
 (c)  $15 \times 9T + 35 \times 3.5T = 3708$        $257.5T = 3708$        $T = 14.4$       M1 A1 M1 A1      12
7. (a)  $60 \times 4 = 60u + 90 \times 6u$        $600u = 240$        $u = 0.4$       M1 A1 M1 A1  
 (b) Change in momentum of  $B = 0.09 \times 2.4 = 0.216 \text{ Ns}$       M1 A1 B1  
 (c)  $60(2) + 90(-8) = 60(-7) + 90v_B$   
 $-180 = 90v_B$        $v_B = -2 \text{ ms}^{-1}$ , direction unchanged      M1 A1 A1      12
8. (a)  $T_P + T_Q = 22g$       M(A):  $1.5(6g) + 3.5(8g) = 4.5T_Q$       B1 M1 A1  
 $T_Q = 37g \div 4.5 = 80.6 \text{ N}$        $T_P = 22g - T_Q = 135 \text{ N}$       M1 A1 A1  
 (b)  $2.5T_Q = 22g + mg$       M(A):  $mg(2.25) + 9g + 28g = 4.5T_Q$       B1 M1 A1  
 $2.25m + 37 = 39.6 + 1.8m$        $0.45m = 2.6$        $m = 5.78$       M1 A1 A1      12