


07 May 2011  
22:28

MI JUNE 02

Q1  $S = 50m$   
 (a)  $u = 5ms^{-1}$   
 $t = 4s$   
 $a = ?$   
 $S = ut + \frac{1}{2}at^2$   
 $50 = 5 \cdot 4 + \frac{1}{2}a \cdot 4^2$   
 $50 = 20 + 8a$   
 $a = 3.75ms^{-2}$  (3)

(b)  $S = ?$   
 $u = 5ms^{-1}$   
 $V = 30ms^{-1}$   
 $a = 3.75ms^{-2}$   
 $V^2 = u^2 + 2as$   
 $30^2 = 5^2 + 2 \cdot 3.75s$   
 $s = 116 \frac{2}{3}m$  (116.7m) (3)



(a) Impulse on A = 3.6Ns  
 Impulse = change in momentum:  
 Before:  $0.5 \times 5 = 2.5$   
 After:  $-0.5u = -0.5u$   
 $3.6 = 2.5 + 0.5u$   
 $u = 2.2ms^{-1}$  (3)


(b) speed of b =  $1ms^{-1}$   
 Momentum before = after:  
 $0.5 \times 5 - m \times 3 = -0.5 \times 2.2 + m \times 1$   
 $4m = 3.6 \Rightarrow m = 0.9kg$   
 (i) B changes direction:  
 $0.5 \times 5 - m \times 3 = -0.5 \times 2.2 - m \times 1$   
 $2m = 3.6 \Rightarrow m = 1.8kg$  (4)

MI JUNE 02  
 Q4  $\mu = 0.4$   
 $F_{max} = \mu R_N$   
 $F_{max} = 0.4R_N$


$R_N = 6g \cos 30 + P \sin 30$   
 $R_N = 50.9(222) + 0.5P$  (i)  
 $6g \sin 30 + F_{max} = P \cos 30$  substitute  $R_N$  into this equation (ii)  
 $29.4 + 0.4R_N = 0.866(0.25)P$  (ii)  
 $29.4 + 0.4(50.9 + 0.5P) = 0.866P$   
 $29.4 + 20.3688 + 0.2P = 0.866P$   
 (b)  $P = 74.7N$  (3)

from (i)  
 (a)  $R_N = 50.9 + 0.5 \times P$   
 $= 88.3N$  (88N) (4)

(c)  $F_{max} = \mu R_N$   
 $R_N = 50.9N$   
 $F_{max} = 0.4 \times 50.9 = 20.36N$   
 $F_{max} < 29.4N$   
 so the box will move towards 29.4 force (obviously!!)



(a) Either take moments about C:  
 $\curvearrowright \quad \curvearrowleft$   
 $30 \times 16 = 20 \times W + 70 \times 5$   
 $480 = 20W + 350$   
 $W = 6.5N$  (3)  
 or  $\uparrow = \downarrow$   $R = 21 + W$   
 and moments about A:  
 $\curvearrowleft \quad \curvearrowright$   
 $30 \times R = 50W + 100 \times 5$   
 $30(21 + W) = 50W + 500$   
 $630 + 30W = 50W + 500$   
 $20W = 130$   
 $W = 6.5N$  (3)



(c) The height of the strings doesn't affect the moments equation (isn't taken into consideration) (1)

JUNE 02 MI

Q5. (a) at  $t = 3$   $v = i - 2j$   
 $d = 90^\circ + \tan^{-1}(\frac{2}{1}) = 153.4^\circ$   
 (3)

Q5(b) acceleration =  $\frac{\text{change in velocity}}{\text{time}}$   
 $a = \frac{(i - 2j) - (-5i + 7j)}{3} = \frac{6i - 9j}{3} = 2i - 3j$   
 $a = 2i - 3j$  (2)

(c)  $F = ma = 2 \times (2i - 3j) = 4i - 6j$   
 $F = \sqrt{4^2 + 6^2} = 2\sqrt{13} = 7.21N$  (3)

(d) Velocity = initial velocity + t \* acceleration  
 $v = (-5 + 2t)i + (7 - 3t)j$  (2)

(e)  $P$  parallel to  $i + j$  when its velocity is a multiple of  $(i + j)$  i.e.  $i$  and  $j$  components are equal  
 $(-5 + 2t)i + (7 - 3t)j = k(i + j)$   
 $\Rightarrow -5 + 2t = 7 - 3t$   
 $5t = 12$   
 $t = 2.4s$  (3)

M1 June 02




(a)  $A+B+C=27m$   
 (b)  $a = \frac{3-0}{t} \Rightarrow 2 = \frac{3}{t}$   
 $t = 1.5s$


(b) Alternatively:  
 $\Delta A = \frac{1}{2} \cdot 1.5 \cdot 3 = 2.25$   
 $\square B = T \cdot 3 = 3T$   
 $\Delta C = \frac{1}{2} \cdot 2.5 \cdot 3 = 3.75$   
 $3T + 6 = 27$   
 $T = 7s$

$A+B+C=27m$   
 Area  $\Delta = \frac{T+(T+4)}{2} \times 3 = \frac{2T+4}{2} \times 3 = 3(T+2)$

$3(T+2) = 27$   
 $T+2=9$   
 $T=7s$  (3)



M1 June 02



(a) B: N2L  $ma = mg \sin 30^\circ - T$   
 A: N2A  $2ma = T - F_{max}$   $F_{max} = \mu R_N$   
 B:  $ma = mg \sin 30^\circ - T$   
 A:  $2ma = T - 2\mu mg$   $= \mu 2mg$   
 $3\mu a = mg(\sin 30^\circ - 2\mu)$   
 $a = \frac{1}{3}(\frac{1}{2} - 2\mu)g$   
 $a = \frac{1}{6}(1 - 4\mu)g$  (7)

(b)  $\mu = 0.2 \Rightarrow a = \frac{1}{6}(1 - 4 \times \frac{1}{5})g = \frac{1}{30}g$   
 When the string snaps the velocity of the system is:  
 $v^2 = u^2 + 2as$  ( $u=0, s=h$ )  
 $v^2 = 2 \times \frac{1}{30}g \times h$   
 $v^2 = \frac{gh}{15}$

M1 June 02

(b) so  $u = \sqrt{\frac{gh}{15}}$  is the initial velocity  
 $v=0$  is the final

$F = ma$    
 $2ma = -F_{max}$   
 $2ma = -2\mu mg$   $\mu = 0.2$   
 $a = -0.2g$

Hence the distance:  $v^2 = u^2 + 2as$   
 $0 = \frac{gh}{15} - 2 \times 0.2g \times s$   
 $0.4g \times s = \frac{gh}{15}$   $\frac{1}{15} = 0.4$   
 $s = \frac{gh}{6} = \frac{h}{6}$   $= \frac{1}{15} \div \frac{2}{5}$   
 $= \frac{1}{15} \times \frac{5}{2}$   
 $= \frac{1}{6}$   
 Hence the total distance:  
 $h + \frac{h}{6} = \frac{7h}{6}$  (6)

(c) weight of pulley or string, friction on slope or pulley; (2)

M1 June 02

(d)   
 $a = 2ms^{-2}$   
 N2L  $F = ma$   
 $T - 200g = 200 \times 2$   
 $T = 400 + 200g$   
 $T = 2360N$  (3)

(e)   
 $a = -1.2ms^{-2}$   
 N2L  $F = ma$   
 $R_N - 80g = 80 \times (-1.2)$   
 $R_N = 80g - 96$   
 $R_N = 688N$  (3)