

Write your name here

Surname

Other names

Pearson
Edexcel GCE

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Mechanics M1

Advanced/Advanced Subsidiary

Wednesday 6 June 2018 – Morning

Time: 1 hour 30 minutes

Paper Reference

6677/01**You must have:**

Mathematical Formulae and Statistical Tables (Pink)

Total Marks

--

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P51517A

©2018 Pearson Education Ltd.

1/1/1/



Pearson

3.




Figure 2

A wooden beam AB , of mass 150 kg and length 9 m, rests in a horizontal position supported by two vertical ropes. The ropes are attached to the beam at C and D , where $AC = 1.5$ m and $BD = 3.5$ m. A gymnast of mass 60 kg stands on the beam at the point P , where $AP = 3$ m, as shown in Figure 2. The beam remains horizontal and in equilibrium.

By modelling the gymnast as a particle, the beam as a uniform rod and the ropes as light inextensible strings,

- (a) find the tension in the rope attached to the beam at C . (3)

The gymnast at P remains on the beam at P and another gymnast, who is also modelled as a particle, stands on the beam at B . The beam remains horizontal and in equilibrium. The mass of the gymnast at B is the largest possible for which the beam remains horizontal and in equilibrium.

- (b) Find the tension in the rope attached to the beam at D . (4)



7.




Figure 4

A particle P of mass $4m$ is held at rest at the point X on the surface of a rough inclined plane which is fixed to horizontal ground. The point X is a distance h from the bottom of the inclined plane. The plane is inclined to the horizontal at an angle α where $\tan \alpha = \frac{3}{4}$.

The coefficient of friction between P and the plane is $\frac{1}{4}$. The particle P is attached to one end of a light inextensible string. The string passes over a small smooth pulley which is fixed at the top of the plane. The other end of the string is attached to a particle Q of mass m which hangs freely at a distance d , where $d > h$, below the pulley, as shown in Figure 4.

The string lies in a vertical plane through a line of greatest slope of the inclined plane. The system is released from rest with the string taut and P moves down the plane.

For the motion of the particles before P hits the ground,

- state which of the information given above implies that the magnitudes of the accelerations of the two particles are the same, (1)
- write down an equation of motion for each particle, (5)
- find the acceleration of each particle. (5)

When P hits the ground, it immediately comes to rest. Given that Q comes to instantaneous rest before reaching the pulley,

- show that $d > \frac{28h}{25}$. (5)



