

GCE Examinations

Mechanics

Module M1

Advanced Subsidiary / Advanced Level

Paper J

Time: 1 hour 30 minutes

Instructions and Information

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 7 questions.

When a numerical value of g is required, use $g = 9.8 \text{ m s}^{-2}$.

Advice to Candidates

You must show sufficient working to make your methods clear to an examiner. Answers without working will gain no credit.



Written by Shaun Armstrong & Chris Huffer

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

1. At time $t = 0$, a particle of mass 2 kg has velocity $(8\mathbf{i} + \lambda\mathbf{j}) \text{ m s}^{-1}$ where \mathbf{i} and \mathbf{j} are horizontal perpendicular unit vectors and $\lambda > 0$.

Given that the speed of the particle at time $t = 0$ is 17 m s^{-1} ,

- (a) find the value of λ . **(3 marks)**

The particle experiences a constant retarding force \mathbf{F} so that when $t = 5$, it has velocity $(3\mathbf{i} + 5\mathbf{j}) \text{ m s}^{-1}$.

- (b) Show that \mathbf{F} can be written in the form $\mu(\mathbf{i} + 2\mathbf{j}) \text{ N}$ where μ is a constant which you should find.

(5 marks)

2. A monk uses a small brush to clean the stone floor of a monastery by pushing the brush with a force of P Newtons at an angle of 60° to the vertical. He moves the brush at a constant speed. The mass of the brush is 0.5 kg and the coefficient of friction between the brush and the floor is $\frac{1}{\sqrt{3}}$. The brush is modelled as a particle and air resistance is ignored.

- (a) Show that $P = \frac{g}{2}$ Newtons. **(7 marks)**

- (b) Explain why it is reasonable to ignore air resistance in this situation. **(1 mark)**
-

3. A small van of mass 1500 kg is used to tow a car of mass 750 kg by means of a rope of length 9 m joined to both vehicles. The van sets off with the rope slack and reaches a speed of 2 m s^{-1} just before the rope becomes taut and jerks the car into motion. Immediately after the rope becomes taut, the van and car travel with common speed $V \text{ m s}^{-1}$.

- (a) Show that $V = \frac{4}{3}$. **(3 marks)**

- (b) Calculate the magnitude of the impulse on the car when the rope tightens. **(2 marks)**

The van and car eventually reach a steady speed of 18 m s^{-1} with the rope taut when a child runs out into the road, 30 m in front of the van. The van driver brakes sharply and decelerates uniformly to rest in a distance of 27 m.

It takes the driver of the car 1 second to react to the van starting to brake. He then brakes and the car decelerates uniformly at $f \text{ m s}^{-2}$, coming to rest before colliding with the van.

- (c) Find the set of possible values of f . **(5 marks)**
-

4.




Fig. 1

Figure 1 shows a weight A of mass 6 kg connected by a light, inextensible string which passes over a smooth, fixed pulley to a box B of mass 5 kg . There is an object C of mass 3 kg resting on the horizontal floor of box B .

The system is released from rest. Find, giving your answers in terms of g ,

- (a) the acceleration of the system, **(4 marks)**
- (b) the force on the pulley. **(3 marks)**
- (c) Show that the reaction between C and the floor of B is $\frac{18}{7}g$ newtons. **(3 marks)**

5. Two flies P and Q , are crawling vertically up a wall. At time $t = 0$, the flies are at the same height above the ground, with P crawling at a steady speed of 4 cm s^{-1} .

Q starts from rest at time $t = 0$ and accelerates uniformly to a speed of 6 cm s^{-1} in 6 seconds. Fly Q then maintains this speed.

- (a) Find the value of t when the two flies are moving at the same speed. **(3 marks)**
- (b) Sketch on the same diagram, speed-time graphs to illustrate the motion of the two flies. **(3 marks)**

Given that the distance of the two flies from the top of the wall at time $t = 0$ is $x\text{ cm}$ and that Q reaches the top of the wall first,

- (c) show that $x > 36$. **(5 marks)**

Turn over

6.




Fig. 2

Figure 2 shows a uniform plank AB of length 8 m and mass 50 kg suspended horizontally by two light vertical inextensible strings attached at either end of the plank. The maximum tension that either string can support is $40g$ N.

A rock of mass M kg is placed on the plank at A and rolled along the plank to B without either string breaking.

- (a) Explain, with the aid of a sketch-graph, how the tension in the string at A varies with x , the distance of the rock from A . **(3 marks)**

- (b) Show that $M \leq 15$. **(5 marks)**

The first rock is removed and a second rock of mass 20 kg is placed on the plank.

- (c) Find the fraction of the plank on which the rock can be placed without one of the strings breaking. **(6 marks)**

7. At 6 a.m. a cargo ship has position vector $(7\mathbf{i} + 56\mathbf{j})$ km relative to a fixed origin O on the coast and moves with constant velocity $(9\mathbf{i} - 6\mathbf{j})$ km h⁻¹.

A ferry sails from O at 6 a.m. and moves with constant velocity $(12\mathbf{i} + 18\mathbf{j})$ km h⁻¹. The unit vectors \mathbf{i} and \mathbf{j} are directed due east and due north respectively.

- (a) Show that the position vector of the cargo ship t hours after 6 a.m. is given by

$$[(7 + 9t)\mathbf{i} + (56 - 6t)\mathbf{j}] \text{ km,}$$

and find the position vector of the ferry in terms of t . **(3 marks)**

- (b) Show that if both vessels maintain their course and speed, they will collide and find the time and position vector at which this occurs.

(6 marks)

At 8 a.m. the captain of the ferry realises that a collision is imminent and changes course so that the ferry now has velocity $(21\mathbf{i} + 6\mathbf{j})$ km h⁻¹.

- (c) Find the distance between the two ships at the time when they would have collided.

(5 marks)

END