

Exercise 2F

- 1 a Take downwards as the positive direction.

$$s = 28, u = 0, a = 9.8, t = ?$$

$$s = ut + \frac{1}{2}at^2$$

$$28 = 0 \times t + \frac{1}{2} \times 9.8 \times t^2 = 4.9t^2$$

$$t = \sqrt{\frac{28}{4.9}} = 2.4 \text{ (to 2 s.f.)}$$

The time taken for the diver to hit the water is 2.4 s.

b $v^2 = u^2 + 2as$

$$v^2 = 0 + 2 \times 9.8 \times 28 = 548.8$$

$$v = \sqrt{548.8} = 23 \text{ (to 2 s.f.)}$$

When the diver hits the water, he is travelling at 23 m s^{-1} .

- 2 Take upwards as the positive direction.

$$u = 20, a = -9.8, s = 0, t = ?$$

$$s = ut + \frac{1}{2}at^2$$

$$0 = 20t - 4.9t^2 = t(20 - 4.9t), \quad t \neq 0$$

$$t = \frac{20}{4.9} = 4.1 \text{ (to 2 s.f.)}$$

The time of flight of the particle is 4.1 s.

- 3 Take downwards as the positive direction.

$$u = 18, a = 9.8, t = 1.6, s = ?$$

$$s = ut + \frac{1}{2}at^2 = 18 \times 1.6 + 4.9 \times 1.6^2 = 41 \text{ (to 2 s.f.)}$$

The height of the tower is 41 m.

- 4 a Take upwards as the positive direction.

$$u = 24, a = -9.8, v = 0, s = ?$$

$$v^2 = u^2 + 2as$$

$$0^2 = 24^2 - 2 \times 9.8 \times s$$

$$s = \frac{24^2}{2 \times 9.8} = 29 \text{ (to 2 s.f.)}$$

The greatest height reached by the pebble above the point of projection is 29 m.

4 b $u = 24, a = -9.8, v = 0, t = ?$

$$v = u + at$$

$$0 = 24 - 9.8t$$

$$t = \frac{24}{9.8} = 2.4 \text{ (to 2 s.f.)}$$

The time taken to reach the greatest height is 2.4 s.

5 a Take upwards as the positive direction.

$$u = 18, a = -9.8, s = 15, v = ?$$

$$v^2 = u^2 + 2as = 18^2 - 2 \times 9.8 \times 15 = 30$$

$$v = \sqrt{30} = \pm 5.5 \text{ (to 2 s.f.)}$$

The speed of the ball when it is 15 m above its point of projection is 5.5 m s^{-1} .

b $u = 18, a = -9.8, s = -4, v = ?$

$$v^2 = u^2 + 2as = 18^2 + 2 \times (-9.8) \times (-4) = 324 + 78.4 = 402.4$$

$$v = -\sqrt{402.2} = -20 \text{ (to 2 s.f.)}$$

The speed with which the ball hits the ground is 20 m s^{-1} .

6 a Take downwards as the positive direction.

$$s = 80, u = 4, a = 9.8, v = ?$$

$$v^2 = u^2 + 2as$$

$$= 4^2 + 2 \times 9.8 \times 80 = 1584$$

$$v = \sqrt{1584} = 40 \text{ (to 2 s.f.)}$$

The speed with which P hits the ground is 40 m s^{-1} .

b $u = 4, a = 9.8, v = \sqrt{1584}, t = ?$

$$v = u + at$$

$$\sqrt{1584} = 4 + 9.8t$$

$$t = \frac{\sqrt{1584} - 4}{9.8} = 3.7 \text{ (to 2 s.f.)}$$

The time P takes to reach the ground is 3.7 s.

- 7 a Take upwards as the positive direction.

$$v = -10, a = -9.8, t = 5, u = ?$$

$$v = u + at$$

$$-10 = u - 9.8 \times 5$$

$$u = 9.8 \times 5 - 10 = 39$$

The speed of projection of P is 39 m s^{-1} .

- b $u = 39, v = 0, a = -9.8, s = ?$

$$v^2 = u^2 + 2as$$

$$0^2 = 39^2 - 2 \times 9.8 \times s$$

$$s = \frac{1521}{2 \times 9.8} = 78 \text{ (to 2 s.f.)}$$

The greatest height above X attained by P during its motion is 78 m .

- 8 Take upwards as the positive direction.

$$u = 21, t = 4.5, a = -9.8, s = ?$$

$$s = ut + \frac{1}{2}at^2 = 21 \times 4.5 - 4.9 \times 4.5^2 = -4.7 \text{ (to 2 s.f.)}$$

The height above the ground from which the ball was thrown is 4.7 m .

- 9 a Take upwards as the positive direction.

$$s = -3, u = 16, a = -9.8, t = ?$$

$$s = ut + \frac{1}{2}at^2$$

$$-3 = 16t - 4.9t^2$$

$4.9t^2 - 16t - 3 = 0$, so using the quadratic formula,

$$t = \frac{-(-16) \pm \sqrt{(-16)^2 - 4 \times (4.9) \times (-3)}}{2 \times (4.9)}$$

$t = 3.4$ (to 2 s.f.) as we may discount the negative answer.

The time of flight of the stone is 3.4 s .

9 b $u = 16, v = 0, a = -9.8, s = ?$

$$v^2 = u^2 + 2as$$

$$0^2 = 16^2 - 2 \times 9.8 \times s$$

$$s = \frac{16^2}{2 \times 4.9} = 13 \text{ (to 2 s.f.)}$$

The total distance travelled by the stone is $(2 \times 13 + 3) \text{ m} = 29 \text{ m}$.

10 Take upwards as the positive direction.

$$u = 24.5, a = -9.8, s = 21, t = ?$$

$$s = ut + \frac{1}{2}at^2$$

$$21 = 24.5t - 4.9t^2$$

$$4.9t^2 - 24.5t + 21 = 0$$

Using the quadratic formula,

$$t = \frac{-(-24.5) \pm \sqrt{(-24.5)^2 - 4 \times (4.9) \times (21)}}{2 \times (4.9)}$$

$$= 1.1 \text{ or } 3.9$$

The difference between these times is

$$(3.9 - 1.1) \text{ s} = 2.8 \text{ s}$$

The total time for which the particle is 21 m or more above its point of projection is 2.8 s.

11 a Take upwards as the positive direction.

$$v = \frac{1}{3}u, a = -9.8, t = 2, u = ?$$

$$v = u + at$$

$$\frac{1}{3}u = u - 9.8 \times 2$$

$$\frac{2}{3}u = 19.6 \Rightarrow u = \frac{3}{2} \times 19.6 = 29.4$$

$$u = 29 \text{ (to 2 s.f.)}$$

11 b $u = 29.4, s = 0, a = -9.8, t = ?$

$$s = ut + \frac{1}{2}at^2$$

$$0 = 29.4t - 4.9t^2 = t(29.4 - 4.9t), t \neq 0$$

$$t = \frac{29.4}{4.9} = 6$$

The time from the instant that the particle leaves O to the instant that it returns to O is 6 s.

12 For A , take downwards as the positive direction, $s_A = ut + \frac{1}{2}at^2 = 5t + 4.9t^2$

For B , take upwards as the positive direction, $s_B = ut + \frac{1}{2}at^2 = 18t - 4.9t^2$

$$s_A + s_B = 46$$

$$(5t + 4.9t^2) + (18t - 4.9t^2) = 46$$

$$23t = 46 \Rightarrow t = 2$$

Substitute $t = 2$ into $s_A = 5t + 4.9t^2$

$$s_A = 5 \times 2 + 4.9 \times 2^2 = 29.6 = 30 \text{ (to 2 s.f.)}$$

The distance of the point where A and B collide from the point where A was thrown is 30 m.

13 a Find the speed, u_1 say, immediately before the ball strikes the floor.

$$u = 0, a = 9.8, s = 10, v = u_1$$

$$v^2 = u^2 + 2as$$

$$u_1^2 = 0^2 + 2 \times 9.8 \times 10 = 196$$

$$u_1 = \sqrt{196} = 14$$

The speed of the first rebound, u_2 say, is given by

$$u_2 = \frac{3}{4}u_1 = \frac{3}{4} \times 14 = 10.5$$

Find the maximum height, h_1 say, reached after the first rebound.

$$u = 10.5, v = 0, a = -9.8, s = h_1$$

$$v^2 = u^2 + 2as$$

$$0^2 = 10.5^2 - 2 \times 9.8 \times h_1$$

$$13 \text{ a } h_1 = \frac{10.5^2}{2 \times 9.8} = 5.6 \text{ (to 2 s.f.)}$$

The greatest height above the floor reached by the ball the first time it rebounds is 5.6 m.

- b** Immediately before the ball strikes the floor for the second time, its speed is again $u_2 = 10.5$ by symmetry. The speed of the second rebound, u_3 , say, is given by

$$u_3 = \frac{3}{4}u_2 = \frac{3}{4} \times 10.5 = 7.875$$

Find the maximum height, h_2 , say, reached after the second rebound.

$$u = 7.875, \quad v = 0, \quad a = -9.8, \quad s = h_2$$

$$v^2 = u^2 + 2as$$

$$0^2 = 7.875^2 - 2 \times 9.8 \times h_2$$

$$h_2 = \frac{7.875^2}{2 \times 9.8} = 3.2 \text{ (to 2 s.f.)}$$

The greatest height above the floor reached by the ball the second time it rebounds is 3.2 m.

Challenge

- 1 a** Take upwards as the positive direction.

$$\text{For } P, s = ut + \frac{1}{2}at^2 \text{ gives } s_P = 12t - 4.9t^2$$

$$\text{For } Q, s = ut + \frac{1}{2}at^2$$

Q has been moving for 1 less second than P , so

$$s_Q = 20(t-1) - 4.9(t-1)^2$$

At the point of collision $s_P = s_Q$

$$\begin{aligned} 12t - 4.9t^2 &= 20(t-1) - 4.9(t-1)^2 \\ &= 20t - 20 - 4.9t^2 + 9.8t - 4.9 \end{aligned}$$

$$24.9 = 17.8t \Rightarrow t = \frac{24.9}{17.8} = 1.4 \text{ (to 2 s.f.)}$$

The time between the instant when P is projected and the instant when P and Q collide is 1.4 s.

Challenge

- 1 b Substitute t into $s_p = 12t - 4.9t^2$ from part a

$$s_p = 12t - 4.9t^2 \approx 12 \times 1.4 - 4.9 \times 1.4^2 = 7.2 \text{ (to 2 s.f.)}$$

The distance of the point where P and Q collide from O is 7.2 m.

- 2 Take downwards as positive.

For 1st stone: $u = 0$, $t = t_1$, $a = 9.8$, $s = h$

$$s = ut + \frac{1}{2}at^2$$

$$h = 0 \times t_1 + \frac{1}{2} \times 9.8 \times t_1^2 = 4.9t_1^2$$

For 2nd stone: $u = 25$, $t = t_1 - 2$, $a = 9.8$, $s = h$

$$s = ut + \frac{1}{2}at^2$$

$$\begin{aligned} h &= 25(t_1 - 2) + \frac{1}{2}(9.8 \times (t_1 - 2)^2) \\ &= 25t_1 - 50 + 4.9 \times (t_1^2 - 4t_1 + 4) \\ &= 25t_1 - 50 + 4.9t_1^2 - 19.6t_1 + 19.6 \\ &= 4.9t_1^2 + 5.4t_1 - 30.4 \end{aligned}$$

Substituting for h from information for first stone:

$$4.9t_1^2 = 4.9t_1^2 + 5.4t_1 - 30.4$$

$$30.4 = 5.4t_1$$

$$t_1 = \frac{30.4}{5.4} = 5.629$$

Putting this value into equation for first stone:

$$h = 4.9 \times 5.629^2 = 4.9 \times 31.69 = 155 \text{ (to 3 s.f.)}$$

The height of the building is 155 m.