

Exercise 7B

1 From symmetry the tension in both strings is the same.

$$R(\uparrow)$$


$$T \sin 45^\circ + T \sin 45^\circ - 5g = 0$$

$$\therefore 2T \sin 45^\circ = 5g$$

$$T = \frac{5g}{2 \sin 45^\circ}$$

$$= \frac{49\sqrt{2}}{2}$$

$$T = 34.6 \text{ N (3 s.f.)}$$




2 a Let the tension in the string be T N

$$R(\leftarrow)$$

$$T \sin 30^\circ - 10 = 0$$

$$\therefore T = \frac{10}{\sin 30^\circ}$$

$$T = 20 \text{ N}$$



b $R(\uparrow)$

$$T \cos 30^\circ - mg = 0$$

$$mg = 20 \cos 30^\circ \quad (\text{since } T = 20 \text{ N})$$

$$\therefore m = \frac{20 \cos 30^\circ}{g}$$

$$= \frac{10\sqrt{3}}{g}$$

$$= 1.8 \text{ kg (2 s.f.)}$$

3 Let the tension in the string be T N.

$$R(\rightarrow)$$


$$8 - T \sin \theta = 0$$

$$\therefore T \sin \theta = 8 \quad (1)$$

$$R(\uparrow)$$

$$T \cos \theta - 12 = 0$$

$$\therefore T \cos \theta = 12 \quad (2)$$



- 3 a Divide equation (1) by equation (2) to eliminate the tension T .

$$\frac{T \sin \theta}{T \cos \theta} = \frac{8}{12}$$

$$\therefore \tan \theta = \frac{2}{3}$$

$$\therefore \theta = 33.7^\circ \text{ (3 s.f.)}$$

- b Substitute into equation (1)

$$T \sin 33.7^\circ = 8$$

$$T = \frac{8}{\sin 33.7^\circ} \\ = 14.4 \text{ (3 s.f.)}$$

- 4 Let the tension in the strings be T N and S N as shown in the figure.

$R(\leftarrow)$

$$T \cos 60^\circ - S \cos 45^\circ = 0$$

$$\therefore \frac{T}{2} - \frac{S}{\sqrt{2}} = 0$$

$$\therefore T = S\sqrt{2} \quad (1)$$

$R(\uparrow)$

$$T \sin 60^\circ + S \sin 45^\circ - 6g = 0$$

$$T \frac{\sqrt{3}}{2} + S \frac{1}{\sqrt{2}} = 6g \quad (2)$$

Substitute $T = S\sqrt{2}$ from (1) into equation (2)

$$S \left(\sqrt{2} \times \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \right) = 6g$$


$$S \left(\frac{\sqrt{3}+1}{\sqrt{2}} \right) = 6g$$

$$S = \frac{6g\sqrt{2}}{(\sqrt{3}+1)}$$

$$= 3g\sqrt{2}(\sqrt{3}-1)$$

$$= 30 \text{ (2 s.f.)}$$

$$\text{and } T = 6g(\sqrt{3}-1) = 43 \text{ (2 s.f.)}$$



Mechanics 1

Solution Bank

- 5 a Let the tension in the string be T and the mass of the bead be m .


Resolve horizontally first to find T :

$R(\rightarrow)$

$$T \cos 30^\circ - T \cos 60^\circ - 2 = 0$$

$$T(\cos 30^\circ - \cos 60^\circ) = 2$$

$$\begin{aligned} \therefore T &= \frac{2}{\cos 30^\circ - \cos 60^\circ} \\ &= \frac{4}{\sqrt{3} - 1} \\ &= \frac{4(\sqrt{3} + 1)}{(\sqrt{3} - 1)(\sqrt{3} + 1)} \\ &= \frac{4(\sqrt{3} + 1)}{2} \\ &= 2(\sqrt{3} + 1) = 5.46 \text{ N (3 s.f.)} \end{aligned}$$



- b $R(\uparrow)$

$$T \sin 60^\circ + T \sin 30^\circ - mg = 0$$

$$mg = T(\sin 60^\circ + \sin 30^\circ)$$

$$m = \frac{2}{g} (\sqrt{3} + 1) \left(\frac{\sqrt{3}}{2} + \frac{1}{2} \right) \quad (\text{using } T = 2(\sqrt{3} + 1) \text{ from part a})$$

$$= \frac{4 + 2\sqrt{3}}{g}$$

$$= 0.76 \text{ kg (2 s.f.)}$$

- c Modelling the bead as smooth assumes there is no friction between it and the string.

6 Let the tension in the string be T and the mass of the bead be m .


a Resolve horizontally first to find T .

$R(\rightarrow)$

$$2 - T \cos 60^\circ - T \cos 30^\circ = 0$$

$$T(\cos 60^\circ + \cos 30^\circ) = 2$$

$$\begin{aligned} \therefore T &= \frac{2}{\cos 60^\circ + \cos 30^\circ} \\ &= \frac{4}{1 + \sqrt{3}} \\ &= \frac{4}{1 + \sqrt{3}} \times \frac{\sqrt{3} - 1}{\sqrt{3} - 1} \quad (\text{to rationalise the denominator}) \\ &= 2(\sqrt{3} - 1) \\ &= 1.46 \text{ (3 s.f.)} \end{aligned}$$



b $R(\uparrow)$

$$T \sin 60^\circ - T \sin 30^\circ - mg = 0$$

$$mg = T(\sin 60^\circ - \sin 30^\circ)$$

$$= 2(\sqrt{3} - 1) \left(\frac{\sqrt{3}}{2} - \frac{1}{2} \right) \quad (\text{using } T = 2(\sqrt{3} - 1) \text{ from a})$$

$$= (\sqrt{3} - 1)^2$$

$$= 4 - 2\sqrt{3}$$

$$m = \frac{(4 - 2\sqrt{3})}{g}$$

$$= 0.055 \text{ kg} = 55 \text{ g}$$

7 $\tan \theta = \frac{12}{5} \Rightarrow \sin \theta = \frac{12}{13}$ and $\cos \theta = \frac{5}{13}$

Let the normal reaction be R N.


a $R(\rightarrow)$

$$P \cos \theta - 1 = 0$$

$$\therefore P = \frac{1}{\cos \theta}$$

$$= \frac{13}{5}$$

$$P = 2.6$$



7 b $R(\uparrow)$

$$R - P \sin \theta - 2 = 0$$

$$\therefore R = P \sin \theta + 2$$

$$= 2.6 \times \frac{12}{13} + 2$$

$$= 2.4 + 2$$

$$= 4.4$$

8 a Consider the particle of mass $2m$ kg first, as it has only two forces acting on it. This enables you to find the tension.

 $R(\uparrow)$

$$T - 2mg = 0$$

$$\therefore T = 2mg$$

Consider the particle of mass m kg:

 $R(\rightarrow)$

$$T - F = 0$$

$$\therefore F = T = 2mg$$

$$= 19.6m$$

 $R(\uparrow)$

$$R - mg = 0$$

$$\therefore R = mg$$

$$= 9.8m$$

b Let T' be the new tension in the string.

Consider the particle of mass $2m$ kg:

$$R(\uparrow): T' = 2mg$$

Consider the particle of mass m kg:

 $R(\rightarrow)$

$$T' \cos 30^\circ - F' = 0$$

$$\therefore F' = 2mg \times \frac{\sqrt{3}}{2}$$

$$= \sqrt{3}mg$$

$$= 17m \text{ (2 s.f.)}$$



 $R(\uparrow)$

$$R' + T' \sin 30 - mg = 0$$

$$\therefore R' = mg - T' \sin 30$$

$$= mg - 2mg \times \frac{1}{2} \text{ (using } T' = 2mg)$$

$$= 0$$



Mechanics 1

Solution Bank

9 Let the normal reaction be R N.


$$R(\nearrow):$$

$$P - 2g \sin 45^\circ = 0$$

$$\therefore P = 2g \sin 45^\circ$$

$$= g\sqrt{2}$$

$$= 14 \text{ N (2 s.f.)}$$



10 Let the normal reaction be R N.


$$R(\nearrow):$$

$$P \cos 45^\circ - 4g \sin 45^\circ = 0$$

$$\therefore P = \frac{4g \sin 45^\circ}{\cos 45^\circ}$$

$$= 4g$$

$$= 39 \text{ (2 s.f.)}$$



11 a Let the normal reaction between the particle P and the plane be R N.
Let the tension in the string be T N.

Consider first the 5 kg mass.

$$R(\uparrow)$$

$$T - 5g = 0$$

$$\therefore T = 5g$$

Consider the 2 kg mass.


$$R(\nwarrow)$$

$$R - 2g \cos \theta = 0$$

$$R = 2g \times \frac{4}{5}$$

$$= \frac{8g}{5}$$

$$= 16 \text{ N (2 s.f.)}$$



b $R(\nearrow)$

$$T - F - 2g \sin \theta = 0$$

$$F = T - 2g \sin \theta$$

$$= 5g - 2g \times \frac{3}{5} \quad (\text{using } T = 5g \text{ from above})$$

$$= \frac{19g}{5}$$

$$= 37 \text{ N (2 s.f.)}$$

c Assuming the pulley is smooth means there is no friction between it and the string.

12 Let the normal reaction be R N.

First, resolve along the plane to find P as it is the only unknown when resolving in that direction.

R (\nearrow)

$$P \cos 30^\circ - 5 \cos 45^\circ - 20 \sin 45^\circ = 0$$

$$\begin{aligned} \therefore P &= \frac{5 \cos 45^\circ + 20 \sin 45^\circ}{\cos 30^\circ} \\ &= \left(5 \times \frac{\sqrt{2}}{2} + 20 \times \frac{\sqrt{2}}{2} \right) \times \frac{2}{\sqrt{3}} \\ &= \frac{25\sqrt{2}}{\sqrt{3}} \\ &= \frac{25\sqrt{6}}{3} \\ &= 20.4 \text{ (3 s.f.)} \end{aligned}$$

R (\nwarrow)

$$R + P \sin 30^\circ + 5 \sin 45^\circ - 20 \cos 45^\circ = 0$$

$$R = 20 \cos 45^\circ - 5 \sin 45^\circ - P \sin 30^\circ \quad \left(\text{as } P = \frac{25\sqrt{6}}{3} \right)$$

$$\begin{aligned} R &= \frac{15}{\sqrt{2}} - \frac{25\sqrt{6}}{6} \\ &= \frac{45\sqrt{2} - 25\sqrt{6}}{6} \\ &= 0.400 \text{ (3 s.f.)} \end{aligned}$$

