



Mechanics 1 Resolving Forces Questions

- 8 A rough slope is inclined at an angle of 25° to the horizontal. A box of weight 80 newtons is on the slope. A rope is attached to the box and is parallel to the slope. The tension in the rope is of magnitude T newtons. The diagram shows the slope, the box and the rope.



- (a) The box is held in equilibrium by the rope.
- Show that the normal reaction force between the box and the slope is 72.5 newtons, correct to three significant figures. *(3 marks)*
 - The coefficient of friction between the box and the slope is 0.32. Find the magnitude of the maximum value of the frictional force which can act on the box. *(2 marks)*
 - Find the least possible tension in the rope to prevent the box from moving down the slope. *(4 marks)*
 - Find the greatest possible tension in the rope. *(3 marks)*
 - Show that the mass of the box is approximately 8.16 kg. *(1 mark)*
- (b) The rope is now released and the box slides down the slope. Find the acceleration of the box. *(3 marks)*
-


- 2 A particle is in equilibrium under the action of four horizontal forces of magnitudes 5 newtons, 8 newtons, P newtons and Q newtons, as shown in the diagram.



- (a) Show that $P = 9$. *(3 marks)*
- (b) Find the value of Q . *(2 marks)*


-
- 4 A block is being pulled up a rough plane inclined at an angle of 22° to the horizontal by a rope parallel to the plane, as shown in the diagram.

The mass of the block is 0.7 kg , and the tension in the rope is T newtons.



- (a) Draw a diagram to show the forces acting on the block. *(1 mark)*
- (b) Show that the normal reaction force between the block and the plane has magnitude 6.36 newtons, correct to three significant figures. *(3 marks)*
- (c) The coefficient of friction between the block and the plane is 0.25 . Find the magnitude of the frictional force acting on the block during its motion. *(2 marks)*
- (d) The tension in the rope is 5.6 newtons. Find the acceleration of the block. *(4 marks)*
-

- 3 The diagram shows three forces which act in the same plane and are in equilibrium.




- (a) Find F . *(3 marks)*
- (b) Find α . *(3 marks)*
-

- 6 A trolley, of mass 100 kg, rolls at a constant speed along a straight line down a slope inclined at an angle of 4° to the horizontal.

Assume that a constant resistance force, of magnitude P newtons, acts on the trolley as it moves. Model the trolley as a particle.

- (a) Draw a diagram to show the forces acting on the trolley. *(1 mark)*
- (b) Show that $P = 68.4\text{ N}$, correct to three significant figures. *(3 marks)*
- (c) (i) Find the acceleration of the trolley if it rolls down a slope inclined at 5° to the horizontal and experiences the same constant force of magnitude P that you found in part (b). *(4 marks)*
- (ii) Make one criticism of the assumption that the resistance force on the trolley is constant. *(1 mark)*
-

- 3 A sign, of mass 2 kg, is suspended from the ceiling of a supermarket by two light strings. It hangs in equilibrium with each string making an angle of 35° to the vertical, as shown in the diagram. Model the sign as a particle.



- (a) By resolving forces horizontally, show that the tension is the same in each string. *(2 marks)*
- (b) Find the tension in each string. *(5 marks)*
- (c) If the tension in a string exceeds 40 N, the string will break. Find the mass of the heaviest sign that could be suspended as shown in the diagram. *(3 marks)*
-

- 6 A box, of mass 3 kg, is placed on a slope inclined at an angle of 30° to the horizontal. The box slides down the slope. Assume that air resistance can be ignored.
- (a) A simple model assumes that the slope is smooth.
- (i) Draw a diagram to show the forces acting on the box. *(1 mark)*
 - (ii) Show that the acceleration of the box is 4.9 m s^{-2} . *(2 marks)*
- (b) A revised model assumes that the slope is rough. The box slides down the slope from rest, travelling 5 metres in 2 seconds.
- (i) Show that the acceleration of the box is 2.5 m s^{-2} . *(2 marks)*
 - (ii) Find the magnitude of the friction force acting on the box. *(3 marks)*
 - (iii) Find the coefficient of friction between the box and the slope. *(5 marks)*
 - (iv) In reality, air resistance affects the motion of the box. Explain how its acceleration would change if you took this into account. *(2 marks)*

Mechanics 1 Resolving Forces Answers

8(a)(i)	$R = 80 \cos 25^\circ$ $R = 72.5\text{N}$	M1 A1 A1		3	component attempted correct component cao
(ii)	$F = 0.32 \times 72.5$ $F = 23.2\text{N}$	M1 A1		2	condone inequality cao
(iii)	$T + F = 80 \cos 65^\circ$ $T = 10.6\text{N}$	M2 A1			3 forces direction correct, component attempted component
(iv)	$T = F + 80 \cos 65^\circ$ $T = 57.0\text{N}$ (57N)	M1 A1			3 forces, direction correct, component attempted component
(iv)	$\text{Mass} = \frac{80}{g} = (8.16\text{kg})$	B1		1	
(b)	$80 \cos 65^\circ - F = \text{mass} \times \text{acceleration}$	M1			3 terms, component attempted
	$10.6 = \frac{80}{g} \times \text{acc}$ $\text{acc} = 1.30 \text{ m s}^{-2}$ (1.3 m s^{-2})	A1 A1		3	all correct cao
Total				16	

2(a)	$P = 5 + 8 \cos 60^\circ$ $P = 9$	M1 A1 A1		3	Both relevant forces, component of 8N attempted All correct CAO
(b)	$Q = 8 \cos 30^\circ$ $Q = 6.93$ or $4\sqrt{3}$	M1 A1		2	Component of 8N attempted AWRT 6.93
Total				5	

4(a)		B1	1	Accept W or mg (or 6.86) for weight Arrows and labels needed (can replace W with 2 correct components)
(b)	$R = 0.7g \cos 22^\circ$ $R = 6.36 \text{ N}$	M1 A1 A1	3	component of weight attempted all correct, including signs CAO
(c)	$F = 0.25 \times 6.36$ $F = 1.59 \text{ N}$	M1 A1	2	CAO
(d)	$5.6 - 0.7g \sin 22^\circ - 1.59 = 0.7a$ $a = 2.06 \text{ ms}^{-2}$	M1 A2 A1F	4	4 terms with weight component attempted A marks -1 each error, accept $\pm 0.7a$ FT one error, accept \pm
Total			10	

3(a)	$F = \sqrt{6^2 + 5^2}$ $= \sqrt{61} = 7.81$ Alt $\alpha = \tan^{-1}\left(\frac{5}{6}\right) = 39.8^\circ$ $F = \frac{6}{\cos 39.8} = 7.81$ or $F = \frac{5}{\sin 39.8} = 7.81$	M1A1 A1 (M1A1) (A1)	3	Obtaining an equation for F with square or root. Correct equation Correct force Equation for F with a value for α . Correct equation Correct force
(b)	$\alpha = \tan^{-1}\left(\frac{5}{6}\right)$ or $\cos^{-1}\left(\frac{6}{7.81}\right)$ or $\sin^{-1}\left(\frac{6}{7.81}\right)$ $= 39.8^\circ$ Alt $\frac{\sin \alpha}{5} = \frac{\sin 90^\circ}{\sqrt{61}}$ $\alpha = 39.8^\circ$	M1 A1 A1	3	Obtaining an equation for α using trigonometry. Correct equation (using their F) Correct angle Accept values between 39.7 and 39.9
Total			6	

6(a)		B1	1	Correct diagram with arrows and labels Must not use F instead of P Condone resistance instead of P
(b)	$P = 100 \times 9.8 \sin 4^\circ$ $= 68.4$	M1 M1 A1	3	Resolving weight (must see 100) Using $\sin 4^\circ$ or $\cos 86^\circ$ AG Correct P from correct working
(c)	$100a = 100 \times 9.8 \sin 5^\circ - 100 \times 9.8 \sin 4^\circ$ $a = \frac{100 \times 9.8 \sin 5^\circ - 100 \times 9.8 \sin 4^\circ}{100}$ $= 0.171$	M1 A1 A1	4	Three term equation of motion Weight resolved correctly Correct equation
(d)	You would expect P to vary with the speed of the car.	B1	1	Correct explanation
			9	

3(a)	$T_1 \sin 35^\circ = T_2 \sin 35^\circ$ $T_1 = T_2$ OR $T_1 \cos 55^\circ = T_2 \cos 55^\circ$ $T_1 = T_2$	M1 A1	2	Resolving two forces and forming an equation, with different tensions for each string Correct result from correct working
(b)	$T_1 \cos 35^\circ + T_2 \cos 35^\circ = 2 \times 9.8$ $T_1 \cos 35^\circ + T_1 \cos 35^\circ = 2 \times 9.8$ $T_1 = \frac{2 \times 9.8}{2 \cos 35^\circ} = 12.0 \text{ N (to 3sf)}$	M1 A1 A1 dM1 A1	5	Resolving forces to form a three term vertical equation Correct equation T_1 or T_2 eliminated correctly Solving for T_1 or T_2 Correct tension Accept 12 N or 11.9 N
(c)	$2 \times 40 \cos 35^\circ = 9.8m$ $m = \frac{80 \cos 35^\circ}{9.8} = 6.69 \text{ kg}$ OR $m = \frac{40}{11.96} \times 2$ $= 6.69 \text{ kg}$	M1 A1 A1	3	Forming an equation with two tensions to find m Correct equation Correct mass Accept 6.68
Total			10	

6(a)(i)		B1	1	Correct diagram with arrows and labels
(ii)	$3a = 3g \sin 30^\circ$ $a = g \sin 30^\circ = 4.9 \text{ ms}^{-2}$	M1 A1	2	Two term equation of motion AG Correct acceleration from correct working (Allow $a = g \sin 30^\circ$)
(b)(i)	$5 = \frac{1}{2}a \times 2^2$ $a = 2.5 \text{ ms}^{-2}$	M1 A1	2	Constant acceleration equation with $u = 0$ AG Correct answer from correct working. (Use of $v = 5$ must be justified)
(ii)	$3 \times 2.5 = 3g \sin 30^\circ - F$ $F = 3g \sin 30^\circ - 7.5$ $= 7.20 \text{ N (to 3 sf)}$	M1 A1 A1	3	Three term equation of motion Correct equation Correct F Accept 7.2 N
(iii)	$R = 3g \cos 30^\circ (= 25.46)$ $7.2 = \mu \times 3g \cos 30^\circ$ $\mu = \frac{7.2}{3g \cos 30^\circ} = 0.283$	M1 A1 M1 A1F A1F	5	Resolving perpendicular to the slope to find R Correct R Use of $F = \mu R$ Correct expression Correct μ Accept 0.282 (Follow through from incorrect F from above, but not an incorrect R)
(iv)	Reduce a , as the air resistance would reduce the magnitude of the resultant force or because the air resistance increases as the velocity increases towards its terminal value	B1 B1	2	Reduces Explanation Second B1 dependent on the first B1 mark
Total			15	