

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN


JUNE 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6678

Paper No. M2

| Question number | Scheme | Marks |
|-----------------|--|---|
| <p>3.</p> | <p>[Wherever \leq or \geq used in scheme, can be replaced by =]</p>  <p>Resolve \rightarrow: $S = F$ Resolve \uparrow: $R = 6mg$</p> <p>$M(A)$: $S \cdot 2a \cos 30^\circ = mg \sin 30^\circ (a + 5x)$ "F $\leq 0.5 R$" $\Rightarrow S \leq 3mg$</p> <p>$\Rightarrow (a + 5x) \tan 30^\circ \leq 6a, \quad x \leq \frac{(6\sqrt{3}-1)a}{5} \Rightarrow k = \frac{(6\sqrt{3}-1)}{5}$ or 1.88 } or 1.9 }</p> <p>[Alternatives: $M(B)$: $R \cdot 2a \sin 30^\circ = F \cdot 2a \cos 30^\circ + mga \sin 30^\circ + 5mgx \sin 30^\circ$ M1A1A1 $d = 2a - x$ B1; "F $\leq 0.5 R$" $\Rightarrow F \leq 3mg$ M1, rest as scheme. $M(\text{centre})$: $Ra \sin 30^\circ + 5mg(x-a) \sin 30^\circ = (F+S)a \cos 30^\circ$; $S \leq 3mg$ etc. Mark as scheme.]</p> <p>[Note (i): MR - 30° to the ground - gives $k = \frac{(6-\sqrt{3})}{5}$ or 0.493 (ii) The same answer is obtained if only error is sin/cos confusion; both score 7/9. (iii) m used for mg throughout, no penalty; inconsistent, as scheme but max -2]</p> | <p>B1 M1A1 M1A1A1 M1 M1A1 (9)</p> |
| <p>4.</p> | <p>(a) Impulse = change in momentum $3.5 \mathbf{i} + 3 \mathbf{j} = 0.1[(10 \mathbf{i} + 25 \mathbf{j}) - (u \mathbf{i} + v \mathbf{j})]$ Answer: $u \mathbf{i} + v \mathbf{j} = (-25 \mathbf{i} - 5 \mathbf{j}) \text{ ms}^{-1}$</p> <p>(b) Complete method to find height s above hit position Correct equation in s only: $0 = 625 - 2(9.8)s$; $s = 25(25/g) - \frac{1}{2}g(25/g)^2$ Answer: 32.9 m or 33m</p> <p>(c) Method for total time: $0 = 25t - 4.9t^2 \Rightarrow t = 5.10 \text{ s}$ or "half time" $0 = 25 - 9.8t' \Rightarrow t' = 2.55 \text{ s}$ Horizontal distance = $10 \times t = 51 \text{ m}$ [$\sqrt{\text{for } 10t \text{ or } 20t'}$]</p> <p>[Notes: If \mathbf{i} and \mathbf{j} interchanged, then can score Ms in (b) and (c); allow $\sqrt{\text{for } 25 \times 2.04 = 51}$. [Use of answer in (a) can score M marks in (b)(c) only [Use of $\frac{V^2 \sin^2 \theta}{2g}$ and $\frac{V^2 \sin 2\theta}{g}$: M1 method for V or θ, A1 both correct for first two marks]</p> | <p>M1A1 A1 (3) M1 A1 A1 (3) M1A1 M1A1 (4)</p> |

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN

JUNE 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6.678

Paper No. M.2

| Question number | Scheme | Marks |
|-----------------|---|--|
| 5. | <p>(a) Using work/energy equation: (i) P.E. = $\pm 0.5gh$, = $\pm g \sin 20^\circ$; (ii) K.E. = $\frac{1}{2} \times 0.5 \times 25$ $\frac{1}{2} \times 0.5 \times 25 = 0.5 gh + 2R$ Solving for R; $R = 1.45$ or 1.4 [Note: $2(R + 0.5 \times 9.8 \times \sin 20^\circ) = \frac{1}{2} (0.5)25$ scores first 5 marks, mark as scheme]</p> <p><i>Alternative method:</i> Speed equation for a : $0 = 25 \pm 2 a (2)$ ($a = \pm 6.25$) Equation of motion: $(R + 0.5 \times 9.8 \times \sin 20^\circ) = \pm 0.5a$ Totally correct equation: $-(R + 0.5 \times 9.8 \times \sin 20^\circ) = 0.5a$, $a = -ve$ Solving for R</p> <p>(b) Complete method for s [Work/energy equation: $\frac{1}{2} \times 0.5 \times 25 = s R + 0.5 \times 9.8 \times s \sin 40^\circ$ or $-(R + 0.5g \sin 40^\circ) = 0.5a$ ($a = -9.2$) and $0 = 25 + 2as$] Answer: $s = 1.36$ m or 1.4 m</p> | <p>M1,A1;B1 M1A1 M1A1 (7)</p> <p>M1A1 M1A1 A1 M1A1</p> <p>M1A1√ A1 (3)</p> |
| 6. | <p>(a) $\rightarrow v_1$ $\rightarrow v_2$ CoM: $4mu + 4mu = 2m v_1 + 4m v_2$ $\rightarrow 2u$ $\rightarrow u$ $\Rightarrow 4u = v_1 + 2 v_2$ A \circ B \circ $2m$ $4m$ NEL: $\frac{1}{2} (2u - u) = v_2 - v_1$</p> <p>Solving to find v_2; $v_2 = \frac{3u}{2}$</p> <p>(b) Substitute for v_2 in one equation; $v_1 = v_2 - \frac{1}{2} u = u$</p> <p>(c) $\rightarrow w_1$ $\rightarrow w_2$ CoM: $4m(\frac{3}{2}u) = 4m w_1 + m w_2$ $\rightarrow \frac{3}{2}u$ $\rightarrow 0$ $\Rightarrow 6u = 4w_1 + w_2$ \circ B \circ C $4m$ m NEL: $e(\frac{3}{2}u) = w_2 - w_1$</p> <p>Solving for w_1 as $f(e)$: $w_1 = \frac{3u(4-e)}{10}$ or e as $f(w_1)$: $e = \frac{2(6u - 5w_1)}{3u}$</p> <p>Requirement is that $w_1 \geq$ candidate's $v_1 = u$; $\Rightarrow e \leq \frac{2}{3}$ [Note: If w_1 or e not found (not asked for): Setting $w_1 = v = u \Rightarrow w_2 = 2u \Rightarrow e = \frac{2}{3}$ is M1A1 but need to deal with inequality for final M1A1]</p> | <p>M1A1 M1A1 M1A1cso(6) M1A1 (2)</p> <p>M1A1 M1A1 M1A1</p> <p>M1;A1 (8)</p> <p><i>whichever equation used</i></p> |

EDEXCEL FOUNDATION

Stewart House 32 Russell Square London WC1B 5DN

JUNE 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6-678

Paper No. M.2.

| Question number | Scheme | Marks |
|------------------------------------|--|--|
| 7. | <p>(a) $U_y = 23.75 \sin \alpha (= 19)$</p> <p>Complete method to find time, e.g. $-2.4 = 23.75 \sin \alpha t - \frac{1}{2}gt^2$</p> <p>Solving to find t; $t = 4$</p> <p>(b) $\frac{dv}{dt} = -\frac{1}{4}t^2$ $\Rightarrow v = -\frac{1}{12}t^3 + c$</p> <p>$t = 0, v = 18 \Rightarrow v = 18 - \frac{1}{12}t^3$</p> <p>(c) Putting $v = 0$ expression in (b)</p> <p>Solving equation [dependent on previous M1 and M1 in (b)]</p> <p>Finding $T = 6$, with no wrong working seen [Allow verification]</p> <p>(d) Distance \rightarrow travelled by package = $23.75 \cos \alpha \times 4_c = 57$ m [$\sqrt{\quad}$ only on $14.25 \times 4_c$]</p> <p>For lorry $s = 18t - \frac{1}{48}t^4$</p> <p>Showing $s = 66\frac{2}{3}$ for lorry, and distance between them is just under 10m</p> <p>[If lorry moving in direction CA, allow final answer of just under 124m]</p> | <p>B1</p> <p>M1A1</p> <p>M1A1 (5)</p> <p>M1A1</p> <p>A1 (3)</p> <p>M1</p> <p>M1</p> <p>A1 cso (3)</p> <p>M1A1\checkmark</p> <p>M1;A1\checkmark</p> <p>A1 cso (5)</p> |
| <p><i>Geoff Staley 25/6/01</i></p> | | |