

**ADVANCED GCE  
MATHEMATICS**

**4729/01**

Mechanics 2

**MONDAY 16 JUNE 2008**

Afternoon

Time: 1 hour 30 minutes

**Additional materials:** Answer Booklet (8 pages)  
List of Formulae (MF1)

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by  $g \text{ m s}^{-2}$ . Unless otherwise instructed, when a numerical value is needed, use  $g = 9.8$ .
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of **6** printed pages and **2** blank pages.

1 A car is pulled at constant speed along a horizontal straight road by a force of 200 N inclined at  $35^\circ$  to the horizontal. Given that the work done by the force is 5000 J, calculate the distance moved by the car. [3]

2 A bullet of mass 9 grams passes horizontally through a fixed vertical board of thickness 3 cm. The speed of the bullet is reduced from  $250 \text{ m s}^{-1}$  to  $150 \text{ m s}^{-1}$  as it passes through the board. The board exerts a constant resistive force on the bullet. Calculate the magnitude of this resistive force. [4]

3 The resistance to the motion of a car of mass 600 kg is  $k v$  N, where  $v \text{ m s}^{-1}$  is the car's speed and  $k$  is a constant. The car ascends a hill of inclination  $\alpha$ , where  $\sin \alpha = \frac{1}{10}$ . The power exerted by the car's engine is 12 000 W and the car has constant speed  $20 \text{ m s}^{-1}$ .

(i) Show that  $k = 0.6$ . [3]

The power exerted by the car's engine is increased to 16 000 W.

(ii) Calculate the maximum speed of the car while ascending the hill. [3]

The car now travels on horizontal ground and the power remains 16 000 W.

(iii) Calculate the acceleration of the car at an instant when its speed is  $32 \text{ m s}^{-1}$ . [3]

4 A golfer hits a ball from a point  $O$  on horizontal ground with a velocity of  $35 \text{ m s}^{-1}$  at an angle of  $\theta$  above the horizontal. The horizontal range of the ball is  $R$  metres and the time of flight is  $t$  seconds.

(i) Express  $t$  in terms of  $\theta$ , and hence show that  $R = 125 \sin 2\theta$ . [5]

The golfer hits the ball so that it lands 110 m from  $O$ .

(ii) Calculate the two possible values of  $t$ . [5]




Fig. 1

A toy is constructed by attaching a small ball of mass 0.01 kg to one end of a uniform rod of length 10 cm whose other end is attached to the centre of the plane face of a uniform solid hemisphere with radius 3 cm. The rod has mass 0.02 kg, the hemisphere has mass 0.5 kg and the rod is perpendicular to the plane face of the hemisphere (see Fig. 1).

- (i) Show that the distance from the ball to the centre of mass of the toy is 10.7 cm, correct to 1 decimal place. [4]

(ii)





Fig. 2

The toy lies on horizontal ground in a position such that the ball is touching the ground (see Fig. 2). Determine whether the toy is lying in equilibrium or whether it will move to a position where the rod is vertical. [4]

6



A particle  $P$  of mass  $0.5\text{ kg}$  is attached to points  $A$  and  $B$  on a fixed vertical axis by two light inextensible strings of equal length. Both strings are taut and each is inclined at  $60^\circ$  to the vertical (see diagram). The particle moves with constant speed  $3\text{ m s}^{-1}$  in a horizontal circle of radius  $0.4\text{ m}$ .

- (i) Calculate the tensions in the two strings. [7]

The particle now moves with constant angular speed  $\omega\text{ rad s}^{-1}$  and the string  $BP$  is on the point of becoming slack.

- (ii) Calculate  $\omega$ . [5]

7



Two small spheres  $A$  and  $B$  of masses  $2\text{ kg}$  and  $3\text{ kg}$  respectively lie at rest on a smooth horizontal platform which is fixed at a height of  $4\text{ m}$  above horizontal ground (see diagram). Sphere  $A$  is given an impulse of  $6\text{ N s}$  towards  $B$ , and  $A$  then strikes  $B$  directly. The coefficient of restitution between  $A$  and  $B$  is  $\frac{2}{3}$ .

- (i) Show that the speed of  $B$  after it has been hit by  $A$  is  $2\text{ m s}^{-1}$ . [6]

Sphere  $B$  leaves the platform and follows the path of a projectile.

- (ii) Calculate the speed and direction of motion of  $B$  at the instant when it hits the ground. [7]

8 (i)




Fig. 1

A uniform lamina  $ABCD$  is in the form of a right-angled trapezium.  $AB = 6$  cm,  $BC = 8$  cm and  $AD = 17$  cm (see Fig. 1). Taking  $x$ - and  $y$ -axes along  $AD$  and  $AB$  respectively, find the coordinates of the centre of mass of the lamina. [8]

(ii)




Fig. 2

The lamina is smoothly pivoted at  $A$  and it rests in a vertical plane in equilibrium against a fixed smooth block of height 7 cm. The mass of the lamina is 3 kg.  $AD$  makes an angle of  $30^\circ$  with the horizontal (see Fig. 2). Calculate the magnitude of the force which the block exerts on the lamina. [5]