Solution Bank



Exercise 2A

- **1** a $s = 9(1) 1^3 = 8$ m
 - **b** $9t t^3 = 0$ $t(9 - t^2) = 0$ Either t = 0 or $t^2 = 9$ $\Rightarrow t = 0$ or $t = \pm 3$
- 2 a At t = 2, $s = 5(2)^2 - 2^3 = 12$ At t = 4, $s = 5(4)^2 - 4^3 = 16$ Change in displacement = 16 - 12 = 4 m
 - **b** At t = 3, $s = 5(3)^2 - 3^3 = 18$ Change in displacement in the third second = 18 - 12 = 6 m
- **3** a $v = 3 + 5(1) 1^2 = 7 \text{ m s}^{-1}$



At t = 0, v = 3 v = 3 again when $5t - t^2 = 0 \Rightarrow t = 5$ Using symmetry, turning point is when t = 2.5. When t = 2.5, $v = 3 + 5(2.5) - 2.5^2 = 9.25$ So in $0 \le t \le 4$, range of v is $3 \le v \le 9.25$ Greatest speed is 9.25 m s⁻¹.

c $v = 3 + 5(7) - 7^2$ = 3 + 35 - 49

$$= -11$$

When t = 7, the velocity of the particle is -11 m s^{-1} . This means it is moving in the opposite direction to that in which it was initially travelling.

Solution Bank



4 **a** s = 0 when $\frac{1}{5}(4t - t^2) = 0$ $\frac{1}{5}t(4 - t) = 0$ $\Rightarrow t = 0$ or t = 4By symmetry, maximum displacement is when t = 2. When t = 2, $s = \frac{1}{5}(4(2) - 2^2)$ $= \frac{4}{5}$

The maximum displacement is 0.8 m.

b When the toy car returns to *P*, s = 0 $\frac{1}{5} (4t - t^2) = 0$ $\frac{1}{5}t(4 - t) = 0$ $\Rightarrow t = 0 \text{ or } t = 4$

The toy car returns to P after 4 s.

- **c** The toy car travels to maximum distance and back again. So total distance = 0.8 + 0.8 = 1.6 m
- **d** The model is valid for $0 \le t \le 4$.
- 5 a When t = 0, v = 0 - 0 + 8 = 8The initial velocity is 8 m s⁻¹.
 - **b** $3t^2 10t + 8 = 0$ (3t - 4)(t - 2) = 0The body is at rest when $t = \frac{4}{3}$ and t = 2.
 - c $3t^2 10t + 8 = 5$ $3t^2 - 10t + 3 = 0$ (3t - 1)(t - 3) = 0Velocity = 5 m s⁻¹ when t = $\frac{1}{3}$ and t = 3.

Solution Bank





Using the answer to part b and symmetry, the body has its maximum/minimum velocity when $t = \frac{5}{3}$ s.

When
$$t = \frac{5}{3}$$
,
 $v = 3\left(\frac{5}{3}\right)^2 - 10\left(\frac{5}{3}\right) + 8$
 $= \frac{25}{3} - \frac{50}{3} + 8$
 $= -\frac{25}{3} + 8$
 $v = -\frac{1}{3}$

So in $0 \le t \le 2$, range of v is $-\frac{1}{3} \le v \le 8$. Greatest speed is 8 m s⁻¹.

6 a $8t - 2t^2 = 0$ 2t(4 - t) = 0

The particle is next at rest after 4 s.

b By symmetry, minimum/maximum velocity is when t = 2. When t = 2, $v = 8(2) - 2(2)^2$ = 8So in $0 \le t \le 4$, greatest speed is 8 m s⁻¹.

Solution Bank



7 $s = 3t^2 - t^3$

Model is valid until particle returns to starting point, i.e. until next point at which s = 0. After this it would have a negative displacement, i.e. be beyond *O*. s = 0 when

 $3T^2 - T^3 = 0$ $T^2(3 - T) = 0$ $T \neq 0$ so T = 3

- 8 a $\frac{1}{5}(3t^2 10t + 3) = 0$ $3t^2 - 10t + 3 = 0$ (3t - 1)(t - 3) = 0Particle is at rest when $t = \frac{1}{3}$ and t = 3.
 - **b** Using answer to part **a** and symmetry, the body has its maximum/minimum velocity when $t = \frac{5}{3}$. When $t = \frac{5}{2}$,

$$v = \frac{1}{5} \left(3 \left(\frac{5}{3} \right)^2 - 10 \left(\frac{5}{3} \right) + 3 \right)$$
$$= \frac{1}{5} \left(\frac{25}{3} - \frac{50}{3} + \frac{9}{3} \right)$$
$$= \frac{1}{5} \left(-\frac{16}{3} \right)$$
$$= -\frac{16}{15}$$

So in $0 \le t \le 3$, greatest speed is $\frac{16}{15}$ m s⁻¹.