## **Mechanics 2**

#### Solution Bank



#### **Exercise 2F**

- 1  $v = \int a dt$ = at + c, where c is a constant of integration.
  - When t = 0, v = 0  $0 = a \times 0 + c \Rightarrow c = 0$  v = at  $s = \int v dt$   $= \int at dt$  $= \frac{1}{2}at^2 + k$ , where k is a constant of integration.
  - When t = 0, s = x  $x = \frac{1}{2} \times a \times 0^2 + k \Rightarrow k = x$  $s = \frac{1}{2}at^2 + x$
- 2 a  $v = \int a dt$ =  $\int 5 dt$ = 5t + c, where c is a constant of integration.
  - When t = 0, v = 12  $12 = 0 + c \Rightarrow c = 12$ v = 12 + 5t
  - **b**  $s = \int v dt$ =  $\int (12+5t) dt$ =  $12t + \frac{5}{2}t^2 + k$ , where k is a constant of integration.
    - When t = 0, s = 7  $7 = 0 + 0 + k \Rightarrow k = 7$   $s = 12t + \frac{5}{2}at^2 + 7$  $= 12t + 2.5t^2 + 7$
- 3  $s = ut + \frac{1}{2}at^2$   $v = \frac{ds}{dt} = u + at$  $a = \frac{dv}{dt} = a$

So acceleration is constant.

# **Mechanics 2**

# Solution Bank



4 A:  $s = 2t^2 - t^3$   $v = \frac{ds}{dt} = 4t - 3t^2$   $a = \frac{dv}{dt} = 4 - 6t$ Not constant

$$\mathbf{B}: s = 4t + 7$$
$$v = \frac{\mathrm{d}s}{\mathrm{d}t} = 4$$
$$a = \frac{\mathrm{d}v}{\mathrm{d}t} = 0$$

No acceleration (i.e. constant acceleration of zero)

C: 
$$s = \frac{t^2}{4}$$
  
 $v = \frac{ds}{dt} = \frac{t}{2}$   
 $a = \frac{dv}{dt} = \frac{1}{2}$ 

Constant acceleration

D: 
$$s = 3t - \frac{2}{t^2}$$
  
 $v = \frac{ds}{dt} = 3 + \frac{4}{t^3}$   
 $a = \frac{dv}{dt} = -\frac{12}{t^4}$ 

Not constant

$$\mathbf{E}: s = 6$$
$$v = \frac{\mathrm{d}s}{\mathrm{d}t} = 0$$

Particle stationary (i.e. constant acceleration of zero)

5 a 
$$v = u + at$$
  
 $u = 5, v = 13, t = 2$   
 $13 = 5 + 2a$   
 $a = \frac{13-5}{2} = 4$ 

The acceleration of the particle is  $4 \text{ m s}^{-2}$ .

## **Mechanics 2**

#### Solution Bank



5 b  $v = \int a dt$   $= \int 4 dt$  = 4t + c, where c is a constant of integration. When t = 0, v = 5  $5 = 0 + c \Rightarrow c = 5$  v = 4t + 5 $s = \int v dt$ 

 $= \int (4t+5) dt$ =  $2t^2 + 5t + k$ , where k is a constant of integration.

When t = 0, s = 0  $0 = 0 + 0 + k \Rightarrow k = 0$  $s = 2t^2 + 5t$ 

This is an equation of the required form with p = 2, q = 5 and r = 0.

6 a  $s = 25t - 0.2t^2$ When t = 40,  $s = 25 \times 40 - 0.2 \times 40^2$ = 680

The distance AB is 680 m.

**b** 
$$v = \frac{ds}{dt} = 25 - 0.4t$$
  
 $a = \frac{dv}{dt} = -0.4$ 

The train has a constant acceleration (of  $-0.4 \text{ m s}^{-2}$ ).

#### **INTERNATIONAL A LEVEL**

## **Mechanics 2**

#### Solution Bank



6 c Taking the direction in which the train travels to be positive: For the bird: a = -0.6, u = -7, initial displacement = 680

 $v_B = \int a dt$ =  $\int -0.6 dt$ = -0.6t + c, where c is a constant of integration.

When t = 0,  $v_B = -7$   $-7 = 0 + c \Rightarrow c = -7$  v = -0.6t - 7  $s_B = \int v_B dt$   $= \int (-0.6t - 7) dt$  $= -0.3t^2 - 7t + k$ , where k is a constant of integration.

When t = 0,  $s_B = 680$   $680 = 0 - 0 + k \Rightarrow k = 680$  $s_B = -0.3t^2 - 7t + 680$ 

When the bird is directly above the train, the displacement of both train and bird are the same.

 $25t - 0.2t^{2} = -0.3t^{2} - 7t + 680$   $0.1t^{2} + 32t - 680 = 0$   $t^{2} + 320t - 6800 = 0$  (t - 20)(t + 340) = 0 t > 0, so t = 20When t = 20,  $s = 25 \times 20 - 0.2 \times 20^{2}$ = 420

The bird is directly above the train 420 m from A.