


Exercise 3F

1 a



Let O be the origin and OA be the positive x -axis.


$$\begin{aligned}
 7 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} &= 5 \begin{pmatrix} 2.5 \\ 0.5 \end{pmatrix} + 2 \begin{pmatrix} 2.5 \\ 2 \end{pmatrix} \\
 &= \frac{1}{7} \begin{pmatrix} 17.5 \\ 6.5 \end{pmatrix} \\
 &= \begin{pmatrix} \frac{5}{2} \\ \frac{13}{14} \end{pmatrix}
 \end{aligned}$$

When the lamina is suspended from O let the angle between OA and the downward vertical be θ .

$$\tan \theta = \frac{\frac{13}{14}}{\frac{5}{2}}$$

$$\begin{aligned}
 \theta &= 20.376\dots \\
 &= 20.4^\circ \text{ (3 s.f.)}
 \end{aligned}$$

1 b



Let O be the origin and OA be the positive x -axis.

$$12 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 10 \begin{pmatrix} 2.5 \\ 1 \end{pmatrix} + 2 \begin{pmatrix} 4 \\ 2.5 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{12} \begin{pmatrix} 33 \\ 15 \end{pmatrix}$$


$$= \begin{pmatrix} \frac{33}{12} \\ \frac{5}{4} \end{pmatrix}$$

When the lamina is suspended from O let the angle between OA and the downward vertical be θ .

$$\tan \theta = \frac{\frac{5}{4}}{\frac{33}{12}}$$

$$\begin{aligned} \theta &= 24.443\dots \\ &= 24.4^\circ \text{ (3 s.f.)} \end{aligned}$$

1 c



Let O be the origin and OA be the positive x -axis.

$$10 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 4 \begin{pmatrix} 0.5 \\ 2 \end{pmatrix} + 6 \begin{pmatrix} 2.5 \\ 3 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{10} \begin{pmatrix} 17 \\ 26 \end{pmatrix}$$

$$= \begin{pmatrix} \frac{17}{10} \\ \frac{13}{5} \end{pmatrix}$$


When the lamina is suspended from O let the angle between OA and the downward vertical be θ .

$$\tan \theta = \frac{\frac{13}{5}}{\frac{17}{10}}$$

$$\theta = 56.821\dots$$

$$= 56.8^\circ \text{ (3 s.f.)}$$

2



Let A be the origin

$$13 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 2 \begin{pmatrix} 1.5 \\ 1 \end{pmatrix} + 8 \begin{pmatrix} 2 \\ -1 \end{pmatrix} + 3 \begin{pmatrix} 2.5 \\ -2.5 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{13} \begin{pmatrix} 26.5 \\ -13.5 \end{pmatrix}$$

$$= \begin{pmatrix} \frac{53}{26} \\ -\frac{27}{26} \end{pmatrix}$$

When the lamina is suspended from A let the angle between AB and the downward vertical be θ .


$$\tan \theta = \frac{\frac{53}{26}}{\frac{27}{26}}$$

$$\theta = 63.004\dots$$


$$= 63.0^\circ \text{ (3 s.f.)}$$

- 3 Angle between AB and downward vertical = $63.004\dots^\circ$ (from Q2)
 Therefore, angle between AB and horizontal = $90 - 63.004\dots = 27.0^\circ$ (3 s.f)

4



Let P be the origin and PQ be the positive x -axis.
Let the point directly below R that lies on PQ be T .



The total area of the shape is:

$$(4 \times 6) + \frac{1}{2}(6 \times 6) = 42$$

Let PQ be the origin.

$$42 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 24 \begin{pmatrix} 2 \\ 3 \end{pmatrix} + 18 \begin{pmatrix} 6 \\ 2 \end{pmatrix}$$


$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{42} \begin{pmatrix} 156 \\ 108 \end{pmatrix}$$

$$= \begin{pmatrix} \frac{26}{7} \\ \frac{18}{7} \end{pmatrix}$$

a Distance from PS is $\frac{26}{7}$ cm

b Distance from PQ is $\frac{18}{7}$ cm

4 c




$$\tan \theta = \frac{\frac{18}{7}}{10 - \frac{26}{7}}$$

$$\theta = 22.249\dots$$

$$\theta = 22.2^\circ \text{ (3 s.f.)}$$


- 5 Let A be the origin and AF be the positive x -axis.



$$1.2M \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 0.5M \begin{pmatrix} 1 \\ 3 \end{pmatrix} + 0.5M \begin{pmatrix} 5 \\ 1 \end{pmatrix} + 0.2M \begin{pmatrix} 8 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{1.2} \begin{pmatrix} 4.6 \\ 2 \end{pmatrix}$$

$$= \begin{pmatrix} \frac{23}{6} \\ \frac{5}{3} \end{pmatrix}$$




$$\tan \theta = \frac{\frac{11}{6}}{\frac{13}{3}}$$

$$\theta = 22.932\dots\dots$$

$$\theta = 22.9^\circ \text{ (3 s.f.)}$$

6




Let F be the origin and FE be the positive x -axis.

$$16 \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = 12 \begin{pmatrix} 1 \\ 3 \end{pmatrix} + 4 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \frac{1}{16} \begin{pmatrix} 24 \\ 40 \end{pmatrix}$$

$$= \begin{pmatrix} \frac{3}{2} \\ \frac{5}{2} \end{pmatrix}$$


- a Therefore centre of mass lies 1.5 cm from FA
- b Therefore centre of mass lies 2.5 cm from EF
- c



$$\tan \theta = \frac{1.5}{2.5}$$

$$\begin{aligned} \theta &= 30.963\dots \\ &= 31.0^\circ \text{ (3 s.f.)} \end{aligned}$$

7




Let C be the origin and CB be the positive x -axis.

The centre of gravity of the lamina lies at the point $(1, 2)$

$$1.2M \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = M \begin{pmatrix} 1 \\ 2 \end{pmatrix} + 0.2M \begin{pmatrix} 0 \\ 6 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \begin{pmatrix} \frac{5}{6} \\ \frac{8}{3} \end{pmatrix}$$




$$\tan \theta = \frac{\frac{5}{6}}{\frac{8}{3}}$$

$$\theta = 17.354\dots$$

$$= 17.4^\circ \text{ (3 s.f.)}$$

8




Let B be the origin.

The centre of gravity of the lamina lies at the point $\left(0, -\frac{2r \sin \alpha}{3\alpha}\right)$

$\alpha = \frac{\pi}{2}$ and $r = 4$ therefore:

$$\begin{aligned} \frac{2r \sin \alpha}{3\alpha} &= \frac{2 \times 4 \times \sin\left(\frac{\pi}{2}\right)}{3 \times \frac{\pi}{2}} \\ &= \frac{16}{3\pi} \end{aligned}$$

Therefore the centre of gravity of the lamina lies at the point $\left(0, -\frac{16}{3\pi}\right)$




$$\sin 45 = \frac{l}{4}$$

$$l = 2\sqrt{2}$$

Therefore C lies at the point $(2\sqrt{2}, -2\sqrt{2})$

8 (continued)


$$\begin{aligned}
 1.5M \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} &= M \begin{pmatrix} 0 \\ -\frac{16}{3\pi} \end{pmatrix} + 0.5M \begin{pmatrix} 2\sqrt{2} \\ -2\sqrt{2} \end{pmatrix} \\
 &= \frac{1}{1.5} \begin{pmatrix} \sqrt{2} \\ -\frac{16}{3\pi} - \sqrt{2} \end{pmatrix} \\
 &= \begin{pmatrix} \frac{2\sqrt{2}}{3} \\ -\frac{32}{9\pi} - \frac{2\sqrt{2}}{3} \end{pmatrix} \\
 &= \begin{pmatrix} 0.9428\dots \\ 2.0745\dots \end{pmatrix}
 \end{aligned}$$



$$\begin{aligned}
 \tan \theta &= \frac{0.9428\dots}{2.0745\dots} \\
 \theta &= 24.4398\dots
 \end{aligned}$$

Therefore the angle made with AB is $45 + 24.4398\dots = 69.4^\circ$ (3 s.f)

Challenge



Let C be the origin and CD be the positive x -axis.

The centre of gravity of the lamina lies at the point $(2, 3)$

A lies at the point $(0, 6)$.

$$(M + kM) \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = M \begin{pmatrix} 2 \\ 3 \end{pmatrix} + kM \begin{pmatrix} 0 \\ 6 \end{pmatrix}$$

$$(1 + k) \begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} + k \begin{pmatrix} 0 \\ 6 \end{pmatrix}$$

$$\begin{pmatrix} \bar{x} \\ \bar{y} \end{pmatrix} = \begin{pmatrix} \frac{2}{1+k} \\ \frac{3+6k}{1+k} \end{pmatrix}$$

$$\tan 30 = \frac{\frac{2}{1+k}}{\frac{3+6k}{1+k}}$$

$$\sqrt{3}(3+6k) = 6$$

$$3+6k = \frac{6}{\sqrt{3}}$$

$$k = \frac{1}{\sqrt{3}} - \frac{1}{2} \text{ as required}$$