

EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN

June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

Paper No. M3

Question number	Scheme	Marks
1. (a)	$v = \int \frac{1}{2} e^{-\frac{1}{6}t} dt$ $= -3e^{-\frac{1}{6}t} (+c)$ <p>use of limits or $t=0, v=10$</p> $v = 13 - 3e^{-\frac{1}{6}t}$	\rightarrow M1 \leftarrow A1 M1 A1 (4)
(b)	$t=3, v = 11.2 \text{ ms}^{-1}$	M1 A1 (2)
(c)	13 (ft. if $v = a \pm b e^{-\frac{1}{6}t}$)	B1 + (1) (7)
2. (a)	$\cos \theta = \frac{3}{4}, 0.75, 6/8$	B1 (1)
(b)	$mg \cos \theta (-R) = \frac{mv^2}{0.8}$ $v^2 = 5.88 \text{ m}^2 \text{ s}^{-2}$	M1 A1 A1 (3)
(c)	$\frac{1}{2} m \cdot 5.88 - \frac{1}{2} mu^2 = mg \times 0.2$ $u = 1.4$	M1 A1 A1 (3) (7)
3. (a)	$\frac{1}{2} \times 1.5 v^2 = \frac{52 \times 0.05^2}{2 \times 0.25}$ $v = 0.589 \text{ ms}^{-1} \text{ (3SF)}$	\rightarrow M1 A1 A1 M1 A1 (5)
(b)	$F = 0.6 \times 1.5g$ $\frac{52x}{0.25} \text{ or } \frac{52x}{25}$ $T = F \Rightarrow x = 0.0424 \text{ m or } 4.24 \text{ cm}$ $\text{Min distance} = 0.208 \text{ m or } 20.8 \text{ cm}$	M1 B1 M1 A1 A1 + (5) (10)

EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN


June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

Paper No. M3

Question number	Scheme	Marks
A.(a)	$g = k/R^2 \Rightarrow k = R^2g$ $a = -\frac{k}{x^2}$ $\frac{v dv}{dx} = -\frac{R^2g}{x^2}$ $\int v dv = -\int \frac{R^2g}{x^2} dx$ $\frac{v^2}{2} = \frac{R^2g}{x} (+ c)$ <p style="text-align: center;"><small>Correct</small></p> $x=R, v=U \text{ or use of limits}$ $\frac{v^2}{2} = \frac{R^2g}{x} + \frac{U^2}{2} - Rg$ <p style="text-align: center;"><small>use of v=0</small></p> $X = \frac{2gR^2}{2gR - U^2}$	<p style="text-align: center;">B1</p> <p style="text-align: center;">→ M1 [M1 A1 c.s.o. (4)</p> <hr/> <p style="text-align: center;">→ M1 A1 → M1 A1 M1 A1 (6) (10)</p>
S.(a)	$\frac{\pi r^2 h}{6} \quad \frac{1}{6} \pi r^2 h \quad \frac{5}{6} \pi r^2 h$ <p style="text-align: center;"><small>(6) (1) (5)</small></p> $\frac{1}{2}h \quad \frac{7h}{8} \quad \bar{x}$ $6 \cdot \frac{1}{2}h - \frac{7h}{8} = 5\bar{x}$ $\bar{x} = \frac{17h}{40}$	<p style="text-align: center;">B2 -1e.e.o.o. B2 -1e.e.o.o. M1 A1 A1 (7)</p> <hr/> <p style="text-align: center;">→ M1 A1 M1 A1 (4) (11)</p>
(b)	 <p style="text-align: center;">$\tan \alpha = \frac{h - \bar{x}}{r}$</p> <p style="text-align: center;"><small>use of h=4r to obtain expression in h or only</small></p> <p style="text-align: center;">$\alpha = 66.5^\circ$ (1DP)</p>	

EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN


June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6679

Paper No. M3

Question number	Scheme	Marks
6. (a)	$r = \frac{1}{2} h \tan 60^\circ = \frac{\sqrt{3}h}{2} *$	M1 A1 (2)
6. (b)	 <p> $R(\uparrow), T_1 \cos 60^\circ - T_2 \cos 60^\circ = mg$ $R(\leftarrow), T_1 \sin 60^\circ + T_2 \sin 60^\circ = m \frac{\sqrt{3}}{2} h \omega^2$ Use of $\cos 60^\circ = \frac{1}{2}$ and $\sin 60^\circ = \frac{\sqrt{3}}{2}$ solving for T_1 or T_2 $T_1 = \frac{1}{2} m (h \omega^2 + 2g); T_2 = \frac{1}{2} m (h \omega^2 - 2g)$ </p>	\rightarrow M1 A1 \rightarrow M1 A1 B1 M1 A1; A1 = (9)
6. (c)	$T_2 > 0 \Rightarrow \omega > \sqrt{\frac{2g}{h}}$ $T = \frac{2\pi}{\omega} \Rightarrow T < 2\pi \sqrt{\frac{h}{2g}} = \frac{\pi \sqrt{2h}}{\sqrt{g}} *$	M1 A1 ✓ M1 A1 c.s.c. (4) (14)
7. (a)	In equilibrium, $T = mg \sin 30^\circ$ $\lambda \frac{1}{8} a = mg \sin 30^\circ \Rightarrow \lambda = 4mg/x$	B1 M1 A1 (3)
7. (b)	$m \ddot{x} = mg \sin 30^\circ - \frac{4mg}{a} \left(\frac{1}{8} a + x \right)$ $\ddot{x} = -\frac{4g}{a} x \Rightarrow \text{SHM}$ Period = $2\pi \sqrt{\frac{a}{4g}} = \frac{\pi \sqrt{a}}{\sqrt{g}} *$	\rightarrow M1 A2 \rightarrow M1 A1 A1 (6)
7. (c)	Max accel = $\omega^2 a = \frac{4g}{a} \cdot \frac{a}{4} = g$	M1 A1 (2)
7. (d)	$x = \frac{a}{4} \sin \omega t; \frac{a}{8} = \frac{a}{4} \sin \omega t$ $\omega t = \sin^{-1} \frac{1}{2} = \frac{\pi}{6}$ $t = \frac{\pi}{2\omega} \sqrt{\frac{a}{g}}$	\rightarrow M1 A1 \rightarrow M1 A1 A1 ✓ (5)
OR:	Circle approach: $\theta = \frac{\pi}{2} - \cos^{-1} \frac{1}{2} = \frac{\pi}{2} - \frac{\pi}{3} = \frac{\pi}{6}$ $\omega t = \frac{\pi}{6}$ $t = \frac{\pi}{2\omega} \sqrt{\frac{a}{g}}$	OR \rightarrow M1 A1 \rightarrow M1 A1 A1 ✓ (5)
OR:	$\cos^{-1} \left(\frac{1}{2} \right) - \cos^{-1} (0) = \frac{\pi}{3} - \frac{\pi}{2} = \frac{\pi}{6}$ $\omega t = \frac{\pi}{6}$ $t = \frac{\pi}{2\omega} \sqrt{\frac{a}{g}}$	OR \rightarrow M1 A1 \rightarrow M1 A1 A1 ✓ (5) (16)

Question	Solution	Markscheme
7.(d)	$\theta = \frac{\pi}{2} - \cos^{-1} \frac{1}{2} = \frac{\pi}{6}$ $\omega t = \frac{\pi}{6}$ $t = \frac{\pi}{6} \sqrt{\frac{a}{Ag}} = \frac{\pi}{12} \sqrt{\frac{a}{g}}$	$\left[\begin{array}{l} M1 A1 \\ M1 A1 \end{array} \right]$ <p>A1 f.t.</p>
<u>OR:</u>	$\cos^{-1} \left(-\frac{1}{2}\right) - \cos^{-1}(0) = \frac{2\pi}{3} - \frac{\pi}{2} = \frac{\pi}{6}$ $\omega t = \frac{\pi}{6}$ $t = \frac{\pi}{6} \sqrt{\frac{a}{Ag}} = \frac{\pi}{12} \sqrt{\frac{a}{g}}$	$\left[\begin{array}{l} M1 A1 \\ M1 A1 \end{array} \right]$ <p>A1 f.t.</p>