



**ADVANCED GCE UNIT
MATHEMATICS (MEI)**

Mechanics 3

MONDAY 21 MAY 2007

4763/01

Morning
Time: 1 hour 30 minutes

Additional materials:
Answer booklet (8 pages)
Graph paper
MEI Examination Formulae and Tables (MF2)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.

ADVICE TO CANDIDATES

- Read each question carefully and make sure you know what you have to do before starting your answer.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.

This document consists of **6** printed pages and **2** blank pages.

2

- 1 (a) (i) Write down the dimensions of the following quantities.

Velocity

Acceleration

Force

Density (which is mass per unit volume)

Pressure (which is force per unit area) [5]

For a fluid with constant density ρ , the velocity v , pressure P and height h at points on a streamline are related by Bernoulli's equation

$$P + \frac{1}{2}\rho v^2 + \rho gh = \text{constant},$$

where g is the acceleration due to gravity.

- (ii) Show that the left-hand side of Bernoulli's equation is dimensionally consistent. [4]
- (b) In a wave tank, a float is performing simple harmonic motion with period 3.49 s in a vertical line. The height of the float above the bottom of the tank is h m at a time t s. When $t = 0$, the height has its maximum value. The value of h varies between 1.6 and 2.2.
- (i) Sketch a graph showing how h varies with t . [2]
- (ii) Express h in terms of t . [4]
- (iii) Find the magnitude and direction of the acceleration of the float when $h = 1.7$. [3]

3

- 2 A fixed hollow sphere with centre O has an inside radius of 2.7 m . A particle P of mass 0.4 kg moves on the smooth inside surface of the sphere.

At first, P is moving in a horizontal circle with constant speed, and OP makes a constant angle of 60° with the vertical (see Fig. 2.1).




Fig. 2.1

- (i) Find the normal reaction acting on P . [2]
- (ii) Find the speed of P . [4]

The particle P is now placed at the lowest point of the sphere and is given an initial horizontal speed of 9 m s^{-1} . It then moves in part of a vertical circle. When OP makes an angle θ with the upward vertical and P is still in contact with the sphere, the speed of P is $v\text{ m s}^{-1}$ and the normal reaction acting on P is $R\text{ N}$ (see Fig. 2.2).




Fig. 2.2

- (iii) Find v^2 in terms of θ . [3]
- (iv) Show that $R = 4.16 - 11.76 \cos \theta$. [5]
- (v) Find the speed of P at the instant when it leaves the surface of the sphere. [4]

4

3 A light elastic string has natural length 1.2 m and stiffness 637 N m^{-1} .

- (i) The string is stretched to a length of 1.3 m. Find the tension in the string and the elastic energy stored in the string. [3]

One end of this string is attached to a fixed point A. The other end is attached to a heavy ring R which is free to move along a smooth vertical wire. The shortest distance from A to the wire is 1.2 m (see Fig. 3).




Fig. 3

The ring is in equilibrium when the length of the string AR is 1.3 m.

- (ii) Show that the mass of the ring is 2.5 kg. [4]

The ring is given an initial speed $u \text{ m s}^{-1}$ vertically downwards from its equilibrium position. It first comes to rest, instantaneously, in the position where the length of AR is 1.5 m.

- (iii) Find u . [7]
- (iv) Determine whether the ring will rise above the level of A. [4]

5

- 4 (a) The region bounded by the curve $y = x^3$ for $0 \leq x \leq 2$, the x -axis and the line $x = 2$, is occupied by a uniform lamina. Find the coordinates of the centre of mass of this lamina. [8]
- (b) The region bounded by the circular arc $y = \sqrt{4 - x^2}$ for $1 \leq x \leq 2$, the x -axis and the line $x = 1$, is rotated through 2π radians about the x -axis to form a uniform solid of revolution, as shown in Fig. 4.1.




Fig. 4.1

- (i) Show that the x -coordinate of the centre of mass of this solid of revolution is 1.35. [6]

This solid is placed on a rough horizontal surface, with its flat face in a vertical plane. It is held in equilibrium by a light horizontal string attached to its highest point and perpendicular to its flat face, as shown in Fig. 4.2.




Fig. 4.2

- (ii) Find the least possible coefficient of friction between the solid and the horizontal surface. [4]