

GCE Examinations
Advanced Subsidiary / Advanced Level

Mechanics
Module M3

Paper D

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.




Written by Shaun Armstrong & Chris Huffer

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

M3 Paper D – Marking Guide

1. (a) $T = \frac{\lambda x}{l} \therefore 4.5 = \frac{0.06\lambda}{0.15}$ M1 A1
giving $\lambda = 11.25 \text{ N}$ A1
- (b) work done = change in EPE = $\frac{\lambda}{2l}(x_2^2 - x_1^2)$ M1 A1
 $= \frac{11.25}{2 \times 0.15}(0.1^2 - 0.06^2) = 0.24 \text{ J}$ M1 A1 (7)
-
2. (a) particle B1
- (b) minimum speed when bead is at highest point B1
- 

con. of ME: $\frac{1}{2} m(u^2 - v^2) = mg \times 1.6$ M1 A1

$v = \frac{3}{5} u$ M1

$\therefore u^2 - \frac{9}{25} u^2 = 3.2g$ M1

$u^2 = 3.2 \times 9.8 \div \frac{16}{25} = 49$ so $u = 7$ A1 (7)
-
3. (a) $v = \int \frac{4}{(1+t)^3} dt \therefore v = \frac{-2}{(1+t)^2} + c$ M1 A1
 $t = 0, v = 1 \therefore c = 3$ M1
giving $v = [3 - \frac{2}{(1+t)^2}] \text{ ms}^{-1}$ A1
- (b) $x = \int 3 - \frac{2}{(1+t)^2} dt \therefore x = 3t + \frac{2}{(1+t)} + d$ M1 A1
 $t = 0, x = 3 \therefore d = 1$ M1
giving $x = 3t + \frac{2}{(1+t)} + 1$
 $t = 3, x = 9 + \frac{1}{2} + 1 = 10.5 \text{ m}$ M1 A1 (9)
-
4. (a) (i) period = $2 \times 3 = 6 \text{ s}$ B1
(ii) amplitude = $\frac{1}{2} \times 4 = 2 \text{ m}$ B1
- (b) period = $6 = \frac{2\pi}{\omega} \therefore \omega = \frac{\pi}{3}$ M1
 $v_{\max} = a\omega = 2 \times \frac{\pi}{3} = \frac{2\pi}{3}$ A1
 $\text{KE}_{\max} = \frac{1}{2} m v_{\max}^2 = \frac{1}{2} \times \frac{1}{2} \times (\frac{2\pi}{3})^2 = \frac{1}{9} \pi^2 \text{ J}$ M1 A1
- (c) $x = a \cos \omega t \therefore -0.8 = 2 \cos \omega t$ M1 A1
 $\cos \omega t = -0.4 \therefore \omega t = 1.9823 \therefore t = 1.9823 \div \frac{\pi}{3} = 1.89 \text{ s (2dp)}$ M1 A1 (10)
-
5. (a) $F = ma = Mv \frac{dv}{dx} = -\frac{4.90 \times 10^{12} \times M}{x^2}$ M1 A1
 $\int v dv = -(4.90 \times 10^{12}) \int x^{-2} dx$ M1
 $\frac{1}{2} v^2 = (4.90 \times 10^{12}) x^{-1} + c$ A1
 $x = (1.74 \times 10^6), v = u \therefore c = \frac{1}{2} u^2 - \frac{4.90 \times 10^{12}}{1.74 \times 10^6}$ M1 A1
so $v^2 = u^2 + (9.80 \times 10^{12}) x^{-1} - (5.63 \times 10^6)$ A1
- (b) we require $v > 0$ as $x \rightarrow \infty$ M1
 $x^{-1} \rightarrow 0 \therefore u^2 - (5.63 \times 10^6) > 0$ M1 A1
giving $u_{\min} = 2400 \text{ ms}^{-1} \text{ (2sf)}$ A1 (11)
-


6. (a)

portion	mass	y	my
large hemisphere	$\frac{1}{2} \times \rho \times \frac{4}{3} \pi (\frac{3}{2} r)^3 = \frac{9}{4} \rho \pi r^3$	$\frac{3}{8} \times \frac{3}{2} r = \frac{9}{16} r$	$\frac{81}{64} \rho \pi r^4$
small hemisphere	$\frac{1}{2} \times \rho \times \frac{4}{3} \pi r^3 = \frac{2}{3} \rho \pi r^3$	$\frac{3}{8} r$	$\frac{1}{4} \rho \pi r^4$
bowl	$\frac{19}{12} \rho \pi r^3$	\bar{y}	$\frac{65}{64} \rho \pi r^4$

$\rho =$ mass per unit volume y coords. taken vert. from plane face M2 A3

$\frac{19}{12} \rho \pi r^3 \times \bar{y} = \frac{65}{64} \rho \pi r^4 \therefore \bar{y} = \frac{65}{64} r \div \frac{19}{12} = \frac{195}{304} r$ M1 A1

(b)




mom. about vert. through pt of contact

$Mg \times \frac{195}{304} r \sin \alpha = \frac{1}{2} Mg \times \frac{3}{2} r \cos \alpha$ M2 A2

$\therefore \tan \alpha = \frac{3}{4} \div \frac{195}{304} = \frac{76}{65}$ M1 A1 (13)

7. (a)




resolve \uparrow : $R \cos 35 - mg = 0, R = \frac{mg}{\cos 35}$ M1

resolve \leftarrow : $R \sin 35 = \frac{mv^2}{r} = \frac{mv^2}{25}$ M1 A1

combining, $v^2 = 25g \tan 35$ M1

giving $v = 13.1 \text{ ms}^{-1}$ (3sf) A1

(b)



resolve \uparrow : $R \cos 35 + \mu R \sin 35 - mg = 0$ M1 A1

$R = \frac{mg}{\cos 35 + \mu \sin 35}$


resolve \leftarrow : $R \sin 35 - \mu R \cos 35 = \frac{mv^2}{r} = \frac{100m}{25}$ M1 A1

combining, $\frac{mg(\sin 35 - \mu \cos 35)}{\cos 35 + \mu \sin 35} = 4m$ M1 A1

giving $g(\sin 35 - \mu \cos 35) = 4(\cos 35 + \mu \sin 35)$ M1

$\therefore \mu = \frac{g \sin 35 - 4 \cos 35}{4 \sin 35 + g \cos 35} = 0.227$ (3sf) A1

(c)



resolve \uparrow : $R \cos 35 - \mu R \sin 35 - mg = 0$ M1

$R = \frac{mg}{\cos 35 - \mu \sin 35}$

resolve \leftarrow : $R \sin 35 + \mu R \cos 35 = \frac{mv^2}{r} = \frac{mv^2}{25}$ M1

combining, $\frac{mg(\sin 35 + \mu \cos 35)}{\cos 35 - \mu \sin 35} = \frac{mv^2}{25}$ M1

giving $v^2 = \frac{25g(\sin 35 + \mu \cos 35)}{\cos 35 - \mu \sin 35}$ M1

$\therefore v = 16 \text{ ms}^{-1}$ (2sf) A1 (18)

Total (75)

