

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**

**Advanced Subsidiary General Certificate of Education  
Advanced General Certificate of Education**

**MATHEMATICS**

**4731**

Mechanics 4

Wednesday

**21 JUNE 2006**

Afternoon

1 hour 30 minutes

Additional materials:

8 page answer booklet

Graph paper

List of Formulae (MF1)

**TIME** 1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by  $g \text{ m s}^{-2}$ . Unless otherwise instructed, when a numerical value is needed, use  $g = 9.8$ .
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

---

**This question paper consists of 4 printed pages.**

- 1 A straight rod  $AB$  of length  $a$  has variable density. At a distance  $x$  from  $A$  its mass per unit length is  $k(a + 2x)$ , where  $k$  is a positive constant. Find the distance from  $A$  of the centre of mass of the rod. [5]

- 2 A flywheel takes the form of a uniform disc of mass  $8 \text{ kg}$  and radius  $0.15 \text{ m}$ . It rotates freely about an axis passing through its centre and perpendicular to the disc. A couple of constant moment is applied to the flywheel. The flywheel turns through an angle of  $75$  radians while its angular speed increases from  $10 \text{ rad s}^{-1}$  to  $25 \text{ rad s}^{-1}$ .


- (i) Find the moment of the couple about the axis. [5]

When the flywheel is rotating with angular speed  $25 \text{ rad s}^{-1}$ , it locks together with a second flywheel which is mounted on the same axis and is at rest. Immediately afterwards, both flywheels rotate together with the same angular speed  $9 \text{ rad s}^{-1}$ .

- (ii) Find the moment of inertia of the second flywheel about the axis. [3]

- 3 The region bounded by the  $x$ -axis, the lines  $x = 1$  and  $x = 2$  and the curve  $y = \frac{1}{x^2}$  for  $1 \leq x \leq 2$ , is occupied by a uniform lamina of mass  $24 \text{ kg}$ . The unit of length is the metre. Find the moment of inertia of this lamina about the  $x$ -axis. [8]

4




A uniform rod  $AB$ , of mass  $m$  and length  $2a$ , is freely hinged to a fixed point at  $A$ . A particle of mass  $2m$  is attached to the rod at  $B$ . A light elastic string, with natural length  $a$  and modulus of elasticity  $5mg$ , passes through a fixed smooth ring  $R$ . One end of the string is fixed to  $A$  and the other end is fixed to the mid-point  $C$  of  $AB$ . The ring  $R$  is at the same horizontal level as  $A$ , and is at a distance  $a$  from  $A$ . The rod  $AB$  and the ring  $R$  are in a vertical plane, and  $RC$  is at an angle  $\theta$  above the horizontal, where  $0 < \theta < \frac{1}{4}\pi$ , so that the acute angle between  $AB$  and the horizontal is  $2\theta$  (see diagram).

- (i) By considering the energy of the system, find the value of  $\theta$  for which the system is in equilibrium. [7]
- (ii) Determine whether this position of equilibrium is stable or unstable. [3]

- 5 A uniform rectangular lamina  $ABCD$  has mass 20 kg and sides of lengths  $AB = 0.6$  m and  $BC = 1.8$  m. It rotates in its own vertical plane about a fixed horizontal axis which is perpendicular to the lamina and passes through the mid-point of  $AB$ .

(i) Show that the moment of inertia of the lamina about the axis is  $22.2 \text{ kg m}^2$ . [3]




The lamina is released from rest with  $BC$  horizontal and below the level of the axis. Air resistance may be neglected, but a frictional couple opposes the motion. The couple has constant moment  $44.1 \text{ N m}$  about the axis. The angle through which the lamina has turned is denoted by  $\theta$  (see diagram).

(ii) Show that the angular acceleration is zero when  $\cos \theta = 0.25$ . [3]

(iii) Hence find the maximum angular speed of the lamina. [5]

6



A ship  $P$  is moving with constant velocity  $7 \text{ m s}^{-1}$  in the direction with bearing  $110^\circ$ . A second ship  $Q$  is moving with constant speed  $10 \text{ m s}^{-1}$  in a straight line. At one instant  $Q$  is at the point  $X$ , and  $P$  is  $7400$  m from  $Q$  on a bearing of  $050^\circ$  (see diagram). In the subsequent motion, the shortest distance between  $P$  and  $Q$  is  $1790$  m.

(i) Show that one possible direction for the velocity of  $Q$  relative to  $P$  has bearing  $036^\circ$ , to the nearest degree, and find the bearing of the other possible direction of this relative velocity. [3]


Given that the velocity of  $Q$  relative to  $P$  has bearing  $036^\circ$ , find

(ii) the bearing of the direction in which  $Q$  is moving, [4]

(iii) the magnitude of the velocity of  $Q$  relative to  $P$ , [2]

(iv) the time taken for  $Q$  to travel from  $X$  to the position where the two ships are closest together, [3]

(v) the bearing of  $P$  from  $Q$  when the two ships are closest together. [1]



A uniform rod  $AB$  has mass  $m$  and length  $6a$ . It is free to rotate in a vertical plane about a smooth fixed horizontal axis passing through the point  $C$  on the rod, where  $AC = a$ . The angle between  $AB$  and the upward vertical is  $\theta$ , and the force acting on the rod at  $C$  has components  $R$  parallel to  $AB$  and  $S$  perpendicular to  $AB$  (see diagram). The rod is released from rest in the position where  $\theta = \frac{1}{3}\pi$ . Air resistance may be neglected.

- (i) Find the angular acceleration of the rod in terms of  $a$ ,  $g$  and  $\theta$ . [4]
- (ii) Show that the angular speed of the rod is  $\sqrt{\frac{2g(1 - 2\cos\theta)}{7a}}$ . [3]
- (iii) Find  $R$  and  $S$  in terms of  $m$ ,  $g$  and  $\theta$ . [6]
- (iv) When  $\cos\theta = \frac{1}{3}$ , show that the force acting on the rod at  $C$  is vertical, and find its magnitude. [4]