


Newton's Laws Questions


- 7 A builder ties two identical buckets, P and Q , to the ends of a light inextensible rope. He hangs the rope over a smooth beam so that the buckets hang in equilibrium, as shown in the diagram.



The buckets are each of mass 0.6 kg .

- (a) (i) State the magnitude of the tension in the rope. *(1 mark)*
- (ii) State the magnitude and direction of the force exerted on the beam by the rope. *(2 marks)*
- (b) The bucket Q is held at rest while a stone, of mass 0.2 kg , is placed inside it. The system is then released from rest and, in the subsequent motion, bucket Q moves vertically downwards with the stone inside.
- (i) By forming an equation of motion for each bucket, show that the magnitude of the tension in the rope during the motion is 6.72 newtons , correct to three significant figures. *(6 marks)*
- (ii) State the magnitude of the force exerted on the beam by the rope while the motion takes place. *(1 mark)*
-

- 5 A small block P is attached to another small block Q by a light inextensible string. The block P rests on a rough horizontal surface and the string hangs over a smooth peg so that Q hangs freely, as shown in the diagram.



The block P has mass 0.4 kg and the coefficient of friction between P and the surface is 0.5 .

The block Q has mass 0.3 kg .

The system is released from rest and Q moves vertically downwards.

- (a) (i) Draw a diagram to show the forces acting on P . (1 mark)
- (ii) Show that the frictional force between P and the surface has magnitude 1.96 newtons . (2 marks)
- (b) By forming an equation of motion for each block, show that the magnitude of the acceleration of each block is 1.4 ms^{-2} . (5 marks)
- (c) Find the speed of the blocks after 3 seconds of motion. (2 marks)
- (d) After 3 seconds of motion, the string breaks. The blocks continue to move. Comment on how the speed of each block will change in the subsequent motion. For each block, give a reason for your answer. (4 marks)
-

- 2 A lift rises vertically from rest with a constant acceleration.


After 4 seconds, it is moving upwards with a velocity of 2 m s^{-1} .

It then moves with a constant velocity for 5 seconds.

The lift then slows down uniformly, coming to rest after it has been moving for a total of 12 seconds.


- (a) Sketch a velocity–time graph for the motion of the lift. (4 marks)
- (b) Calculate the total distance travelled by the lift. (2 marks)
- (c) The lift is raised by a single vertical cable. The mass of the lift is 300 kg . Find the maximum tension in the cable during this motion. (4 marks)
-

- 4 The diagram shows a block, of mass 13 kg, on a rough horizontal surface. It is attached by a string that passes over a smooth peg to a sphere of mass 7 kg, as shown in the diagram.



The system is released from rest, and after 4 seconds the block and the sphere both have speed 6 m s^{-1} , and the block has **not** reached the peg.

- (a) State **two** assumptions that you should make about the string in order to model the motion of the sphere and the block. *(2 marks)*
- (b) Show that the acceleration of the sphere is 1.5 m s^{-2} . *(2 marks)*
- (c) Find the tension in the string. *(3 marks)*
- (d) Find the coefficient of friction between the block and the surface. *(6 marks)*
-
- 4 A car, of mass 1200 kg, is connected by a tow rope to a truck, of mass 2800 kg. The truck tows the car in a straight line along a horizontal road. Assume that the tow rope is horizontal. A horizontal driving force of magnitude 3000 N acts on the truck. A horizontal resistance force of magnitude 800 N acts on the car. The car and truck accelerate at 0.4 m s^{-2} .




- (a) Find the tension in the tow rope. *(3 marks)*
- (b) Show that the magnitude of the horizontal resistance force acting on the truck is 600 N. *(4 marks)*
- (c) In fact, the tow rope is **not** horizontal. Assume that the resistance forces and the driving force are unchanged.

Is the tension in the tow rope greater or less than in part (a)?

Explain why. *(2 marks)*

Newtons Laws Answers

7(a)(i)	$T = 0.6 \times 9.8 = 5.88N$ Or $0.6g$	B1	1	
(ii)	Force = $2T = \downarrow 11.76N$ Or $11.8 N$ Or $1.2g$	B1		Magnitude
		B1	2	Direction
(b)(i)	$Q: 0.8g - T = 0.8a$	M1		Either equation
	$T - 0.6g = 0.6a$	A1		
	$0.2g = 1.4a$	m1		Alternative for m1 A1 if solving for T
	$a = 1.4$	A1		m1 method for solving, A1 accurate attempt
	$T = 6.72N$	A1	6	cao SC whole string to find $a: 0.2g = 1.4a$ M1 $a = 1.4$ A1
(ii)	Force = $2T = 13.44N$	B1	1	cao to find $T: M1 A1$
Total			10	

5(a)(i)		B1	1	Accept mg , $0.4g$ or 3.92 for weight Arrows and labels needed
(ii)	$F = 0.5 \times (0.4 \times 9.8)$ $F = 1.96N$	M1		Need to see 0.4×9.8 or 3.92 used
		A1	2	
(b)	$T - 1.96 = 0.4a$ $0.3g - T = 0.3a$ $a = 1.4ms^{-2}$	M1A1		Consistent reversal of signs in both equations 4 marks; reversal of signs in one equation, M1 A1 M1 A0 Sign change needs justification (whole string: equation, $0.3g - 1.96 = 0.7a$ M1A1 $a = 1.4$ A1) max 3/5
		M1A1		
		A1	5	
(c)	$v = 1.4 \times 3$ $v = 4.2ms^{-2}$	M1		Full method
		A1	2	CAO
(d)	P : Friction will cause speed to decrease	M1 A1		Accept decelerate or comes to rest
	Q : Gravity will cause speed to increase	M1 A1	4	Accept accelerate
Total			14	

2(a)		B1 B1 B1 B1	4	Starts and finishes at rest Correct shape Correct values on t -axis Correct values on v -axis Condone omission of the origin
(b)	$s = \frac{1}{2}(5+12) \times 2$ <p>or $s = \frac{1}{2} \times 2 \times 4 + 5 \times 2 + \frac{1}{2} \times 2 \times 3 = 17$ $= 17$</p>	M1 A1	2	Use of the area under the graph (or equivalent) to find s Correct distance SC When 21 used instead of 12 allow full marks for $s = 26$
(c)	$\max a = \frac{2}{4} = 0.5$ $300 \times 0.5 = T - 300 \times 9.8$ $T = 2940 + 150 = 3090$	B1 M1 A1 A1	4	Maximum acceleration Three term equation of motion using their a Correct equation using $a = 0.5$ Correct tension
			10	

4(a)	The string is light and inextensible or inelastic or taut	B1 B1	2	First assumption Second assumption
(b)	$6 = 0 + 4a$ $a = \frac{6}{4} = 1.5$	M1 A1	2	Finding a using a CA equation Correct a from correct working
(c)	$7 \times 9.8 - T = 7 \times 1.5$ $T = 68.6 - 10.5 = 58.1$	M1A1 A1	3	Three term equation of motion with F for the 7 kg particle. Correct equation Correct tension
(d)	$58.1 - F = 13 \times 1.5$ $F = 58.1 - 19.5 = 38.6$ $R = 13.98 = 127.4$ $38.6 = \mu \times 127.4$ $\mu = \frac{38.6}{127.4} = 0.303$	M1A1 A1 B1 dM1 A1	6	Three term equation of motion with F for the 13 kg particle. Correct equation Correct F Correct R Use of $F = \mu R$ Correct coefficient of friction
			13	

4(a)	$T - 800 = 1200 \times 0.4$ $T = 800 + 480$ $= 1280 \text{ N}$	M1 A1 A1		Three term equation of motion for the car Correct equation
			3	Correct tension Treat calculation of two tensions as two methods unless one selected Treat sum or difference of two tensions as an incorrect method
(b)	$3000 - 800 - F = 4000 \times 0.4$ $F = 3000 - 800 - 1600$ $F = 600 \text{ N}$ OR $3000 - 1280 - F = 2800 \times 0.4$ $F = 3000 - 1280 - 1120$ $F = 600 \text{ N}$	M1 A1 A1 A1		Four term equation of motion (truck or both) Correct terms Correct signs AG Correct resistance force from correct working
(c)	Increase, because a greater tension would be needed so that the horizontal component would be the same as the tension above.	B1 B1	2	Greater Reason Second B1 dependent on the first B1 mark
Total		9		