

Figure 2

A beam *AB* has mass *m* and length 2*a*.

The beam rests in equilibrium with *A* on rough horizontal ground and with *B* against a smooth vertical wall.

The beam is inclined to the horizontal at an angle  $\theta$ , as shown in Figure 2.

The coefficient of friction between the beam and the ground is  $\mu$ 

The beam is modelled as a uniform rod resting in a vertical plane that is perpendicular to the wall.

Using the model,

(a) show that  $\mu \ge \frac{1}{2}\cot\theta$ 

A horizontal force of magnitude kmg, where k is a constant, is now applied to the beam at A.

This force acts in a direction that is perpendicular to the wall and towards the wall.

Given that  $\tan \theta = \frac{5}{4}$ ,  $\mu = \frac{1}{2}$  and the beam is now in limiting equilibrium,

(b) use the model to find the value of *k*.

(5)

(5)

2.

(3)




Figure 2

A uniform rod AB has mass M and length 2a

A particle of mass 2M is attached to the rod at the point C, where AC = 1.5a

The rod rests with its end A on rough horizontal ground.

The rod is held in equilibrium at an angle  $\theta$  to the ground by a light string that is attached to the end *B* of the rod.

The string is perpendicular to the rod, as shown in Figure 2.

(a) Explain why the frictional force acting on the rod at *A* acts horizontally to the right on the diagram.

(1)

The tension in the string is T

(b) Show that  $T = 2Mg\cos\theta$ 

Given that  $\cos\theta = \frac{3}{5}$ 

(c) show that the magnitude of the vertical force exerted by the ground on the rod at A

is 
$$\frac{57Mg}{25}$$
 (3)

The coefficient of friction between the rod and the ground is  $\mu$ 

Given that the rod is in limiting equilibrium,

(d) show that 
$$\mu = \frac{8}{19}$$
 (4)

3.




Figure 3

A rod AB has mass M and length 2a.

The rod has its end A on rough horizontal ground and its end B against a smooth vertical wall.

The rod makes an angle  $\theta$  with the ground, as shown in Figure 3.

The rod is at rest in limiting equilibrium.

(a) State the direction (left or right on Figure 3 above) of the frictional force acting on the rod at *A*. Give a reason for your answer.

(1)

The magnitude of the normal reaction of the wall on the rod at B is S.

In an initial model, the rod is modelled as being uniform.

## Use this initial model to answer parts (b), (c) and (d).

(b) By taking moments about A, show that

$$S = \frac{1}{2} Mg \cot\theta$$
(3)

The coefficient of friction between the rod and the ground is  $\mu$ 

Given that  $\tan \theta = \frac{3}{4}$ 

- (c) find the value of  $\mu$
- (d) find, in terms of *M* and *g*, the magnitude of the resultant force acting on the rod at *A*.

(3)

(5)

In a new model, the rod is modelled as being **non-uniform**, with its centre of mass closer to *B* than it is to *A*.

A new value for S is calculated using this new model, with  $\tan \theta = \frac{3}{4}$ 

(e) State whether this new value for S is larger, smaller or equal to the value that S would take using the initial model. Give a reason for your answer.

(1)