| Please write clearly, in block capitals. |                  |
|--|------------------|
| Centre number                            | Candidate number |
| Surname                                  |                  |
| Forename(s)                              |                  |
| Candidate signature                      |                  |

# AS FURTHER MATHEMATICS

Paper 2 – Statistics



Time allowed: 1 hour 30 minutes

## **Materials**

For this paper you must have:

- You must ensure you have the other optional question paper/answer booklet for which you are entered (**either** Mechanics **or** Discrete). You will have 1 hour 30 minutes to complete both papers.
- The AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

## Instructions

- Use black ink or black ball-point pen. Pencil should be used for drawing.
- Answer all questions.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

#### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 40.

# Advice

Unless stated otherwise, you may quote formulae, without proof, from the booklet. You do not necessarily need to use all the space provided.

2

Answer **all** questions in the spaces provided.

 $f(t) = \begin{cases} \frac{t}{8} & 0 \le t \le k \\ 0 & \text{otherwise} \end{cases}$ Find the value of k [1 mark]  $\frac{1}{16} & \frac{1}{4} & 4 & 16$ 

The random variable T has probability density defined by

2 The discrete random variable *X* has probability distribution defined by

 $\mathsf{P}(X=x) = \begin{cases} 0.1 & x = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9\\ 0 & \text{otherwise} \end{cases}$ 

Find the value of  $P(4 \le X \le 7)$ 

Circle your answer.

[1 mark]

| 0.2 | 0.3 | 0.4 | 0.5 |
|-----|-----|-----|-----|
|-----|-----|-----|-----|

1

| The discrete random variable $R$ has the following probability distribution. |  |
|--|--|
|--|--|

| r                                | -2  | 0 | а | 4   |
|----------------------------------|-----|---|---|-----|
| <b>P</b> ( <i>R</i> = <i>r</i> ) | 0.3 | b | С | 0.1 |

It is known that E(R) = 0.2 and Var(R) = 3.56

Find the values of a, b and c.

3

[4 marks]

Turn over for the next question

4 The number of printers, *V*, bought during one day from the *Verigood* store can be modelled by a Poisson distribution with mean 4.5

The number of printers, W, bought during one day from the *Winnerprint* store can be modelled by a Poisson distribution with mean 5.5

**4 (a)** Find the probability that the total number of printers bought during one day from *Verigood* and *Winnerprint* stores is greater than 10.

[2 marks]

**4 (b)** State the circumstance under which the distributional model you used in part **(a)** would not be valid.

[1 mark]

**5** Participants in a school jumping competition gain a total score for each jump based on the length, *L* metres, jumped beyond a fixed point and a mark, *S*, for style.

L may be regarded as a continuous random variable with probability density function

$$f(l) = \begin{cases} wl & 0 \le l \le 15 \\ 0 & \text{otherwise} \end{cases}$$

where w is a constant.

S may be regarded as a discrete random variable with probability function

$$P(S = s) = \begin{cases} \frac{1}{15}s & s = 1, 2, 3, 4, 5\\ 0 & \text{otherwise} \end{cases}$$

Assume that *L* and *S* are independent.

The total score for a participant in this competition, *T*, is given by  $T = L^2 + \frac{1}{2}S$ 

Show that the expected total score for a participant is  $114\frac{1}{3}$ 

[5 marks]


| <br> |
|------|
|      |
| <br> |
| <br> |
| <br> |
|      |
|      |
|      |
| <br> |
|      |

6 The continuous random variable *T* has probability density function defined by

$$f(t) = \begin{cases} \frac{1}{3} & 0 \le t \le \frac{3}{2} \\ \frac{9-2t}{18} & \frac{3}{2} \le t \le \frac{9}{2} \\ 0 & \text{otherwise} \end{cases}$$

6 (a) (i) Sketch this probability density function below.

[2 marks]



6 (a) (ii) State the median of *T*.

[1 mark]

| • |
|---|
|   |

| 6 | (b) (i)  | Find E(T) [2 mark  | s] |
|---|----------|--|----|
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |
| 6 | (b) (ii) | Given that $E(T^2) = \frac{15}{4}$ , find Var $(4T-5)$ [3 mark | s] |
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |
|   |          |  |    |

Turn over for the next question

- 9
- 7 A dairy industry researcher, Robyn, decided to investigate the milk yield, classified as low, medium or high, obtained from four different breeds of cow, A, B, C and D.

The milk yield of a sample of 105 cows was monitored and the results are summarised in contingency **Table 1**.

| Table 1 |       | Yield |        |      |       |
|---------|-------|-------|--------|------|-------|
|         |       | Low   | Medium | High | Total |
| Breed   | Α     | 4     | 5      | 12   | 21    |
|         | В     | 10    | 6      | 4    | 20    |
|         | С     | 8     | 17     | 7    | 32    |
|         | D     | 5     | 20     | 7    | 32    |
|         | Total | 27    | 48     | 30   | 105   |

The sample of cows may be regarded as random.

Robyn decides to carry out a  $\chi^2$ -test for association between milk yield and breed using the information given in **Table 1**.

7 (a) Contingency **Table 2** gives some of the expected frequencies for this test.

Complete **Table 2** with the missing expected values.

[2 marks]

| Table 2 |   | Yield |        |      |  |
|---------|---|-------|--------|------|--|
|         |   | Low   | Medium | High |  |
|         | Α |       |        | 6    |  |
| Breed   | В | 5.14  | 9.14   | 5.71 |  |
|         | С |       |        |      |  |
|         | D | 8.23  | 14.63  | 9.14 |  |

| 7 (b) (i)  | For Robyn's test, the test statistic $\sum \frac{(O-E)^2}{E} = 19.4$ correct to three significant figures.   |
|------------|--|
|            | Use this information to carry out Robyn's test, using the 1% level of significance. [5 marks]  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
|            |  |
| 7 (b) (ii) | By considering the observed frequencies given in <b>Table 1</b> with the expected frequencies in <b>Table 2</b> , interpret, in context, the association, if any, between milk yield and breed.<br>[2 marks] |
|            |  |
|            |  |
|            |  |

8 In a small town, the number of properties sold during a week in spring by a local estate agent, Keith, can be regarded as occurring independently and with constant mean  $\mu$ . Data from several years have shown the value of  $\mu$  to be 3.5.

A new housing development was built on the outskirts of the town and the properties on this development were offered for sale by the builder of the development, not by the local estate agents.

During the first **four** weeks in spring, when properties on the new development were offered for sale by the builder, Keith sold a total of 8 properties.

Keith claims that the sale of new properties by the builder reduced his mean number of properties sold during a week in spring.

8 (a) Investigate Keith's claim, using the 5% level of significance.

[6 marks]

8 (b) For your test carried out in part (a) state, in context, the meaning of a Type II error. [1 mark]
[1 mark]
[2 mark]
[2 marks]

## END OF QUESTIONS

Copyright © 2017 AQA and its licensors. All rights reserved.