



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/13

Paper 1 Further Pure Mathematics 1

May/June 2020

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

1 The cubic equation $7x^3 + 3x^2 + 5x + 1 = 0$ has roots α, β, γ .

(a) Find a cubic equation whose roots are $\alpha^{-1}, \beta^{-1}, \gamma^{-1}$. [3]

.....

.....

.....

.....

.....

.....

.....

(b) Find the value of $\alpha^{-2} + \beta^{-2} + \gamma^{-2}$. [2]

.....

.....

.....

.....

.....

.....

.....

(c) Find the value of $\alpha^{-3} + \beta^{-3} + \gamma^{-3}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3

2 The sequence u_1, u_2, u_3, \dots is such that $u_1 = 1$ and $u_{n+1} = 2u_n + 1$ for $n \geq 1$.

(a) Prove by induction that $u_n = 2^n - 1$ for all positive integers n . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Deduce that u_{2^n} is divisible by u_n for $n \geq 1$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 Let $S_n = 2^2 + 6^2 + 10^2 + \dots + (4n-2)^2$.

(a) Use standard results from the List of Formulae (MF19) to show that $S_n = \frac{4}{3}n(4n^2 - 1)$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Express $\frac{n}{S_n}$ in partial fractions and find $\sum_{n=1}^N \frac{n}{S_n}$ in terms of N . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Deduce the value of $\sum_{n=1}^{\infty} \frac{n}{S_n}$. [1]

.....

.....

.....

.....

.....

.....

.....

4 The matrix \mathbf{A} is given by

$$\mathbf{A} = \begin{pmatrix} k & 0 & 2 \\ 0 & -1 & -1 \\ 1 & 1 & -k \end{pmatrix},$$

where k is a real constant.

(a) Show that \mathbf{A} is non-singular.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The matrices \mathbf{B} and \mathbf{C} are given by

$$\mathbf{B} = \begin{pmatrix} 0 & -3 \\ -1 & 3 \\ 0 & 0 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} -3 & -1 & 1 \\ 1 & 1 & 2 \end{pmatrix}.$$

It is given that $\mathbf{CAB} = \begin{pmatrix} -2 & -\frac{3}{2} \\ -1 & -\frac{3}{2} \end{pmatrix}$.

(b) Find the value of k .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 The curve C has polar equation $r = a \tan \theta$, where a is a positive constant and $0 \leq \theta \leq \frac{1}{4}\pi$.

(a) Sketch C and state the greatest distance of a point on C from the pole. [2]

(b) Find the exact value of the area of the region bounded by C and the half-line $\theta = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show that C has Cartesian equation $y = \frac{x^2}{\sqrt{a^2 - x^2}}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Using your answer to part (b), deduce the exact value of $\int_0^{\frac{1}{2}a\sqrt{2}} \frac{x^2}{\sqrt{a^2 - x^2}} dx$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The curve C has equation $y = \frac{10+x-2x^2}{2x-3}$.

(a) Find the equations of the asymptotes of C . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that C has no turning points. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Sketch C , stating the coordinates of the intersections with the axes.

[3]

- (d) Sketch the curve with equation $y = \left| \frac{10+x-2x^2}{2x-3} \right|$ and find in exact form the set of values of x for which $\left| \frac{10+x-2x^2}{2x-3} \right| < 4$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 7 The lines l_1 and l_2 have equations $\mathbf{r} = -5\mathbf{j} + \lambda(5\mathbf{i} + 2\mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k} + \mu(\mathbf{j} + \mathbf{k})$ respectively. The plane Π contains l_1 and is parallel to l_2 .

(a) Find the equation of Π , giving your answer in the form $ax + by + cz = d$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the distance between l_2 and Π . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The point P on l_1 and the point Q on l_2 are such that PQ is perpendicular to both l_1 and l_2 .

- (c) Show that P has position vector $\frac{55}{27}\mathbf{i} - 5\mathbf{j} + \frac{22}{27}\mathbf{k}$ and state a vector equation for PQ . [8]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

