



Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/11

Paper 1 Further Pure Mathematics 1

May/June 2023

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

BLANK PAGE

2 The cubic equation $x^3 + 4x^2 + 6x + 1 = 0$ has roots α, β, γ .

(a) Find the value of $\alpha^2 + \beta^2 + \gamma^2$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use standard results from the list of formulae (MF19) to show that

$$\sum_{r=1}^n ((\alpha+r)^2 + (\beta+r)^2 + (\gamma+r)^2) = n(n^2 + an + b),$$

where a and b are constants to be determined. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that **M** represents the sequence of two transformations in the x - y plane given by an enlargement, centre the origin, scale factor 5 followed by a shear, x -axis fixed, with $(0, 1)$ mapped to $(5, 1)$.

(c) Find **M**. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) The triangle DEF in the x - y plane is transformed by **M** onto triangle PQR .
Given that the area of triangle DEF is 12 cm^2 , find the area of triangle PQR . [2]

.....

.....

.....

.....

.....

10

5 The curve C has polar equation $r^2 = \frac{1}{\theta^2 + 1}$, for $0 \leq \theta \leq \pi$.

(a) Sketch C and state the polar coordinates of the point of C furthest from the pole. [3]

.....

(b) Find the area of the region enclosed by C , the initial line, and the half-line $\theta = \pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Show that, at the point of C furthest from the initial line,

$$\left(\theta + \frac{1}{\theta}\right)\cot\theta - 1 = 0$$

and verify that this equation has a root between 1.1 and 1.2.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The curve C has equation $y = \frac{x^2 + 2x - 15}{x - 2}$.

(a) Find the equations of the asymptotes of C . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that C has no stationary points. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Sketch C , stating the coordinates of the intersections with the axes. [3]

-
- (d) Sketch the curve with equation $y = \left| \frac{x^2 - 2x - 15}{x - 2} \right|$. [2]

BLANK PAGE

7 The plane Π_1 has equation $r = -4\mathbf{j} - 3\mathbf{k} + \lambda(\mathbf{i} - \mathbf{j} + \mathbf{k}) + \mu(\mathbf{i} + \mathbf{j} - \mathbf{k})$.

(a) Obtain an equation of Π_1 in the form $px + qy + rz = d$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The plane Π_2 has equation $\mathbf{r} \cdot (-5\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}) = 4$.

Find a vector equation of the line of intersection of Π_1 and Π_2 . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line l passes through the point A with position vector $a\mathbf{i} + a\mathbf{j} + (a - 7)\mathbf{k}$ and is parallel to $(1 - b)\mathbf{i} + b\mathbf{j} + b\mathbf{k}$, where a and b are positive constants.

- (c) Given that the perpendicular distance from A to Π_1 is $\sqrt{2}$, find the value of a . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Given that the obtuse angle between l and Π_1 is $\frac{3}{4}\pi$, find the exact value of b . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.