



## Cambridge International AS & A Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



### FURTHER MATHEMATICS

9231/01

Paper 1 Further Pure Mathematics 1

For examination from 2020

SPECIMEN PAPER

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

### INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

1 (a) Given that  $f(r) = \frac{1}{(r+1)(r+2)}$ , show that

$$f(r-1) - f(r) = \frac{2}{r(r+1)(r+2)}. \quad [2]$$

• • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •

(b) Hence find  $\sum_{r=1}^n \frac{1}{r(r+1)(r+2)}$ . [3]

• • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •  
 • • •

(c) Deduce that  $\sum_{r=1}^{\infty} \frac{1}{r(r+1)(r+2)}$ . [1]

• • •  
 • • •

2 It is given that  $\phi(n) = 5^n(4n + 1) - 1$  for  $n \in \mathbb{Z}_{>0}$ .

Prove, by mathematical induction, that  $\phi(n)$  is divisible by 5 for every positive integer  $n$ . [7]

•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•

3 The curve  $C$  has polar equation  $r = 2 + 2 \cos \theta$ , for  $0 \leq \theta \leq \pi$ .

(a) Sketch  $C$ .

[3]

(b) Find the area of the region enclosed by  $C$  and the initial line.

[4]

.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •

(c) Show that the Cartesian equation of the circle expressed as  $4x^2 + y^2 = (x^2 + y^2 - 2x)^2$ . [3]

• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •  
• • •

4 The cubic equation

$$z^3 - z^2 - z - 5 = 0$$

has roots  $\alpha, \beta$  and  $\gamma$ .

(a) Show that the value of  $\alpha^3 + \beta^3 + \gamma^3$  is 9

[4]

.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.
.	.	.

(b) Find the value of  $\alpha^4 + \beta^4 + \gamma^4$ .

[2]

•		•	•
•		•	•
•		•	•
•		•	•
•		•	•
•		•	•
•		•	•
•		•	•
•		•	•

(c) Find the cubic equation in  $x$  whose roots are  $\alpha + 1$ ,  $\beta + 1$  and  $\gamma + 1$  if the answer is in the form

$$px^3 + qx^2 + rx + s = 0$$

where  $p, q, r$  and  $s$  are constants to be determined

[3]

•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•
•	•	•	•

5 The matrix  $A$  is given by

$$A = \begin{pmatrix} 5 & k \\ -3 & -4 \end{pmatrix}.$$

(a) Find the value of  $k$  for which  $A$  is singular.

[2]

•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•

It is given that  $k = 6$  so that  $A = \begin{pmatrix} 5 & 6 \\ -3 & -4 \end{pmatrix}$ .

(b) Find the equations of the invariant lines, the origin of the transformation in the  $x$ - $y$  plane represented by  $A$ .

[6]

•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•
•	•	•





(c) The triangle  $DEF$  in the  $x$ - $y$  plane is transformed into triangle  $PQR$ .

(i) Given that the area of triangle  $DEF$  is  $0 \text{ cm}^2$ , find the area of triangle  $PQR$ . [2]



(ii) Find the matrix which transforms triangle  $PQR$  into triangle  $DEF$ . [2]



6 The position vectors  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$ ,  $\mathbf{d}$  are

$$2\mathbf{i} + 4\mathbf{j} - 3\mathbf{k}, \quad -2\mathbf{i} + 5\mathbf{j} - 4\mathbf{k}, \quad \mathbf{i} + 4\mathbf{j} + \mathbf{k}, \quad \mathbf{i} + 5\mathbf{j} + m\mathbf{k},$$

respectively, where  $m$  is an integer. It is given that the shortest distance between the lines  $\mathbf{a}$  and  $\mathbf{b}$  and the line  $\mathbf{c}$  and  $\mathbf{d}$  is 3.

(a) Show that the hypothesis for  $m$  is 2. [7]

.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.
.	.	.	.	.	.	.	.	.	.

- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 

(b) Find the standard deviation of  $D$  from the line through  $A$  and  $C$ . [3]

- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
-

(c) Show that the acute angle between the planes  $ACD$  and  $BCD$  is  $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ . [4]

- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
-

7 The curve C has equation  $y = \frac{2x^2 - 3x - 2}{x^2 - 2x + 1}$ .

(a) State the equation of the asymptotes of C. [2]

.	.	.
.	.	.
.	.	.
.	.	.
.	.	.

(b) Show that  $y \leq \frac{25}{12}$  at all points on C. [4]

.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

(c) Find the coordinates of any stationary points of  $C$ . [3]

- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 
- 

(d) Sketch  $C$ , stating the coordinates of any intersections of  $C$  with the coordinate axes and the asymptotes. [4]

(e) Sketch the curve with equation  $y = \left| \frac{2x^2 - 3x - 2}{x^2 - 2x + 1} \right|$  and find the set of values of  $x$  for which

$$\left| \frac{2x^2 - 3x - 2}{x^2 - 2x + 1} \right| < 2 \quad [4]$$

.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.
.	.	.	.

Additional page

If possible following instructions to complete the answer(s) to any question(s), then question number(s) must be clearly written

- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •
- • •

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.