



Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/22

Paper 2 Further Pure Mathematics 2

May/June 2021

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS


- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

3



The diagram shows the curve with equation $y = x^3$ for $0 \leq x \leq 1$, together with a set of n rectangles of width $\frac{1}{n}$.

(a) By considering the sum of the areas of these rectangles, show that $\int_0^1 x^3 dx < U_n$, where

$$U_n = \left(\frac{n+1}{2n}\right)^2. \quad [4]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Use a similar method to find, in terms of n , a lower bound L_n for $\int_0^1 x^3 dx$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find the least value of n such that $U_n - L_n < 10^{-3}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 (a) State the sum of the series $z + z^2 + z^3 + \dots + z^n$, for $z \neq 1$. [1]

.....

(b) Given that z is an n th root of unity and $z \neq 1$, deduce that $1 + z + z^2 + \dots + z^{n-1} = 0$. [2]

.....

(c) Given instead that $z = \frac{1}{3}(\cos \theta + i \sin \theta)$, use de Moivre's theorem to show that

$$\sum_{m=1}^{\infty} 3^{-m} \cos m\theta = \frac{3 \cos \theta - 1}{10 - 6 \cos \theta}. \quad [7]$$

.....

A series of horizontal dotted lines for writing, spanning the width of the page.

7 (a) It is given that $y = \operatorname{sech}^{-1}\left(x + \frac{1}{2}\right)$.

Express $\cosh y$ in terms of x and hence show that $\sinh y \frac{dy}{dx} = -\frac{1}{\left(x + \frac{1}{2}\right)^2}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the first three terms in the Maclaurin’s series for $\operatorname{sech}^{-1}\left(x + \frac{1}{2}\right)$ in the form

$$\ln a + bx + cx^2,$$

where a, b and c are constants to be determined. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of 25 horizontal dotted lines spanning the width of the page, intended for writing.

