

Cambridge International AS & A Level

FURTHER MATHEMATICS**9231/42**

Paper 4 Further Probability & Statistics

May/June 2024

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **15** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED**Mathematics Specific Marking Principles**

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	$s^2 = \frac{941.5}{13}$	B1	72.423.
	CI: $42.8 \pm 2.160\sqrt{\frac{s^2}{14}}$	M1 A1	Correct formula with a t value. 2.160 seen in a CI formula for A1.
	[37.9, 47.7]	A1	(37.9, 47.7) A1 $37.9 \leq 'x' \leq 47.7$ A1 (ignore symbol used inside the inequality) Condone [47.7, 37.9] A1 Final answer 42.8 ± 4.91 A0
		4	

PUBLISHED

Question	Answer	Marks	Guidance
2	H_0 : practical results and written results are equal H_1 : practical results are greater than written results	B1	Or $H_0: m_p - m_w = 0$, $H_1: m_p - m_w > 0$, oe (m is population median).
	There are 10 + signs; $B(15, 0.5)$ soi	M1	
	$P(X \geq 10) = 1 - P(X \leq 9) = 1 - 0.849 = 0.151$	A1	$P(X \leq 5) = 0.151$ (implies M1)
	Compare with 0.1: '0.151' > 0.1 accept H_0	M1	'0.151' must come from a valid binomial calculation Correct ft conclusion for their 0.151 and 0.1 Condone Reject H_1 'Accept H_0 ' can be implied by a conclusion that is consistent with their 0.151 and 0.1
	Insufficient evidence to suggest that practical results are greater than written results	A1	Correct work only, ignoring their hypotheses, conclusion in context with level of uncertainty in language. Not 'prove'. Condone 'no sufficient' 'not enough'. Do not accept statements such as 'there is sufficient evidence to suggest.....'.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
3(a)	Signed differences 0.4 -1.1 0.8 -0.5 1.2 0.9 1.9 -0.3 -2.2 1.6 0.6 1.0	M1	Allow one error.
	Ranks: 2 -8 5 -3 9 6 11 -1 -12 10 4 7	M1	Attempt at ranking.
	($P_+ = 54$) $P_- = 24$	A1	All ranks must be correct.
	H_0 : population median = 22.0 H_1 : population median \neq 22.0	B1	Accept m used as ‘population median’
	Critical value from table = 17	B1	
	‘24’ > ‘17’ Accept H_0	M1	‘24’ must come from a ranking of signed differences, ‘17’ must be critical value from tables (7, 9, 10, 13, 17 or 21). Correct fit conclusion for their 24 and their 17. Condone Reject H_1 . ‘Accept H_0 ’ can be implied by a conclusion that is consistent with their 24 and their 17.
	Insufficient evidence to support (population) median not being 22.0 Insufficient evidence against the manager’s claim	A1	Correct work only , ignoring their hypotheses, conclusion in context with level of uncertainty in language. Not ‘prove’. Condone ‘no sufficient’ ‘not enough’. Do not accept statements such as ‘there is sufficient evidence to suggest.....’.
		7	

PUBLISHED

Question	Answer	Marks	Guidance
3(b)	Distribution is symmetric about the population median Underlying/population distribution is symmetric about the median	B1	Need to see population or underlying distribution mentioned. Underlying distribution is symmetric B0. Population distribution is symmetric B0. ‘data’ implies B0 but condone ‘population data’ ‘mean’ implies B0.
		1	

Question	Answer	Marks	Guidance
4(a)	$G_Y(t) = \frac{t^2}{(4-3t)^4}, \quad G'_Y(t) = \frac{(4-3t)^4(2t) - 12t^2(4-3t)^3}{(4-3t)^8} \text{ or}$ $\frac{2t}{(4-3t)^4} + \frac{12t^2}{(4-3t)^5}$	M1	‘Two’ terms obtained, correct denominator(s).
	$E(Y) = G'_Y(1) = 14$	M1 A1	$t = 1$ in their expression. CWO
		3	
4(b)	$G_X(t) = \frac{t}{(4-3t)^2}$	B1	
		1	

PUBLISHED

Question	Answer	Marks	Guidance
4(c)	$G_X(t) = t(4-3t)^{-2} = \frac{t}{16} \left(1 - \frac{3}{4}t\right)^{-2} = \frac{t}{16} \left(1 + \frac{3}{2}t - 3 \times \frac{9}{16}t^2 + 4 \times \frac{27}{64}t^3 + \dots\right)$	M1	Expansion as far as t^3 OR term in t^4 calculated.
	Use $P(X=4)$ = their coefficient of t^4	M1	Find the numerical value of their coefficient of t^4 .
	$\frac{27}{256}$	A1	Accept 0.105.
		3	

PUBLISHED

Question	Answer				Marks	Guidance												
5(a)	<table><tr><td>Company</td><td><i>Poor</i></td><td><i>Satisfactory</i></td><td><i>Good</i></td></tr><tr><td><i>P</i></td><td>(18) 25</td><td>(43) 40.625</td><td>(64) 59.375</td></tr><tr><td><i>Q</i></td><td>(22) 15</td><td>(22) 24.375</td><td>(31) 35.625</td></tr></table>				Company	<i>Poor</i>	<i>Satisfactory</i>	<i>Good</i>	<i>P</i>	(18) 25	(43) 40.625	(64) 59.375	<i>Q</i>	(22) 15	(22) 24.375	(31) 35.625	M1 A1	Calculate E values to at least 2dp, allow one error. These must be seen. For A1, working to at least 2dp.
	Company	<i>Poor</i>	<i>Satisfactory</i>	<i>Good</i>														
	<i>P</i>	(18) 25	(43) 40.625	(64) 59.375														
	<i>Q</i>	(22) 15	(22) 24.375	(31) 35.625														
	Chi-squared contributions: <table><tr><td>Company</td><td><i>Poor</i></td><td><i>Satisfactory</i></td><td><i>Good</i></td></tr><tr><td><i>P</i></td><td>1.96</td><td>0.1388</td><td>0.3603</td></tr><tr><td><i>Q</i></td><td>3.2667</td><td>0.2314</td><td>0.6004</td></tr></table>				Company	<i>Poor</i>	<i>Satisfactory</i>	<i>Good</i>	<i>P</i>	1.96	0.1388	0.3603	<i>Q</i>	3.2667	0.2314	0.6004		
Company	<i>Poor</i>	<i>Satisfactory</i>	<i>Good</i>															
<i>P</i>	1.96	0.1388	0.3603															
<i>Q</i>	3.2667	0.2314	0.6004															
Test statistic = 6.56				A1	AWRT 6.56.													
H ₀ : Quality (of items) is independent of company H ₁ : Quality (of items) is not independent of company				B1	Condone ‘no association/relationship between....’ ‘association/relationship between....’.													
6.56 > 5.991 Reject H ₀				M1	‘6.56’ from their attempt at chi-squared, 5.991 must be correct. Consistent signs in comparison. FT conclusion from their 6.56. Condone ‘Accept H ₁ ’. ‘Reject H ₀ ’ can be implied by a conclusion that is consistent with their 6.56.													

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	There is sufficient evidence to suggest that the quality of brushes is not independent of company.	A1	Correct work only, conclusion in context with level of uncertainty in language. Not ‘prove’. Accept ‘enough evidence’. Condone ‘some evidence’. Do not accept ‘there is insufficient...’.
		7	
5(b)	<i>P</i> produces better quality brushes (than <i>Q</i>) or <i>Q</i> produces worse quality brushes (than <i>P</i>) <i>P</i> has fewer poor quality brushes than expected AND <i>Q</i> has more poor quality brushes than expected	B1	Condone <i>P</i> is better (than <i>Q</i>).
		1	

PUBLISHED

Question	Answer	Marks	Guidance
6(a)	Signed differences: 4.1 1.1 –0.4 0.2 3.9 2.6 –0.6 3.5 0.1 0	M1	Allow one error, must be signed differences.
	$\Sigma d = 14.5, \quad \Sigma d^2 = 52.81 \quad s^2 = \frac{1}{9} \left(52.81 - \frac{14.5^2}{10} \right) = 3.532$	M1	
	$H_0: \mu_B - \mu_A = 1$ $H_1: \mu_B - \mu_A > 1$	B1	Accept $H_0: \mu_d = 1, \quad H_1: \mu_d > 1.$
	$t = \frac{1.45 - 1}{\sqrt{\frac{s^2}{10}}}$	M1	The – 1 must be present in the numerator.
	0.757	A1	
	Critical value is 1.383: ‘0.757’ < 1.383 Accept H_0	M1	‘0.757’ must come from a paired sample t -calculation, 1.383 must be correct Correct ft conclusion for their 0.757 and 1.383 Condone Reject H_1 ‘Accept H_0 ’ can be implied by a conclusion that is consistent with their 0.757 and 1.383
	Insufficient evidence to support Jade’s claim	A1	Correct work only , ignoring their hypotheses, conclusion in context with level of uncertainty in language. Not ‘prove’ Condone ‘no sufficient’ ‘not enough’ Do not accept statements such as ‘there is sufficient evidence to suggest.....’
		7	

PUBLISHED

Question	Answer	Marks	Guidance
6(b)	Distribution of population differences is normal	B1	Underlying distribution of differences normal B1 'data' implies B0.
		1	
7(a)	$f(x) = \begin{cases} \frac{x}{4}(4-x^2) & 0 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$ $(E(X) =) \int_0^2 \frac{1}{4}(4x^2 - x^4) dx = \frac{1}{4} \left[\frac{4}{3}x^3 - \frac{1}{5}x^5 \right] = \frac{16}{15} \quad (= 1.067)$	B1	
	$(E(\sqrt{X}) =) \int_0^2 \frac{1}{4}(4x^{1.5} - x^{7.2}) dx = \frac{1}{4} \left[\frac{8}{5}x^{2.5} - \frac{2}{9}x^{4.5} \right]_0^2 \quad \left[= \frac{32}{45}\sqrt{2} = 1.005(7) \right]$	M1	Integration, correct powers required, ignore limits.
	$\text{Var}(\sqrt{X}) = E(X) - (E(\sqrt{X}))^2 \text{ used}$	M1	Correct formula used following attempt at integration of both terms.
	$\left[\frac{16}{15} - \left(\frac{32}{45}\sqrt{2} \right)^2 \right] = 0.0553$	A1	$\frac{112}{2025}$
		4	

PUBLISHED

Question	Answer	Marks	Guidance
7(b)	$F(x) = \frac{1}{16}(8x^2 - x^4) \quad [0 \leq x \leq 2]$	B1	
	$G(y) = \frac{1}{16}(8y - y^2) \quad [0 \leq y \leq 4]$	M1	Change variable in their F(x).
	$g(y) = \frac{1}{16}(8 - 2y) \quad [0 \leq y \leq 4]$	M1	Differentiate their G(y) .
	$g(y) = \begin{cases} \frac{1}{16}(8 - 2y) & 0 \leq y \leq 4, \\ 0 & \text{otherwise.} \end{cases}$	A1	Complete and correct.
		4	
7(c)	$\frac{1}{16}(8m - m^2) = \frac{1}{2}$	M1	Equate their G(y) to $\frac{1}{2}$.
	$[m^2 - 8m + 8 = 0,] \quad m = 4 - 2\sqrt{2}$	A1	Single answer in an exact form.
		2	