Cambridge International AS & A Level

FURTHER MATHEMATICS

9231/42

Paper 4 Further Probability & Statistics 42

October/November 2020

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE[™], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© UCLES 2020 Page 2 of 12

Mathematics Specific Marking Principles					
Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation no marks will be awarded for a scale drawing.	is required then				
Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the accuracy is not affected.	he degree of				
Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.					
Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignor	red (isw).				
Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the method required, award all marks earned and deduct just 1 mark for the misread.	ne difficulty or				
Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent	clear.				

© UCLES 2020 Page 3 of 12

PUBLISHED

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- **B** Mark for a correct result or statement independent of method marks.
- DM or DB When a part of a question has two or more 'method' steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
 - FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
 - A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

© UCLES 2020 Page 4 of 12

Cambridge International AS & A Level – Mark Scheme **PUBLISHED**

October/November 2020

Abbreviations

AEF/OE Any Equivalent Form (of answer is equally acceptable) / Or Equivalent

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

CAO Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)

CWO Correct Working Only

ISW Ignore Subsequent Working

SOI Seen Or Implied

SC Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the

light of a particular circumstance)

WWW Without Wrong Working

AWRT Answer Which Rounds To

© UCLES 2020 Page 5 of 12

Question	Answer	Marks	Guidance
1	$s^2 = \frac{313.1}{10} = 31.3$	B1	Can be implied
	$t = \frac{176.2 - 172.5}{s / \sqrt{11}}$	M1	
	= 2.19	A1	
	'2.19' < 2.228	M1	Comparison of their t value with 2.228
	accept H ₀ : insufficient evidence to reject mean height is 172.5 OR sufficient evidence to accept mean height is 172.5	A1 FT	Correct conclusion in context, level of uncertainty in language used. No contradictions. CWO Do not accept: 'mean height is 172.5' Do not accept use of symbols that are undefined e.g. insufficient evidence to reject $\mu = 172.5$
		5	

© UCLES 2020 Page 6 of 12

PUBLISHED						
Question	Answer	Marks	Guidance			
2(a)	H_0 : difference in (population) medians = 0 H_1 : difference in (population) medians < 0 (or >0)	B1 B1	Correct hypotheses (median or m) All notation identified e.g. m = population median if used			
	Differences: -3 4 5 -1 -8 -19 -11 10 -13 -7 9 -16	M1	At most 3 errors			
	Ranks: -2 3 4 -1 -6 -12 -9 8 -10 -5 7 -11	A1	Award for correct rank order, ignore signs			
	Sum T = 3 + 4 + 8 + 7 = 22	A1	cwo			
	Compare with critical value 17: '22' > 17	M1	Compare their T with 17			
	Accept H ₀ Data does not support student's claim	A1	In context, all correct, except possibly second B1 Level of uncertainty in language used. No contradictions.			
		7				
2(b)	Rank for A becomes + 2,	M1	T = 24			
	Changing sign of difference can only reduce evidence in favour of the claim.	A1	still > 17 and test result unchanged			
		2				

© UCLES 2020 Page 7 of 12

Question		Answer								Guidance
3(a)	$\int_{0}^{0.5} \left(\frac{4}{9}x - \frac{1}{9}x^{2}\right) dx \qquad \left[= \frac{1}{9} \left[2 \times 0.5^{2} - \frac{1}{3}0.5^{3}\right]\right]$									Statement of integration with correct limits
	$(=0.0509(26) \text{ or } \frac{11}{216})$									
	Freq = $p = 0.0509(26) \times 200 = 10.19$ AG By addition to 200, $q = 26.85$ $\left(\frac{725}{27}\right)$							A1	Requires sight of 0.0509 or $\frac{11}{216}$	
								B1	Or by integration	
									3	
3(b)	О	5	23	40	41	46	45		M1	At least 4 correct
	Е	10.19	26.85	37.96	43.52	43.52	37.96	=		
	Test statistic = 2.6433 + 0.5520 + 0.1096 + 0.1459 + 0.1413 + 1.3056							13 +		
	4.89 or 4.90								A1	
'4.90' < 11.07:							M1	Compare their value with 11.07		
	PDF is a satisfactory model for the data.					A1 FT	Correct conclusion in context, ft only their 4.90			
							4			

© UCLES 2020 Page 8 of 12

Question	Answer	Marks	Guidance
4(a)	$F(6) - F(3) = \frac{9}{20}$ or 0.45	B1	
		1	
4(b)	$f(x) = \begin{cases} \frac{1}{30}x, & 2 \le x \le 8\\ 0, & otherwise \end{cases}$	B1	May be implied. Only need to see $f(x) = \frac{1}{30}x$
	$E(\sqrt{X}) = \int_{2}^{8} \frac{1}{30} x^{\frac{3}{2}} dx = \left[\frac{1}{75} x^{\frac{5}{2}} \right]$	M1	Integrated
	= 2.34	A1	2.338
			Answer of 2.34 without $\frac{1}{75}x^{\frac{5}{2}}$ scores B1B1 (2/3)
		3	
4(c)	Var(\sqrt{X}) = $\left[\int_{2}^{8} \frac{1}{30} x^{2} dx = \left[\frac{1}{90} x^{3}\right]\right] - E(\sqrt{X})^{2}$	M1	
	$\operatorname{Var}(\sqrt{X}) = \left[\int_{2}^{8} \frac{1}{30} x^{2} dx = \left[\frac{1}{90} x^{3}\right]_{2}^{8}\right] - E(\sqrt{X})^{2}$		
	$= 5.6 - 2.338^{2}$ $= 0.133 \text{ or } 0.134$	A1	
		2	

© UCLES 2020 Page 9 of 12

	1022		
Question	Answer	Marks	Guidance
4(d)	$G(y) = \frac{1}{60}y^{\frac{2}{3}} - \frac{1}{15}$	M1	CDF for Y
	$g(y) = \frac{1}{90} y^{-\frac{1}{3}}$ for $8 \le y \le 512$ (0 otherwise)	M1 A1	
		3	

© UCLES 2020 Page 10 of 12

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	$P(X = r) = {}^{n}C_{r}p^{r} (1-p)^{n-r} \text{ or } {}^{n}C_{r}p^{r}q^{n-r}$	B1	
	$G_X(t) = \sum_{0}^{n} n\operatorname{Cr} p^{r} (1-p)^{n-r} t^{r}$	M1	Accept minimum of 4 terms including the last. Shown with specific value of n is M0
	$\sum_{0}^{n} \operatorname{nCr}(pt)^{r} (1-p)^{n-r}$ $= (q+pt)^{n}$	A1	At least one intermediate step to be shown, with <i>p</i> and <i>t</i> grouped AG
		3	
5(b)	$G'_X(t) = n(q+pt)^{n-1} \times p$	M1	
	So $E(X) = G'_X(I) = np(q+p)^{n-1}$ and $q+p=1$ so $E(X) = np$	A1	
	$G''_X(t) = n(n-1)(q+pt)^{n-2} \times p \times p$	M1	
	$Var(X) = n(n-1)p^{2} + np - (np)^{2}$	M1	
	np(1-p)	A1	
		5	

© UCLES 2020 Page 11 of 12

PUBLISHED Question Marks Guidance Answer 6(a) **B1** $\overline{x} = \frac{1}{2}(4.22 + 3.36) = 3.79$ Using z value implies M0 $4.22 - 3.36 = \frac{2ts}{\sqrt{10}}$ With t = 1.833**A1** (s = 0.7418) variance = 0.55(0)**A1** 4 **B1** Not \overline{x} 6(b) H_0 : $\mu_A = \mu_B$ H_1 : $\mu_A > \mu_B$ $=\frac{1}{7}(98.02-97.02)$ Pooled estimate = $\frac{9 \times 0.550316 + 7 \times 0.142507}{}$ **M1** =0.372**A1** M1A1 '1.063' < 1.337 oe Compare their value with 1.337 M1Accept H₀ A1ft Correct conclusion in context, level of uncertainty in language used. No contradictions. Nassa's claim is not supported

8

© UCLES 2020 Page 12 of 12