



# Cambridge International AS & A Level

---

**FURTHER MATHEMATICS****9231/42**

Paper 4 Further Probability &amp; Statistics

**October/November 2022**

MARK SCHEME

Maximum Mark: 50

---

<b>Published</b>
------------------

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2022 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

---

This document consists of **13** printed pages.

**PUBLISHED****Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**PUBLISHED**

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

**PUBLISHED****Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

**Types of mark**

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
  - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
  - The total number of marks available for each question is shown at the bottom of the Marks column.
  - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
  - Square brackets [ ] around text or numbers show extra information not needed for the mark to be awarded.

**Abbreviations**

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

**PUBLISHED**

Question	Answer	Marks	Guidance
1	$2.02 = \bar{x} + 1.833\sqrt{\frac{s^2}{10}}$ $1.78 = \bar{x} - 1.833\sqrt{\frac{s^2}{10}}$ Add, $\bar{x} = \frac{2.02 + 1.78}{2} = 1.90$	<b>M1</b>	Add.
	$\sum x = 19$	<b>A1</b>	
	Subtract: $2.02 - 1.78 = 2 \times 1.833\sqrt{\frac{s^2}{10}}$	<b>M1</b>	Allow 1.372, 1.383, 1.812 instead of 1.833.
	$s^2 = 0.042859$	<b>A1</b>	May be implied. $s = 0.20702$
	But $s^2 = \frac{1}{9}(\sum x^2 - \frac{(\sum x)^2}{10})$	<b>M1</b>	
	$\sum x^2 = 9s^2 + \frac{19^2}{10} = 36.5$	<b>A1</b>	36.486
			Using 1.645: maximum M1A1 M0A0 M1A0 3/6
		<b>6</b>	

**PUBLISHED**

Question	Answer	Marks	Guidance									
2	$H_0$ : performance is independent of region $H_1$ : performance is not independent of region	<b>B1</b>										
	<table border="1" data-bbox="434 316 1032 512"> <tbody> <tr> <td data-bbox="434 316 633 379">62 (53.9)</td> <td data-bbox="633 316 833 379">41 (44.1)</td> <td data-bbox="833 316 1032 379">44 (49)</td> </tr> <tr> <td data-bbox="434 379 633 443">102 (106.7)</td> <td data-bbox="633 379 833 443">94 (87.3)</td> <td data-bbox="833 379 1032 443">95 (97)</td> </tr> <tr> <td data-bbox="434 443 633 512">56 (59.4)</td> <td data-bbox="633 443 833 512">45 (48.6)</td> <td data-bbox="833 443 1032 512">61 (54)</td> </tr> </tbody> </table>	62 (53.9)	41 (44.1)	44 (49)	102 (106.7)	94 (87.3)	95 (97)	56 (59.4)	45 (48.6)	61 (54)	<b>M1 A1</b>	Calculate expected frequencies, allow 1 error.
62 (53.9)	41 (44.1)	44 (49)										
102 (106.7)	94 (87.3)	95 (97)										
56 (59.4)	45 (48.6)	61 (54)										
	Test stat = 1.217 + 0.218 + 0.510 + 0.207 + 0.514 + 0.041 + 0.195 + 0.267 + 0.907	<b>M1</b>	Calculate test statistic.									
	4.08	<b>A1</b>	SCB2 for 4.08 if totally unsupported. SCM1A1 B1 for 4.08 supported only by expected frequencies.									
	Tabular value, 4df = 7.779 1.08 < 7.779 Accept $H_0$	<b>M1</b>	Compare with 7.779 and conclusion.									
	Insufficient evidence that performance is not independent of region	<b>A1</b>	Correct conclusion in context, following correct work, level of uncertainty in language. 'Prove' gives A0.									
		<b>7</b>										

**PUBLISHED**

Question	Answer	Marks	Guidance
3	$s_x^2 = \frac{1}{49} \left( 115.2 - \frac{75.5^2}{50} \right) = 0.02439$ $s_y^2 = \frac{1}{79} \left( 172.6 - \frac{116.8^2}{80} \right) = 0.02623$	<b>M1</b>	$\frac{239}{9800}$ $\frac{259}{9875}$
		<b>A1</b>	Both correct
	$s^2 = \frac{0.02439}{50} + \frac{0.02623}{80} \quad [= 0.00081565]$	<b>M1 A1</b>	May be implied by 1.75 for $z$
	$z = \frac{\frac{75.5}{50} - \frac{116.8}{80}}{s}$	<b>M1</b>	
	1.75	<b>A1</b>	
	Compare with 1.645: $1.75 > 1.645$ Reject null hypothesis	<b>M1</b>	Using areas, $0.04 < 0.05$ .
	Sufficient evidence to suggest that population mean in country $X$ is greater than population mean in country $Y$	<b>A1</b>	Correct conclusion in context, following correct work, level of uncertainty in language. 'Prove' gives A0.
		<b>8</b>	Pooled variance: $z = 1.74$ M1A1 M0A0M0A0 M1A0 max 3/8



**PUBLISHED**

Question	Answer	Marks	Guidance
4(a)	$\int_0^1 k dx + \int_1^2 kx dx = [kx] + \left[\frac{k}{2}x^2\right], \quad k + \frac{3}{2}k = 1, \quad k = \frac{2}{5}$	<b>B1</b>	Integration and $k + 3k/2 = 1$ oe seen.
		<b>1</b>	
4(b)	$F(x) = \begin{cases} kx & 0 \leq x < 1, \\ \frac{k}{2}x^2 + \frac{1}{5} & 1 \leq x \leq 2 \end{cases}$	<b>M1</b>	CDF or integration with $\frac{1}{5}x^2 + \frac{1}{5}$ seen.
	UQ: $\frac{k}{2}x^2 + \frac{1}{5} = \frac{3}{4},$	<b>M1</b>	Method for UQ
	$\left[ x^2 + 1 = 5 \times \frac{3}{4}, \quad x^2 = \frac{11}{4} \right] \quad x = \frac{1}{2}\sqrt{11}$	<b>A1</b>	UQ
	LQ: $kx = \frac{1}{4}, \quad x = \frac{5}{8}$	<b>B1</b>	LQ
	$\text{IQR} = \frac{1}{2}\sqrt{11} - \frac{5}{8} = 1.03(3)$	<b>A1</b>	CAO
		<b>5</b>	

**PUBLISHED**

Question	Answer	Marks	Guidance
4(c)	$E(X) = \int_0^1 kx \, dx + \int_1^2 kx^2 \, dx = \left[ \frac{k}{2}x^2 \right] + \left[ \frac{k}{3}x^3 \right] = \frac{17}{15}$	<b>M1</b>	With correct limits.
	$E(X^2) = \int_0^1 kx^2 \, dx + \int_1^2 kx^3 \, dx = \left[ \frac{k}{3}x^3 \right] + \left[ \frac{k}{4}x^4 \right] = \frac{49}{30}$	<b>M1</b>	With correct limits.
	$\text{Var}(X) = \frac{49}{30} - \left( \frac{17}{15} \right)^2$	<b>M1</b>	Using correct formula with numerical values.
	$\frac{157}{450} \quad (= 0.349)$	<b>A1</b>	
		<b>4</b>	

Question	Answer	Marks	Guidance
5(a)	$G_x(t) = \frac{9}{16} + \frac{6}{16}t + \frac{1}{16}t^2$	<b>M1 A1</b>	2 probabilities correct, in a polynomial
		<b>2</b>	

**PUBLISHED**

Question	Answer	Marks	Guidance
5(b)	$G_Z(t) = \left(\frac{9}{16} + \frac{6}{16}t + \frac{1}{16}t^2\right)\left(\frac{25}{36} + \frac{10}{36}t + \frac{1}{36}t^2\right)$	<b>M1</b>	Second PGF correct and multiplied by part (a).
	$\frac{1}{576}(225 + 240t + 94t^2 + 16t^3 + t^4)$	<b>M1</b>	Obtains a quartic polynomial.
	Or $\frac{25}{64} + \frac{5}{12}t + \frac{47}{288}t^2 + \frac{1}{36}t^3 + \frac{1}{576}t^4$	<b>A1</b>	Note: $\frac{1}{576}(t+3)^2(t+5)^2$ scores M1M1A0.
		<b>3</b>	
5(c)	$G_Z'(t) = \frac{1}{576}(240 + 188t + 48t^2 + 4t^3)$ $G_Z''(t) = \frac{1}{576}(188 + 96t + 12t^2)$	<b>M1</b>	Differentiate twice.
	$\text{Var}(Z) = \frac{1}{576}(188 + 96 + 12) + \frac{5}{6} - \frac{25}{36}$	<b>M1</b>	Use correct formula.
	$\frac{47}{72}$	<b>A1</b>	
	OR: $\text{Var}(Z) = 2 \times \frac{1}{4} \times \frac{3}{4} + 2 \times \frac{1}{6} \times \frac{5}{6}$	<b>M1</b> <b>M1</b>	One term correct. Two terms present and added.
	$\frac{47}{72}$	<b>A1</b>	
		<b>3</b>	
5(d)	1	<b>B1 FT</b>	FT their power with greatest coefficient in part (b).
		<b>1</b>	

**PUBLISHED**

Question	Answer	Marks	Guidance																																																
6(a)	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">359</td> <td style="width: 25%;">3</td> <td style="width: 25%;">342</td> <td style="width: 25%;">1</td> </tr> <tr> <td>374</td> <td>5</td> <td>355</td> <td>2</td> </tr> <tr> <td>398</td> <td>7</td> <td>361</td> <td>4</td> </tr> <tr> <td>415</td> <td>10</td> <td>384</td> <td>6</td> </tr> <tr> <td>427</td> <td>11</td> <td>401</td> <td>8</td> </tr> <tr> <td>452</td> <td>14</td> <td>414</td> <td>9</td> </tr> <tr> <td>461</td> <td>16</td> <td>437</td> <td>12</td> </tr> <tr> <td>482</td> <td>18</td> <td>443</td> <td>13</td> </tr> <tr> <td>502</td> <td>20</td> <td>454</td> <td>15</td> </tr> <tr> <td>512</td> <td>22</td> <td>472</td> <td>17</td> </tr> <tr> <td>545</td> <td>23</td> <td>491</td> <td>19</td> </tr> <tr> <td>612</td> <td>24</td> <td>506</td> <td>21</td> </tr> </table>	359	3	342	1	374	5	355	2	398	7	361	4	415	10	384	6	427	11	401	8	452	14	414	9	461	16	437	12	482	18	443	13	502	20	454	15	512	22	472	17	545	23	491	19	612	24	506	21	<b>M1</b>	<p>Wrong test: max B1 for hypotheses, B1B1 for correct mean and variance</p> <p>Attempt at rankings (allow up to 4 errors)</p>
359	3	342	1																																																
374	5	355	2																																																
398	7	361	4																																																
415	10	384	6																																																
427	11	401	8																																																
452	14	414	9																																																
461	16	437	12																																																
482	18	443	13																																																
502	20	454	15																																																
512	22	472	17																																																
545	23	491	19																																																
612	24	506	21																																																
	Test statistic = 127	<b>A1</b>	Clearly identified																																																
	$H_0$ : population medians are equal $H_1$ : population median for $X$ is greater than population median for $Y$	<b>B1</b>																																																	
	Mean = $\frac{1}{2} \times 12 \times 25 = 150$	<b>B1</b>																																																	
	Variance = $\frac{1}{12} \times 12 \times 12 \times 25 = 300$	<b>B1</b>																																																	

**PUBLISHED**

<b>Question</b>	<b>Answer</b>	<b>Marks</b>	<b>Guidance</b>
6(a)	$\frac{127.5 - 150}{\sqrt{300}}$	<b>M1</b>	Allow incorrect or no continuity correction.
	-1.299	<b>A1</b>	
	Compare with $-1.645$ : $-1.299 > -1.645$ , or $0.097 > 0.05$ Accept $H_0$	<b>M1</b>	Valid comparison with 1.645 or 0.05 and reach correct ft conclusion.
	Insufficient evidence to support manager's claim	<b>A1</b>	Correct conclusion in context, following correct work, level of uncertainty in language. 'Prove' is A0.
		<b>9</b>	
6(b)	Not appropriate/no, not the same people	<b>B1</b>	OE No and reason needed, e.g. individuals in the samples cannot be paired up.
		<b>1</b>	