

Cambridge  
International  
**A Level**

**Cambridge International Examinations**  
Cambridge International Advanced Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**FURTHER MATHEMATICS**

**9231/13**

Paper 1

**May/June 2017**

**3 hours**

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF10)

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of a calculator is expected, where appropriate.

Results obtained solely from a graphic calculator, without supporting working or reasoning, will not receive credit.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **24** printed pages.



1 The roots of the cubic equation  $x^3 + 2x^2 - 3 = 0$  are  $\alpha$ ,  $\beta$  and  $\gamma$ .

(i) By using the substitution  $y = \frac{1}{x^2}$ , find the cubic equation with roots  $\frac{1}{\alpha^2}$ ,  $\frac{1}{\beta^2}$  and  $\frac{1}{\gamma^2}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence find the value of  $\frac{1}{\alpha^2} + \frac{1}{\beta^2} + \frac{1}{\gamma^2}$ . [1]

.....

.....

.....

.....

(iii) Find also the value of  $\frac{1}{\alpha^2\beta^2} + \frac{1}{\beta^2\gamma^2} + \frac{1}{\gamma^2\alpha^2}$ . [1]

.....

.....

.....

.....

.....

## 3

2 (i) Verify that  $\frac{2r+1}{r(r+1)(r+2)} = \frac{1}{2} \left\{ \frac{(2r+1)(2r+3)}{(r+1)(r+2)} - \frac{(2r-1)(2r+1)}{r(r+1)} \right\}$ . [2]

.....

.....

.....

.....

.....

.....

(ii) Hence show that  $\sum_{r=1}^n \frac{2r+1}{r(r+1)(r+2)} = \frac{1}{2} \left\{ \frac{(2n+1)(2n+3)}{(n+1)(n+2)} - \frac{3}{2} \right\}$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Deduce the value of  $\sum_{r=1}^{\infty} \frac{2r+1}{r(r+1)(r+2)}$ . [2]

.....

.....

.....

.....

.....



4 A curve  $C$  has equation  $x^3 - 3xy + y^2 = 4$ . Find the value of  $\frac{d^2y}{dx^2}$  at the point  $(0, 2)$  of  $C$ . [7]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

5 A curve  $C$  has parametric equations

$$x = \frac{2}{5}t^{\frac{5}{2}} - 2t^{\frac{1}{2}}, \quad y = \frac{4}{3}t^{\frac{3}{2}}, \quad \text{for } 1 \leq t \leq 4.$$

(i) Find the exact value of the arc length of  $C$ .

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Find also the exact value of the surface area generated when  $C$  is rotated through  $2\pi$  radians about the  $x$ -axis. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 Let  $I_n$  denote  $\int_0^2 (4 + x^2)^{-n} dx$ .

(i) Find  $\frac{d}{dx} (x(4 + x^2)^{-n})$  and hence show that

$$8nI_{n+1} = (2n - 1)I_n + 2 \times 8^{-n}. \tag{5}$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



9

- (ii) Use the result for integrating  $\frac{1}{x^2 + a^2}$  with respect to  $x$ , in the List of Formulae (MF10), to find the value of  $I_1$  and deduce that

$$I_3 = \frac{3}{1024}\pi + \frac{1}{128}. \quad [5]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 10

7 (i) Use de Moivre's theorem to prove that

$$\tan 4\theta = \frac{4 \tan \theta - 4 \tan^3 \theta}{1 - 6 \tan^2 \theta + \tan^4 \theta} \quad [5]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(ii)** Hence find the solutions of the equation

$$t^4 - 4t^3 - 6t^2 + 4t + 1 = 0,$$

giving your answers in the form  $\tan k\pi$ , where  $k$  is a rational number.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8 Find the solution of the differential equation

$$\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 9x = 18t^2 + 6t + 1,$$

given that, when  $t = 0$ ,  $x = 3$  and  $\frac{dx}{dt} = 0$ . [10]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



9 The plane  $\Pi_1$  passes through the points  $(1, 2, 1)$  and  $(5, -2, 9)$  and is parallel to the vector  $\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ .

(i) Find the cartesian equation of  $\Pi_1$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The plane  $\Pi_2$  contains the lines

$$\mathbf{r} = 2\mathbf{i} - 3\mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} - 2\mathbf{j} - \mathbf{k}) \quad \text{and} \quad \mathbf{r} = 2\mathbf{i} - 3\mathbf{j} + \mathbf{k} + \mu(2\mathbf{i} + 3\mathbf{j} - \mathbf{k}).$$

(ii) Find the cartesian equation of  $\Pi_2$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the acute angle between  $\Pi_1$  and  $\Pi_2$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 The matrix  $\mathbf{A}$  is given by

$$\mathbf{A} = \begin{pmatrix} 6 & -8 & 7 \\ 7 & -9 & 7 \\ 6 & -6 & 5 \end{pmatrix}.$$

- (i) Given that  $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$  is an eigenvector of  $\mathbf{A}$ , find the corresponding eigenvalue. [2]

.....

.....

.....

.....

.....

.....

- (ii) Given also that  $-1$  is an eigenvalue of  $\mathbf{A}$ , find a corresponding eigenvector. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....







11 Answer only **one** of the following two alternatives.

**EITHER**

A curve  $C$  has polar equation  $r = 2a \cos(2\theta + \frac{1}{2}\pi)$  for  $0 \leq \theta < 2\pi$ , where  $a$  is a positive constant.

(i) Show that  $r = -2a \sin 2\theta$  and sketch  $C$ . [4]

.....

.....

.....

(ii) Deduce that the cartesian equation of  $C$  is

$$(x^2 + y^2)^{\frac{3}{2}} = -4axy. \quad [2]$$

.....

.....

.....

.....

.....

.....

.....

(iii) Find the area of one loop of  $C$ .

[5]

A series of horizontal dotted lines for writing the solution.



OR

The matrix  $\mathbf{A}$ , given by

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 0 & 2 \\ 3 & -1 & 4 & 0 \\ 5 & -8 & -6 & 19 \\ -2 & 3 & 2 & -7 \end{pmatrix},$$

represents a transformation from  $\mathbb{R}^4$  to  $\mathbb{R}^4$ .

- (i) Find the rank of  $\mathbf{A}$  and show that  $\left\{ \begin{pmatrix} 2 \\ 2 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}$  is a basis for the null space of the transformation. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that if

$$\mathbf{Ax} = p \begin{pmatrix} 1 \\ 3 \\ 5 \\ -2 \end{pmatrix} + q \begin{pmatrix} -1 \\ -1 \\ -8 \\ 3 \end{pmatrix},$$

where  $p$  and  $q$  are given real numbers, then

$$\mathbf{x} = \begin{pmatrix} p + 2\lambda + \mu \\ q + 2\lambda + 3\mu \\ -\lambda \\ \mu \end{pmatrix},$$

where  $\lambda$  and  $\mu$  are real numbers. [2]

.....

.....

.....

.....

.....

.....

(iii) Find the values of  $p$  and  $q$  such that

$$p \begin{pmatrix} 1 \\ 3 \\ 5 \\ -2 \end{pmatrix} + q \begin{pmatrix} -1 \\ -1 \\ -8 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 7 \\ 18 \\ -7 \end{pmatrix}. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

(iv) Find the solution of the equation  $\mathbf{Ax} = \begin{pmatrix} 3 \\ 7 \\ 18 \\ -7 \end{pmatrix}$  of the form  $\mathbf{x} = \begin{pmatrix} 4 \\ 9 \\ \alpha \\ \beta \end{pmatrix}$ , where  $\alpha$  and  $\beta$  are positive integers to be found. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.