

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**GCE Advanced Level**

## **MARK SCHEME for the May/June 2014 series**

### **9231 FURTHER MATHEMATICS**

**9231/22**

Paper 2, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – May/June 2014</b>	<b>9231</b>	<b>22</b>

Question Number	Mark Scheme Details		Part Mark	Total	
<b>1</b>	Equate impulse to momentum to find initial speed $v$ and Newton's law of restitution to find new speed:	$v = 4u, v' = ev = [-] 3u$	M1 A1	2	<b>2</b>
<b>2</b>	Find $v^2$ at both $A$ and $B$ :  Find amplitude $a$ m from given K.E. ratio:  Find $\omega$ from $v_{\max} = a\omega$ : Find time ( $\sqrt{\phantom{x}}$ on $a$ ) at $A$ or at $B$ , e.g.: Combine correctly to find time from $A$ to $B$ :  Evaluate to 3 d.p.:	$v_A^2 = \omega^2(a^2 - 0.5^2)$ and $v_B^2 = \omega^2(a^2 - 0.75^2)$  $\frac{1}{2}mv_A^2 = (12/11) \frac{1}{2}mv_B^2$  $11(a^2 - 0.5^2) = 12(a^2 - 0.75^2)$  $a^2 = \frac{1}{4}(27 - 11) = 4, a = 2$  $0.6 = 2\omega, \omega = 0.3$  $\omega^{-1} \sin^{-1}(0.5/2)$ or $\omega^{-1} \cos^{-1}(0.5/2)$  $\omega^{-1} \sin^{-1}(0.75/2)$ or $\omega^{-1} \cos^{-1}(0.75/2)$  $\omega^{-1} \sin^{-1}(0.75/2) - \omega^{-1} \sin^{-1}(0.5/2)$  or $\omega^{-1} \cos^{-1}(0.5/2) - \omega^{-1} \cos^{-1}(0.75/2)$  $= \omega^{-1}(0.3844 - 0.2527)$ or $\omega^{-1}(1.318 - 1.186)$  $= 1.2813 - 0.8423$  $4.3937 - 3.9547 = 0.439$ [s]	B1  M1 A1  B1  M1 A1 $\sqrt{\phantom{x}}$  M1  A1	3	<b>8</b>

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

3	Use conservation of momentum, e.g.:	$mv_A + 9mv_B = mu$	M1		
	Use Newton's law of restitution (consistent signs):	$v_B - v_A = eu$	M1		
	Relate $v_A$ to $v_B$ using K.E. (A.E.F.):	$\frac{1}{2}mv_A^2 + \frac{1}{2}9mv_B^2 = \frac{1}{2}mu^2$	M1		
	Combine two eqns to find $v_A$ and $v_B$ e.g.:	$v_A = (1 - 9e)u/10, v_B = (1 + e)u/10$ or $v_A, v_B = -u/2, u/6$ [or $7u/10, u/30$ ]	M1 A1		
	Use in 3rd eqn to find $e$ , e.g.:	$(1 - 9e)^2 + 9(1 + e)^2 = 50$			
	(A0 if finally $\pm\frac{2}{3}$ )	$90e^2 = 40, e = \frac{2}{3}$	M1 A1	7	
	Use Newton's law of restitution with	$v_C = 2v_B', \text{ e.g.: } v_C - v_B' = ev_B, v_B' = \frac{2}{3}v_B$ [ $v_B = u/6, v_B = u/9, v_C = 2u/9$ ]	B1		
Use conservation of momentum to find $k$ :	$9mv_B' + kmv_C = 9mv_B$ $9v_B' + 2kv_B' = 13.5v_B', k = 9/4$	M1 A1	3	<b>10</b>	

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

4	(i)	Use conservation of energy at lowest point: Use $F = ma$ radially at lowest point: Eliminate $v^2$ to find $R$ [ $v^2 = 2.3 ga$ ]:	$\frac{1}{2}mv^2 = \frac{1}{2}mu^2 + mga$ $R - mg = mv^2/a$ $R = mu^2/a + 3mg = 3.3mg$	B1 B1 B1	3	10	
	(ii)	Use conservation of energy at $B$ to find $V_B$ :  (A.E.F.)	$\frac{1}{2}mV_B^2 = \frac{1}{2}mu^2 + mga \sin \theta$ $V_B^2 = (0.3 + 0.5)ga, V_B = \sqrt{(0.8ga)}$ or $2\sqrt{(ga/5)}$ or $0.894\sqrt{(ga)}$	M1A1  A1			3
	(iii)	Use vertical component $v_B$ of speed $V_B$ at $B$ : Find height $h$ reached above $B$ : Find height $h$ reached above level of $O$ :	$v_B = V_B \cos \theta$ [= $\frac{1}{4}\sqrt{15}V_B = \sqrt{(3/4ga)}$ ] $h = v_B^2/2g = 3a/8$ $h - a \sin \theta = 3a/8 - \frac{1}{4}a = a/8$ <b>A.G.</b>	M1 M1 A1 A1			4
5	Find MI of components about $A$ :  (M1 for $BC$ or $CD$ )	Glass $(3M/5) \{ \frac{1}{3}(5a)^2 + 25a^2 \} = 20Ma^2$ $AB$ $M\{ \frac{1}{3}(4a)^2 + (4a)^2 \} = 64Ma^2/3$ $AD$ $\frac{1}{3}M\{ \frac{1}{3}(3a)^2 + (3a)^2 \} = 4Ma^2$ $BC$ $\frac{1}{3}M\{ \frac{1}{3}(3a)^2 + 73a^2 \} = 76Ma^{2/3}$ $CD$ $M\{ \frac{1}{3}(4a)^2 + 52a^2 \} = 172Ma^{2/3}$	M1 A1 B1 B1 M1 A1 A1	8	13		
	Find total MI about $A$ : (OR can first find total MI about centre of mass) State or imply total mass acts at mid-point of $AC$	$I = 128Ma^2$ A.G.	A1  M1				
	Use eqn of circular motion to find $d^2\theta/dt^2$ : Approximate $\sin \theta$ by $\theta$ and substitute for $I$ :	$I d^2\theta/dt^2 = [-] (49Mg/15) 5a \sin \theta$ $d^2\theta/dt^2 = -(49g/384a) \theta$	M1 A1 A1				
	Find period $T = 2\pi/\omega$ with $\omega = \sqrt{(49g/384a)}$ :	$T = 2\pi\sqrt{(384a/49g)}$ or $(16\pi/7)\sqrt{(6a/g)}$ or $17.6\sqrt{(a/g)}$ (A.E.F.)	B1			5	

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

6		State or find the expected value of $X$ : using $p = \frac{1}{4}$ :	$E(X) = 1/p = 1/\frac{1}{4} = 4$	B1	1	5
	(i)	Find $P(X = 4)$ :	$P(X = 4) = (\frac{3}{4})^3 \frac{1}{4} = 27/256$ or 0.105	M1 A1	2	
	(ii)	Find $P(X < 6)$ :	$P(X < 6) = 1 - (\frac{3}{4})^5$ or $\{1 + \frac{3}{4} + (\frac{3}{4})^2 + (\frac{3}{4})^3 + (\frac{3}{4})^4\} \frac{1}{4}$ $= 781/1024$ or 0.763	M1 A1	2	
		S.R. Using $p = \frac{1}{2}$ can earn B0 M1 A0 M0 A0				
7	(i)	State probability density function of $T$ :	$f(t) = 0.001 \exp(-0.001t) \quad (t \geq 0)$ [ = 0 (otherwise or $t < 0$ )]	B1	1	8
	(ii)	Find $P(T > 2000)$ : S.R. $1 - e^{-2} = 0.865$ earns B1 only (max 1/3) State inequality for $t$ (lose A1 if = or $\leq$ ): Solve for $t_{\max}$ : (Omitting power 10 earns 0/4; using $1 - (\exp(-0.001t))^{10}$ can earn M1 A0 M1 A0 only)	$P(t > 2000) = 1 - F(2000)$ $= 1 - (1 - e^{-2}) = e^{-2}$ or 0.135 $(\exp(-0.001t))^{10} \geq [or >] 0.9$ $t_{\max} = (\ln 0.9) / (-0.01) = 10.5$	M1 M1 A1 M1 A1 M1 A1	3 4	

Page 6	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

8	<p>State hypotheses (B0 for <math>\bar{\chi}</math> ...):</p> <p>Estimate both popln. variances using two samples: (allow use of biased: <math>\sigma_{X,60}^2 = 236</math> or <math>15.36^2</math>)</p> <p>(allow use of biased: <math>\sigma_{Y,50}^2 = 265</math> or <math>16.28^2</math>)</p> <p>Estimate population variance for combined sample:</p> <p>(allow <math>\sigma_{X,60}^2/60 + \sigma_{Y,50}^2/50</math>: <math>9.233</math> or <math>3.039^2</math>)</p> <p>Calculate value of <math>z</math> (to 2 d.p., either sign):</p> <p>State or use correct tabular <math>z</math> – value (to 2 d.p.): (or can compare 6 with e.g. <math>2.326</math> <math>s = 7.13</math> or <math>7.07</math>)</p> <p>Correct conclusion (A.E.F, <math>\checkmark</math> on <math>z</math> – values):</p> <p><b>S.R.</b> Assuming equal population variances: Find pooled estimate of common variance <math>s^2</math>:</p> <p>Calculate value of <math>z</math> (to 2 d.p., either sign):</p> <p>Tabular value; conclusion</p>	<p><math>H_0: \mu_X = \mu_Y, H_1: \mu_X \neq \mu_Y</math></p> <p><math>S_x^2 = (626220 - 6060^2/60) / 59</math> [= <math>240</math> or <math>15.49^2</math>]</p> <p>And <math>s_y^2 = (464500 - 4750^2/50) / 49</math> [= <math>270.4</math> or <math>16.44^2</math>]</p> <p><math>s^2 = s_x^2/60 + s_y^2/50</math> = <math>9.408</math> or <math>3.067^2</math></p> <p><math>z = (101 - 95) / s</math> = <math>6/3.067 = 1.96</math> (or <math>1.97</math>)</p> <p><math>z_{0.99} = 2.326</math> or <math>2.33</math> (allow <math>2.36</math>)</p> <p>[Accept <math>H_0</math>] Claims are the same Hypotheses; Explicit assumption : <math>s^2 = (626220 - 6060^2/60 +</math> <math>464500 - 4750^2/50) / 108</math></p> <p><math>z = 6 / s\sqrt{(1/60+1/50)} = 1.97</math> = <math>253.8</math> or <math>15.93^2</math></p> <p>As above )</p>	<p>B1</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1 A1 B1</p> <p>B1<math>\checkmark</math> (B1; B1)</p> <p>(M1 A1)</p> <p>(M1 A1) (A1)</p> <p>(B1; B1<math>\checkmark</math>)</p>	<p>9</p>
---	---	---	---	----------

Page 7	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

<b>9</b>	<p>Find expected frequency <math>p</math>:</p> <p>Find <math>q</math> by similar method <i>or</i> by using total of 200:</p> <p>State (at least) null hypothesis: Calculate <math>\chi^2</math> (to 3 s.f.):</p> <p>State or use correct tabular <math>\chi^2</math> value (to 3 s.f.): Valid method for reaching conclusion: Conclusion consistent with correct values (A.E.F):</p>	$p = 200 \int_2^3 (1/x \ln 8) dx$ $= (200 / \ln 8) [\ln x]_2^3$ $= 200 \times 0.1950 = 39.00 \text{ A.G.}$ $q = 21.46 \text{ or } 21.45$ $H_0: f(x) \text{ fits data (A.E.F.)}$ $\chi^2 = 0.202 + 0.923 + 0.678 + 0.584$ $+ 1.134 + 4.134 + 3.644 = 11.3$ $\chi_{6,0.95}^2 = 12.59$ <p>Accept <math>H_0</math> if <math>\chi^2 \leq</math> tabular value Distribution fits observations</p>	<p>M1A1 M1A1</p> <p>B1</p> <p>M1A1 B1 M1 A1</p>	<p>4</p> <p>6</p> <p><b>10</b></p>
<b>10</b>	<p>Find correlation coefficient <math>r</math>: (A.E.F.; A0 if only 3 s.f. clearly used)</p> <p>State both hypotheses (B0 for <math>r \dots</math>): State or use correct tabular two-tail <math>r</math>-value: Valid method for reaching conclusion: Correct conclusion (A.E.F, dep *A1, *B1): Calculate gradient <math>p</math> in <math>x - \bar{x} = p(y - \bar{y})</math>: Find regression line of <math>x</math> on <math>y</math>:</p>	$r = (73\,527 - 866 \times 639 / 10) / \sqrt{\{(121\,276 - 866^2 / 10)(55\,991 - 639^2 / 10)\}}$ $= 18\,189.6 / \sqrt{(46\,280.4 \times 15\,158.9)}$ $= 0.687$ $H_0: \rho = 0, H_1: \rho \neq 0$ $r_{10,5\%} = 0.632$ <p>Reject <math>H_0</math> if <math> r  &gt;</math> tabular value There is non-zero correlation</p> $p = 18\,189.6 / 15\,158.9 = 1.20$ $x = 86.6 + 1.20(y - 63.9)$ $= 1.20y + 9.92$	<p>M1 A1 A1 *A1</p> <p>B1 *B1 M1 A1 B1</p> <p>M1 A1</p>	<p>4</p> <p>4</p> <p>3</p> <p><b>11</b></p>

Page 8	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

11 A	(i)	Use Pythagoras to find $AB$ :	$AB = \sqrt{(4a^2 + 12a^2)} = 4a$	A.G.	M1 A1		
	(ii)	Find $\angle SAB$ :	$\angle CAB = \sin^{-1} 2a\sqrt{3}/4a$ or $\cos^{-1} 2a/4a$ or $\tan^{-1} 2a\sqrt{3}/2a$ $= 60^\circ$ so $\angle SAB = 30^\circ$	A.G.	M1 A1	4	
		<i>EITHER</i>					
		Resolve vertically and horizontally, e.g.:	$\frac{1}{2} N_A + \frac{1}{2}\sqrt{3} N_B + \frac{1}{2}\sqrt{3} F_A = W$				
		( $F_A$ may be in either direction)	and $\frac{1}{2}\sqrt{3} N_A = \frac{1}{2} N_B + \frac{1}{2} F_A$		M1 A1		
		Eliminate $N_B + F_A$ to find $N_A$ :	$N_A = \frac{1}{2} W$	A.G.	A1		
		<i>OR</i>				3	
	(iii)	Resolve in dirn. $PQ$ to find $N_A$ :	$N_A = \frac{1}{2} W$	A.G.	(M1 A1)		
		Second resolution, e.g. in dirn. $PS$ :	$N_B + F_A = \frac{1}{2}\sqrt{3} W$		(A1)		
		Take moments, e.g. about $A$ :	$\frac{1}{2}\sqrt{3} W \times 3a/2 + \frac{1}{2} W \times (2\sqrt{3} - 3)a$ $= N_B \times 2a$		M1 A1 A1		
		(A1 for each side of eqn)					
		Solve to find $N_B$ :	$N_B = \{(7\sqrt{3} - 6)/8\} W$		M1 A1		
		Use $N_B$ to find $F_A$ :	$F_A = \sqrt{3} N_A - N_B$ or $\frac{1}{2}\sqrt{3} W - N_B$ $= \{3(2 - \sqrt{3})/8\} W$ (A.E.F.)		M1 A1	7	14



Page 9	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – May/June 2014	9231	22

<b>B</b>	Estimate population variance:	$s_P^2 = (236.0 - 42 \cdot 8^2 / 8) / 7$			
	(allow biased here: 0.8775 or 0.9367 <sup>2</sup> )	$= 351 / 350$ or 1.003 or 1.001 <sup>2</sup>	M1		
	Find confidence interval (allow $z$ in place of $t$ ) e.g.:	$42.8 / 8 \pm t \sqrt{s_P^2 / 8}$	M1		
	Use correct tabular $t$ -value:	$t_{7, 0.975} = 2.365$	A1		
	Evaluate C.I. correct to 2 d.p.:	$5.35 \pm 0.84$ or [4.51, 6.19]	A1	4	
	Formulate inequality for $k$ (or equality for $k_{\max}$ ):	$(5.35 - k) / \sqrt{s_P^2 / 8} \geq [\text{or } >] t$	M1		
	Use correct tabular $t$ -value:	$t_{7, 0.9} = 1.415$	A1		
	Solve for $k_{\max}$ (A0 if = or $\leq$ was used for $k$ above):	$5.35 - k \geq 0.50, k_{\max} = 4.85$	A1	3	
	State hypotheses (B0 for $\bar{x}$ ...), e.g.:	$H_0: \mu_P = \mu_Q, H_1: \mu_P > \mu_Q$	B1		
	State assumption (A.E.F.):	Normal distns. for [ $P$ and] $Q$  <i>and</i> equal variances	B1		
Estimate (pooled) common variance:	$s^2 = (7 \times 1.003 + 11 \times 1.962) / 18$ $= 1.589$ or 1.261 <sup>2</sup>	M1 A1			
Calculate value of $t$ (to 3 s.f.):	$t = (5.35 - 4.60) / (s \sqrt{1/8 + 1/12})$ $= 1.30$	M1 A1			
Correct conclusion (A.E.F., $\sqrt{\dagger}$ on $t$ ):	$t < t_{18, 0.9} = 1.33$ so $Q$ 's mean is not less than $P$ 's	B1 $\sqrt{\dagger}$	7	<b>14</b>	