

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
GCE Advanced Level

**MARK SCHEME for the May/June 2012 question paper  
for the guidance of teachers**

**9231 FURTHER MATHEMATICS**

**9231/22**

Paper 2, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



<b>Page 2</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – May/June 2012</b>	<b>9231</b>	<b>22</b>

### Mark Scheme Notes

Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – May/June 2012</b>	<b>9231</b>	<b>22</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{\phantom{x}}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2012	9231	22

Question Number	Mark Scheme Details	Part Mark	Total	
1	<p><i>EITHER:</i> Use <math>C = I\alpha</math> to find angular acceleration <math>\alpha</math>: (omitting minus sign loses this A1 only) Integrate or use <math>\omega_1 = \omega_0 + \alpha t</math> to find time <math>t</math>:</p> <p><i>OR:</i> Use energy to find angle <math>\theta</math>: Use <math>\theta = \frac{1}{2}(\omega_0 + \omega_1)t</math> to find time <math>t</math>:</p>	$\alpha = -48 \times 0.3 / 18 [= -0.8]$ M1 A1  $t = (2 - 6)/\alpha = 5$ [s] M1 A1  $\theta = \frac{1}{2} 18(6^2 - 2^2)/(48 \times 0.3) [= 20]$ (M1 A1) $t = \theta / \frac{1}{2}(6 + 2) = 5$ [s] (M1 A1)	4	[4]
2	<p>Find two momentum eqns, e.g.:</p> <p>Solve to find both speeds after colln.:</p> <p>Find total loss in KE:</p>	$3mv_{3m} = 3m \times 2u - 4mu$ or $mv_m = -mu + 4mu$ or $3mv_{3m} + mv_m = 3m \times 2u - mu$ M1 M1 $v_{3m} = 2u/3$ and $v_m = 3u$ A1 $\frac{1}{2}m \{3(2u)^2 + u^2 - 3v_{3m}^2 - v_m^2\}$ M1 A1 $= \frac{1}{2}m (12 + 1 - 4/3 - 9) u^2$ $= (13/2 - 31/6 \text{ or } 16/3 - 4) mu^2$ $= (4/3) mu^2$ A.G. A1	6	[6]
3	<p>Use conservation of energy:</p> <p>Equate radial forces:</p> <p>Eliminate <math>v</math> to find <math>R</math>:</p> <p>(i) Find <math>\cos \theta_1</math> when <math>R = 0</math>: Find speed at this point:</p> <p>(iii) <i>EITHER:</i> Use energy to find reqd speed <math>v_2</math>: Simplify:</p> <p><i>OR:</i> Find horiz. comp. of <math>v_2</math>: Find vertical comp. of <math>v_2</math>: Combine comps. to find <math>v_2</math>:</p>	$\frac{1}{2}mv^2 = \frac{1}{2}m(7ga/2) - mga(1 - \cos \theta)$ B1 $[v^2 = 3ga/2 + 2ga \cos \theta]$ $R = mv^2/a + mg \cos \theta$ B1 $R = (3mg/2)(1 + 2 \cos \theta)$ A.G. M1 A1  $\cos \theta_1 = -\frac{1}{2}$ [ $\theta_1 = 2\pi/3$ ] B1 $v_1^2 = \frac{1}{2}ga$ , $v_1 = \sqrt{\frac{1}{2}ga}$ M1 A1  $\frac{1}{2}mv_2^2 = \frac{1}{2}mv_1^2 - mga \cos \theta_1$ M1 or $\frac{1}{2}m(7ga/2) - mga$ M1 $\frac{1}{2}v_2^2 = \frac{1}{4}ga + \frac{1}{2}ga$ , $v_2 = \sqrt{3ga/2}$ M1 A1  $v_1 \cos(\pi - \theta) [= \frac{1}{2}v_1 = \sqrt{ga/8}]$ (M1) $\sqrt{\{v_1^2 \sin^2(\pi - \theta) + ga\}} [= \sqrt{11ga/8}]$ (M1) $v_2 = \sqrt{ga/8 + 11ga/8} = \sqrt{3ga/2}$ (A1)	4 3 3	[10]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2012	9231	22

Question Number	Mark Scheme Details	Part Mark	Total
4	<p>Find MI of disc <math>D</math> about <math>O</math>:  <math>I_D = \frac{1}{2}(9m)(3a)^2 = (81/2)ma^2</math> B1</p> <p>Find MI of removed disc <math>D'</math> about <math>O</math>:  <math>I_{D'} = \frac{1}{2}ma^2</math> B1</p> <p>Find MI of circular lamina about <math>O</math>:  <math>I_O = I_D - I_{D'} [= 40ma^2]</math> M1</p> <p>Find MI of circular lamina about <math>A</math>:  <math>I_A = I_O + 8m(3a)^2</math> or <math>\frac{1}{2}(243 - 19)ma^2</math>  <math>= 112ma^2</math> <b>A.G.</b> M1 A1</p> <p>Find MI of lamina plus particle about <math>A</math>:  <math>I_A' = I_A + (3m)(6a)^2 = 220ma^2</math> B1</p> <p>State or imply that speed is max when <math>AB</math> vertical M1</p> <p>Use energy when <math>AB</math> vertical (or at general point):  <math>\frac{1}{2}I_A'\omega^2 = 8mg \times 6a + 3mg \times 12a</math>  or <math>2 \times 11mg \times 42a/11 [= 84mga]</math> M1 A1</p> <p>Substitute for <math>I_A'</math> and find max ang. speed <math>\omega</math> (or <math>\omega^2</math>):  <math>\omega^2 = 84mga/110ma^2 = 42g/55a</math> A1</p> <p>Equate <math>6a\omega</math> to <math>k\sqrt{ga}</math> to find <math>k</math>:  <math>k = \sqrt{(36 \times 42/55)} = 5.24</math> M1 A1</p>	5	[12]
5	<p>Resolve forces vertically:  <math>R_C + F_A = 3mg + 5mg</math> B1</p> <p>Take moments about <math>B</math> for <math>AB</math>:  <math>3aF_A = (3a/2)3mg</math> [<math>F_A = 3mg/2</math>] M1 A1</p> <p>Combine to find <math>R_C</math>:  <math>R_C = 8mg - 3mg/2 = 13mg/2</math> <b>A.G.</b> M1 A1</p> <p>Substitute for <math>R_C</math> in above resln. eqn to find <math>F_A</math>:  <math>F_A = 8mg - 13mg/2 = 3mg/2</math> B1</p> <p>Take moments about <math>B</math> for <math>BC</math>:  <math>4aF_C = 3aR_C - (3a/2)5mg</math> M1</p> <p>Substitute for <math>R_C</math> to find <math>F_C</math>:  <math>F_C = 3mg</math> A1</p> <p>Find limiting value <math>\mu_C</math> for <math>\mu</math> at <math>C</math> [or <math>A</math>] (A.E.F.):  <math>\mu_C = 6/13 [= 0.462</math> or <math>\mu_A = 0.5]</math> M1 A1</p> <p>Relate <math>R_A, F_C</math> by e.g. horizontal resolution:  <math>R_A = F_C [= 3mg]</math> B1</p> <p>Deduce least possible value of <math>\mu</math> for system:  <math>\mu_{min} = \max[\mu_A, \mu_C] = 0.5</math> B1</p>	5	[12]
6	<p>Find prob. that 10<sup>th</sup> bulb is first defective one:  <math>(1 - 0.01)^9 \times 0.01 = 0.00914</math>  (allow 0.00913) M1 A1</p> <p>State or find <math>E(N)</math>:  <math>E(N) = 1/0.01 = 100</math> B1</p> <p>Formulate condition for <math>n</math>:  (equality throughout loses this M1 only)  <math>P(N \leq n) = 1 - P(N &gt; n)</math>  <math>= 1 - 0.99^n &gt; 0.9, \quad 0.99^n &lt; 0.1</math> M1</p> <p>Take logs (any base) to give inequality for <math>n</math>:  <math>n &gt; \log 0.1 / \log 0.99</math> M1</p> <p>Find <math>n_{min}</math>:  <math>n &gt; 229.1, \quad n_{min} = 230</math> A1</p>	2 1 3	[6]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2012	9231	22

Question Number	Mark Scheme Details	Part Mark	Total	
7	<p>Consider differences e.g. outdoor – indoor times:            Calculate sample mean:            Estimate population variance:            (allow biased here: 2.679 or 1.637<sup>2</sup>)            State hypotheses (A.E.F.):            Calculate value of <math>t</math> (to 3 sf):            State or use correct tabular <math>t</math> value:            (or can compare <math>\bar{d}</math> with 1.46[3])            Correct conclusion (AEF, dep *A1, *B1):</p>	<p>0.1 2.1 -0.1 0.2 2.4 0.5 2.8 -2.6 M1  <math>\bar{d} = 5.4 / 8 = 0.675</math> M1  <math>s^2 = (25.08 - 5.4^2/8) / 7</math>  <math>= 3.062</math> or <math>1.750^2</math> M1  <math>H_0: \mu_o - \mu_i = 0, H_1: \mu_o - \mu_i \neq 0</math> B1  <math>t = \bar{d}/(s/\sqrt{8}) = 1.09</math> M1 *A1  <math>t_{7,0.975} = 2.36[5]</math> *B1</p>	8	[8]
8	<p>Relate <math>F(x)</math> to Poisson distribution (ignore <math>x &lt; 0</math>):            Equate <math>F(x)</math> to <math>1 - e^{-\lambda x}</math> or <math>f(x)</math> to <math>e^{-\lambda x}</math> to find mean <math>\lambda</math>:            (i) Formulate eqn for median <math>m</math> of <math>X</math>:            Find value of <math>m</math>:            (ii) Find <math>P(X \geq 50)</math> (or <math>&gt; 50</math>):</p>	<p><math>F(x) = 1 - P(X &gt; x)</math>  <math>= 1 - P(\text{no flaws in length } x)</math> M1  <math>= 1 - e^{-(x/100)1.6} = 1 - e^{-0.016x}</math> A.G. M1 A1  <math>\lambda = 1/0.016 = 62.5</math> B1</p> <p><math>1 - e^{-0.016m} = 1/2</math> M1  <math>m = -\ln 1/2 / 0.016 = 43.3</math> M1 A1</p> <p><math>1 - F(50) = e^{-0.8} = 0.449</math> M1 A1</p>	4	[9]
9	<p>Calculate sample mean:            Estimate population variance:            (allow biased here: 1.940 or 1.393<sup>2</sup>)            State hypotheses (A.E.F.):            Calculate value of <math>t</math> (to 3 sf):            State or use correct tabular <math>t</math> value:            (or can compare <math>\bar{d}</math> with <math>4.5 + 0.998 = 5.49[8]</math>)            Correct conclusion (AEF, dep *A1, *B1):            Find confidence interval (allow <math>z</math> in place of <math>t</math>) e.g.:            Use of correct tabular value:            Evaluate C.I. correct to 3 s.f.:</p>	<p><math>\bar{d} = 42.5 / 8 = 5.3125</math> M1  <math>s^2 = 15.519 / 7</math>  <math>= 2.217</math> or <math>1.489^2</math> M1  <math>H_0: \mu = 4.5, H_1: \mu &gt; 4.5</math> B1  <math>t = (\bar{d} - 4.5)/(s/\sqrt{8}) = 1.54</math> M1 *A1  <math>t_{7,0.95} = 1.89[5]</math> *B1</p> <p>Mean is not greater than 4.5 B1</p> <p><math>5.3125 \pm t \sqrt{\{15.519/(7 \times 8)\}}</math> M1  <math>t_{7,0.975} = 2.36[5]</math> A1  <math>5.31 \pm 1.24[5]</math> or <math>[4.07, 6.56]</math> A1</p>	7	[10]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2012	9231	22

Question Number	Mark Scheme Details	Part Mark	Total		
10	Find expected values (to 1 d.p.): (lose A1 if rounded to integers) State (at least) null hypothesis (A.E.F.): Calculate value of $\chi^2$ :  State or use correct tabular $\chi^2$ value (to 3 sf): Conclusion consistent with values (A.E.F): Calculate new value $\chi_{\text{new}}^2$ of $\chi^2$ : State or use correct tabular $\chi^2$ value: Find $N_{\text{min}}$ :	$54.72 \quad 52.992 \quad 36.288$ $40.28 \quad 39.008 \quad 26.712$ $H_0$ : Preferences are independent $\chi^2 = 0.5095 + 0.0186 + 0.5067$ $+ 0.6921 + 0.0252 + 0.6883$ $= 2.44 \text{ or } 2.45$  $\chi_{2, 0.95}^2 = 5.99[1]$ Preferences are independent $\chi_{\text{new}}^2 = N \times \chi^2$ $\chi_{2, 0.99}^2 = 9.21$ $N > 9.21/2.45, N_{\text{min}} = 4$	M1 A1 B1  M1 A1 B1 A1 ✓ M1 B1 M1 A1	7       4	[11]
11 (a)	Resolve vertically at equilibrium with extn. $e$ : Use Newton's Law at general point:  Simplify to give standard SHM eqn: S.R.: Stating this without derivation (max 3/4): Find period $T$ using SHM with $\omega = \sqrt{4g/l}$ : Find speed $v_E$ at $E$ using $v^2 = \omega^2 (A^2 - x^2)$ with $x = 0$ : Find speed $v_P$ before striking plane (A.E.F.): Find comps. of speed $V$ after striking plane:  Combine to find $V$ :	$4mge/l = mg \quad [e = 1/4l]$ $m \, d^2x/dt^2 = mg - 4mg(e+x)/l$ [ or $-mg + 4mg(e-x)/l$ ] $d^2x/dt^2 = -(4g/l)x$  $T = 2\pi/\sqrt{4g/l} = \pi\sqrt{l/g}$ <b>A.G.</b>  $v_E = \omega l/8 = 1/4\sqrt{gl}$ $v_P = \sqrt{(gl/16 + 14gl/16)} = 1/4\sqrt{15gl}$ Parallel to plane: $v_P \sin 30^\circ$ or $1/2 v_P$ or $\sqrt{15gl/64}$ Normal to plane: $1/3 v_P \cos 30^\circ$ or $1/3(\sqrt{3}/2) v_P$ or $\sqrt{5gl/64}$ $V^2 = 15gl/64 + 5gl/64 = 5gl/16$ $V = 1/4\sqrt{5gl}$ <b>A.G.</b>	B1 M1 A1  (B1) B1 M1 A1 M1 A1  B1 B1 M1 A1	4	[12]

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2012	9231	22

Question Number	Mark Scheme Details	Part Mark	Total	
<b>(b)</b>	<b>(i)</b> Find correlation coefficient $r$ :  (A0 if only 3 s.f. used)	$\Sigma x = 8.5, \Sigma x^2 = 9.67, \Sigma xy = 3.955,$ $\Sigma y = 3.85, \Sigma y^2 = 2.0775$ $r = (3.955 - 8.5 \times 3.85/8) / \sqrt{\{(9.67 - 8.5^2/8)(2.0775 - 3.85^2/8)\}}$ M1 A1 $= -0.1356 / \sqrt{(0.6387 \times 0.2247)}$ A1 $= -0.1356 / (0.7992 \times 0.4740)$ [or $-0.01695 / \sqrt{(0.07984 \times 0.02809)}$ $= -0.01695 / (0.2826 \times 0.1676)$ ] $= -0.358$ *A1	4	
	<b>(ii)</b> State both hypotheses: State or use correct tabular one-tail $r$ value: Valid method for reaching conclusion: Correct conclusion (AEF, dep *A1, *B1): Valid comment, consistent with values	$H_0: \rho = 0, H_1: \rho < 0$ B1 $r_{8, 2.5\%} = 0.707$ *B1 Accept $H_0$ if $ r  <$ tabular value M1 There is no negative correlation A1 No effect of $S$ on $R$ (A.E.F.) B1√	5	
	<b>(iii)</b> Find critical tabular one-tail $r$ value: Deduce range of possible values of $N$ :	$r_{16, 5\%} = 0.426$ or $r_{15, 5\%} = 0.441$ M1 A1 $N \geq 16$ A1	3	<b>[12]</b>