



Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/42

Paper 4 Mechanics

May/June 2024

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.



- 1 A cyclist and bicycle have a total mass of 72 kg. The cyclist rides along a horizontal road against a total resistance force of 28 N.

Find the total work done by the cyclist to increase his speed from 8 ms^{-1} to 16 ms^{-1} while travelling a distance of 100 metres. [3]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

2 A particle P moves in a straight line. At time t s after leaving a point O on the line, P has velocity v ms^{-1} , where $v = 44t - 6t^2 - 36$.

(a) Find the set of values of t for which the acceleration of the particle is positive. [2]

[illegible]

(b) Find the two values of t at which P returns to O . [3]

[illegible]

- 4 A car has mass 1400 kg . When the speed of the car is $v\text{ ms}^{-1}$ the magnitude of the resistance to motion is $kv^2\text{ N}$ where k is a constant.

(a) The car moves at a constant speed of 24 ms^{-1} up a hill inclined at an angle of α to the horizontal where $\sin \alpha = 0.12$. At this speed the magnitude of the resistance to motion is 480 N .

(i) Find the value of k . [1]

.....

.....

.....

.....

(ii) Find the power of the car's engine. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The car now moves at a constant speed on a straight level road.

Given that its engine is working at 54 kW , find this speed. [3]

.....

.....

.....

.....

.....

.....

.....

.....

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 6 Three particles A , B and C of masses 5 kg , 1 kg and 2 kg respectively lie at rest in that order on a straight smooth horizontal track XYZ . Initially A is at X , B is at Y and C is at Z . Particle A is projected towards B with a speed of 6 ms^{-1} and at the same instant C is projected towards B with a speed of $v\text{ ms}^{-1}$. In the subsequent motion, A collides and coalesces with B to form particle D . Particle D then collides and coalesces with C to form particle E and E moves towards Z .

(a) Show that after the second collision the speed of E is $\frac{15-v}{4} \text{ ms}^{-1}$. [3]

This image shows a full page of primary-ruled paper. It contains ten identical horizontal rows. Each row is defined by three lines: a solid top line, a dashed middle line, and a solid bottom line. The rows are evenly spaced across the entire page, providing a template for handwriting practice.

(b) The total loss of kinetic energy of the system due to the two collisions is 63 J.

Use the result from **(a)** to show that $v = 3$. [3]

This image shows a single sheet of white paper with ten evenly spaced horizontal dotted lines. The lines are black and extend across the full width of the page, providing a guide for handwriting practice. There is no text or other markings on the paper.

.....

.....

.....

.....

.....

.....

.....

.....

(c) It is given that the distance XY is 36 m and the distance YZ is 98 m.

(i) Find the time between the two collisions. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the time between the instant that A is projected from X and the instant that E reaches Z . [1]


.....

.....

.....

.....

.....



Two particles P and Q of masses 2.5 kg and 0.5 kg respectively are connected by a light inextensible string that passes over a small smooth pulley fixed at the top of a plane inclined at an angle of 30° to the horizontal. Particle P is on the plane and Q hangs below the pulley such that the level of Q is 2 m below the level of P (see diagram).

Particle P is released from rest with the string taut and slides down the plane. The plane is rough with coefficient of friction 0.2 between the plane and P .

- (a) Find the acceleration of P . [5]

[illegible]

- (b)** Use an energy method to find the speed of the particles at the instant when they are at the same vertical height. [5]

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

