



## Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**MATHEMATICS**

**9709/43**

Paper 4 Mechanics

**May/June 2022**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.



- 1 Two particles  $P$  and  $Q$ , of masses  $0.3\text{ kg}$  and  $0.2\text{ kg}$  respectively, are at rest on a smooth horizontal plane.  $P$  is projected at a speed of  $4\text{ m s}^{-1}$  directly towards  $Q$ . After  $P$  and  $Q$  collide,  $Q$  begins to move with a speed of  $3\text{ m s}^{-1}$ .

(a) Find the speed of  $P$  after the collision. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

After the collision,  $Q$  moves directly towards a third particle  $R$ , of mass  $m\text{ kg}$ , which is at rest on the plane. The two particles  $Q$  and  $R$  coalesce on impact and move with a speed of  $2\text{ m s}^{-1}$ .

(b) Find  $m$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 3

2 A particle  $P$  is projected vertically upwards from horizontal ground.  $P$  reaches a maximum height of 45 m. After reaching the ground,  $P$  comes to rest without rebounding.

(a) Find the speed at which  $P$  was projected. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the total time for which the speed of  $P$  is at least  $10 \text{ ms}^{-1}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(b) Find the acceleration of the particle between  $t = 0$  and  $t = 5$ , given that it is constant. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the average speed of the particle during its motion. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....









The cyclist comes to the top of a hill inclined at  $5^\circ$  to the horizontal. The cyclist stops pedalling and freewheels down the hill (so that the cyclist is no longer supplying any power). The magnitude of the resistance force remains at  $30\text{ N}$ . Over a distance of  $d\text{ m}$ , the speed of the cyclist increases from  $6\text{ m s}^{-1}$  to  $12\text{ m s}^{-1}$ .

- (b) Find the change in kinetic energy. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Use an energy method to find  $d$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

.....

.....

.....

- (b) It is given instead that the plane  $BC$  is rough. A force of magnitude  $3\text{ N}$  is applied to  $Q$  directly up the plane along a line of greatest slope of the plane.

Find the least value of the coefficient of friction between  $Q$  and the plane  $BC$  for which the particles remain at rest. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....









**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.