



## Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**MATHEMATICS**

**9709/42**

Paper 4 Mechanics

**February/March 2020**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.



## 3

- 2 A particle  $P$  of mass  $0.4 \text{ kg}$  is on a rough horizontal floor. The coefficient of friction between  $P$  and the floor is  $\mu$ . A force of magnitude  $3 \text{ N}$  is applied to  $P$  upwards at an angle  $\alpha$  above the horizontal, where  $\tan \alpha = \frac{3}{4}$ . The particle is initially at rest and accelerates at  $2 \text{ m s}^{-2}$ .

- (a) Find the time it takes for  $P$  to travel a distance of  $1.44 \text{ m}$  from its starting point. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find  $\mu$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....















- 6 On a straight horizontal test track, driverless vehicles (with no passengers) are being tested. A car of mass 1600 kg is towing a trailer of mass 700 kg along the track. The brakes are applied, resulting in a deceleration of  $12 \text{ m s}^{-2}$ . The braking force acts on the car only. In addition to the braking force there are constant resistance forces of 600 N on the car and of 200 N on the trailer.

(a) Find the magnitude of the force in the tow-bar. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the braking force. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 11

- (c) At the instant when the brakes are applied, the car has speed  $22 \text{ m s}^{-1}$ . At this instant the car is  $17.5 \text{ m}$  away from a stationary van, which is directly in front of the car.

Show that the car hits the van at a speed of  $8 \text{ m s}^{-1}$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) After the collision, the van starts to move with speed  $5 \text{ m s}^{-1}$  and the car and trailer continue moving in the same direction with speed  $2 \text{ m s}^{-1}$ .

Find the mass of the van. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 7 A particle moves in a straight line through the point  $O$ . The displacement of the particle from  $O$  at time  $t$  s is  $s$  m, where

$$s = t^2 - 3t + 2 \quad \text{for } 0 \leq t \leq 6,$$

$$s = \frac{24}{t} - \frac{t^2}{4} + 25 \quad \text{for } t \geq 6.$$

- (a) Find the value of  $t$  when the particle is instantaneously at rest during the first 6 seconds of its motion. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

At  $t = 6$ , the particle hits a barrier at a point  $P$  and rebounds.

- (b) Find the velocity with which the particle arrives at  $P$  and also the velocity with which the particle leaves  $P$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....







**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.