



Cambridge International AS & A Level

MATHEMATICS**9709/51**

Paper 5 Probability & Statistics 1

May/June 2020

MARK SCHEME

Maximum Mark: 50

Published

Students did not sit exam papers in the June 2020 series due to the Covid-19 global pandemic.

This mark scheme is published to support teachers and students and should be read together with the question paper. It shows the requirements of the exam. The answer column of the mark scheme shows the proposed basis on which Examiners would award marks for this exam. Where appropriate, this column also provides the most likely acceptable alternative responses expected from students. Examiners usually review the mark scheme after they have seen student responses and update the mark scheme if appropriate. In the June series, Examiners were unable to consider the acceptability of alternative responses, as there were no student responses to consider.

Mark schemes should usually be read together with the Principal Examiner Report for Teachers. However, because students did not sit exam papers, there is no Principal Examiner Report for Teachers for the June 2020 series.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the June 2020 series for most Cambridge IGCSE™ and Cambridge International A & AS Level components, and some Cambridge O Level components.

This document consists of **13** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

Abbreviations


AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

Question	Answer	Marks
1(a)	Prob of 4 (from 1,3, 3,1 or 2,2) = $\frac{3}{36} = \frac{1}{12}$ AG	B1
		1
1(b)	Mean = $\frac{1}{\frac{1}{12}} = 12$	B1
		1
1(c)	$\left(\frac{11}{12}\right)^5 \times \frac{1}{12} = 0.0539$ or $\frac{161051}{2985984}$	B1
		1
1(d)	$1 - \left(\frac{11}{12}\right)^7$	M1
	0.456 or $\frac{16344637}{35831808}$	A1
		2

Question	Answer	Marks	
2(a)	6!	M1	
	720	A1	
		2	
2(b)	Total number: $\frac{9!}{3!2!}(30240)$	M1	
	Number with Ls together = $\frac{8!}{3!}(6720)$	M1	
	Number with Ls not together = $\frac{9!}{3!2!} - \frac{8!}{3!}$ = 30 240 – 6720	M1	
	23 520	A1	
	Alternative method for question 2(b)		
	$\frac{7!}{3!} \times \frac{8 \times 7}{2}$		
	$7! \times k$ in numerator, k integer ≥ 1	M1	
	$8 \times 7 \times m$ in numerator or $8C2 \times m$, m integer ≥ 1	M1	
	3! in denominator	M1	
	23 520	A1	
	4		

Question	Answer	Marks										
3(a)	<table border="1" data-bbox="365 215 1209 391"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>Probability</td> <td>$\frac{1}{56}$</td> <td>$\frac{15}{56}$</td> <td>$\frac{30}{56}$</td> <td>$\frac{10}{56}$</td> </tr> </table> <p data-bbox="365 427 1142 459">(B1 for probability distribution table with correct outcome values)</p> $P(0) = \frac{3}{8} \times \frac{2}{7} \times \frac{1}{6} = \frac{1}{56}$ $P(1) = \frac{5}{8} \times \frac{3}{7} \times \frac{2}{6} \times 3 = \frac{15}{56}$ $P(2) = \frac{5}{8} \times \frac{4}{7} \times \frac{3}{6} \times 3 = \frac{30}{56}$ $P(3) = \frac{5}{8} \times \frac{4}{7} \times \frac{3}{6} = \frac{10}{56}$ <p data-bbox="365 805 712 837">(M1 for denominator $8 \times 7 \times 6$)</p> <p data-bbox="365 874 967 906">Any one probability correct (with correct outcome)</p> <p data-bbox="365 938 649 970">All probabilities correct</p>	x	0	1	2	3	Probability	$\frac{1}{56}$	$\frac{15}{56}$	$\frac{30}{56}$	$\frac{10}{56}$	<p data-bbox="2011 215 2060 247">B1</p> <p data-bbox="2011 491 2060 523">M1</p> <p data-bbox="2011 874 2060 906">A1</p> <p data-bbox="2011 938 2060 970">A1</p> <p data-bbox="2033 1002 2060 1034">4</p>
x	0	1	2	3								
Probability	$\frac{1}{56}$	$\frac{15}{56}$	$\frac{30}{56}$	$\frac{10}{56}$								
3(b)	$1 - P(8, 9, 10) = 1 - \left[{}^{10}C_8 0.64^8 0.36^2 + {}^{10}C_9 0.64^9 0.36^1 + 0.64^{10} \right]$ $1 - (0.164156 + 0.064852 + 0.11529)$ <p data-bbox="365 1225 436 1257">0.759</p>	<p data-bbox="2011 1066 2060 1098">M1</p> <p data-bbox="2011 1161 2060 1193">M1</p> <p data-bbox="2011 1225 2060 1257">A1</p> <p data-bbox="2033 1289 2060 1321">3</p>										

Question	Answer	Marks
4	Scenarios: 2P 3V 2G ${}^8C_2 \times {}^4C_2 \times {}^6C_3 = 28 \times 6 \times 20 = 3360$ 2P 4V 1G ${}^8C_2 \times {}^4C_1 \times {}^6C_4 = 28 \times 4 \times 15 = 1680$ 3P 3V 1G ${}^8C_3 \times {}^4C_1 \times {}^6C_3 = 56 \times 4 \times 20 = 4480$ 4P 2V 1G ${}^8C_4 \times {}^4C_1 \times {}^6C_2 = 70 \times 4 \times 15 = 4200$ (M1 for ${}^8C_r \times {}^4C_r \times {}^6C_r$ with $\sum r = 7$)	M1
	Two unsimplified products correct	B1
	Summing the number of ways for 3 or 4 correct scenarios	M1
	Total: 13 720	A1
		4

Question	Answer	Marks
5(a)	 <p>Fully correct labelled tree for method of transport with correct probabilities.</p>	<p>B1</p> <hr/> <p>Fully correct labelled branches with correct probabilities for lateness with either 1 branch after W or 2 branches with the prob 0</p> <p>B1</p> <hr/> <p>2</p>
5(b)	<p>$0.35 \times 0.3 + 0.44 \times 0.8 (+ 0)$</p> <hr/> <p>0.457</p>	<p>M1</p> <hr/> <p>A1</p> <hr/> <p>2</p>

Question	Answer	Marks
5(c)	$P(\text{not B} \text{not fruit}) = \frac{P(B' \cap F')}{P(F')}$	M1
	$\frac{0.35 \times 0.7 + 0.21 \times 1}{1 - \text{their}(\mathbf{b})}$	M1
	$\frac{0.455}{0.543}$ (M1 for 1 – <i>their</i> (b) or summing three appropriate 2-factor probabilities, correct or consistent with <i>their</i> tree diagram as denominator)	M1
	0.838 or $\frac{455}{543}$	A1
		4

Question	Answer	Marks
6(a)	$P\left(\frac{50 - 54}{6.1} < z < \frac{60 - 54}{6.1}\right) = P(-0.6557 < Z < 0.9836)$	M1
	Both values correct	A1
	$\Phi(0.9836) - \Phi(-0.6557) = \Phi(0.9836) + \Phi(0.6557) - 1$ $= 0.8375 + 0.7441 - 1$ (Correct area)	M1
	0.582	A1
		4

Question	Answer	Marks
6(b)	$\frac{45 - \mu}{\sigma} = -0.994$	B1
	$\frac{56 - \mu}{\sigma} = 1.372$	B1
	One appropriate standardisation equation with μ, σ, z -value (not probability) and 45 or 56.	M1
	$11 = 2.366 \sigma$ (M1 for correct algebraic elimination of μ or σ from <i>their</i> two simultaneous equations to form an equation in one variable)	M1
	$\sigma = 4.65, \mu = 49.6$	A1
		5

Question	Answer	Marks
7(a)	Class widths: 10, 5, 15, 20, 10	M1
	Frequency density = frequency/ <i>their</i> class width: 1.8, 4.8, 2, 1, 0.8	M1
	All heights correct on diagram (using a linear scale)	A1
	Correct bar ends	B1
	Bar ends: 10.5, 15.5, 30.5, 50.5, 60.5	B1
		5
7(b)	11 – 15 and 31 – 50	B1
	Greatest IQR = 50 – 11 = 39	B1
		2
7(c)	Mean = $\frac{18 \times 5.5 + 24 \times 13 + 30 \times 23 + 20 \times 40.5 + 8 \times 55.5}{100} = \frac{2355}{100} = 23.6$	B1
	Var = $\frac{18 \times 5.5^2 + 24 \times 13^2 + 30 \times 23^2 + 20 \times 40.5^2 + 8 \times 55.5^2}{100} - \text{mean}^2$	M1
	$\frac{77917.5}{100} - \text{mean}^2 = 224.57$	A1
	Standard deviation = 15.0 (FT <i>their</i> variance)	A1 FT
		4