



## Cambridge International AS & A Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 7 9 7 7 4 4 6 4 4 5 2 9 \*

**MATHEMATICS**

**9709/62**

Paper 6 Probability & Statistics 2

**May/June 2021**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **12** pages. Any blank pages are indicated.

**BLANK PAGE**



2 The random variable  $X$  has the distribution  $B(400, 0.01)$ .

(a) Find  $\text{Var}(4X + 2)$ . [3]

.....

.....

.....

.....

.....

(b) (i) State an appropriate approximating distribution for  $X$ , giving the values of any parameters. Justify your choice of approximating distribution. [2]

.....

.....

.....

.....

.....

.....

.....

(ii) Use your approximating distribution to find  $P(2 \leq X \leq 5)$ . [2]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

3



The random variable  $X$  takes values in the range  $1 \leq x \leq p$ , where  $p$  is a constant. The graph of the probability density function of  $X$  is shown in the diagram.

(a) Show that  $p = 2$ . [2]

.....

.....

.....

.....

(b) Find  $E(X)$ . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 6

- 4 Wendy's journey to work consists of three parts: walking to the train station, riding on the train and then walking to the office. The times, in minutes, for the three parts of her journey are independent and have the distributions  $N(15.0, 1.1^2)$ ,  $N(32.0, 3.5^2)$  and  $N(8.6, 1.2^2)$  respectively.

- (a) Find the mean and variance of the total time for Wendy's journey. [2]

.....

.....

.....

If Wendy's journey takes more than 60 minutes, she is late for work.

- (b) Find the probability that, on a randomly chosen day, Wendy will be late for work. [3]

.....

.....

.....

.....

.....

.....

.....

- (c) Find the probability that the mean of Wendy's journey times over 15 randomly chosen days will be less than 54.5 minutes. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....







(b) Calculate a 99% confidence interval for the mean height of animals of this species. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Four random samples were taken and a 99% confidence interval for the population mean,  $\mu$ , was found from each sample.

(c) Find the probability that all four of these confidence intervals contain the true value of  $\mu$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 Customers arrive at a particular shop at random times. It has been found that the mean number of customers who arrive during a 5-minute interval is 2.1.

(a) Find the probability that exactly 4 customers arrive during a 10-minute interval. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the probability that at least 4 customers arrive during a 20-minute interval. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Use a suitable approximating distribution to find the probability that fewer than 40 customers arrive during a 2-hour interval. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

