

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Subsidiary Level

## **MARK SCHEME for the May/June 2015 series**

### **9709 MATHEMATICS**

**9709/22**

Paper 2 maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS Level – May/June 2015	9709	22

### Mark Scheme Notes

Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\nabla$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS Level – May/June 2015</b>	<b>9709</b>	<b>22</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only - often written by a “fortuitous” answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through” marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS Level – May/June 2015	9709	22
1	(i) Introduce logarithms and use power law Obtain $x = 21.6$	M1 A1	[2]
	(ii) Obtain or imply $-21.6$ or $-21$ as lower value State 43	B1 B1	[2]
2	(i) Substitute $x = -2$ into expression and equate to zero Obtain $-32 + 4a + 2(a + 1) - 18 = 0$ or equivalent Obtain $a = 8$	M1 A1 A1	[3]
	(ii) Attempt to find quadratic factor by division, inspection, ... Obtain $4x^2 - 9$ State $(x + 2)(2x - 3)(2x + 3)$	M1 A1 A1	[3]
3	(i) Use identity $\sec^2 \theta = 1 + \tan^2 \theta$ Solve three-term quadratic equation in $\tan \theta$ Obtain at least $\tan \theta = \frac{5}{2}$	B1 M1 A1	[3]
	(ii) Substitute numerical values into $\tan(A + B)$ identity Obtain $\frac{\frac{5}{2} + (-1)}{1 - \frac{5}{2}(-1)}$ or equivalent, following their positive answer from part (i) Obtain $\frac{3}{7}$ or exact equivalent and no other answers	M1 A1 <sup>ft</sup> A1	[3]
4	(i) Differentiate to obtain $e^x - 8e^{-2x}$ Use correct process to solve equation of form $ae^x + be^{-2x} = 0$ Confirm given answer $\ln 2$ correctly	B1 M1 A1	[3]
	(ii) Integrate to obtain expression of form $pe^x + qe^{-2x}$ Obtain correct $e^x - 2e^{-2x}$ Apply both limits correctly Confirm given answer $\frac{5}{2}$	M1 A1 M1 depM A1	[4]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS Level – May/June 2015	9709	22

- 5 (i) Draw recognisable sketch of  $y = 16 - x^4$  B1  
 Draw recognisable sketch of  $y = |3x|$  B1  
 Indicate in some way the two points of intersection B1 depBB  
 [3]
- (ii) Use iterative process correctly at least once M1  
 Obtain final answer 1.804 A1  
 Show sufficient iterations to justify answer or show sign change in the interval (1.8035, 1.8045) A1 [3]
- (iii) State (1.804, 5.412) B1  
 State  $(-1.804, 5.412)$ , following their first point B1<sup>dep</sup> [2]
- 6 (i) Solve three-term quadratic equation for  $\sin x$  M1  
 Obtain at least  $\sin x = -\frac{1}{2}$  and no errors seen A1  
 Obtain  $x = \frac{7}{6}\pi$  A1 [3]
- (ii) State  $\sin^2 x = \frac{1}{2} - \frac{1}{2}\cos 2x$  B1  
 Obtain given  $5 + 8\sin x - 2\cos 2x$  with necessary detail seen B1  
 Integrate to obtain expression of form  $ax + b\cos x + c\sin 2x$  M1  
 Obtain correct  $5x - 8\cos x - \sin 2x$  A1  
 Apply limits 0 and their  $x$ -value correctly M1 depM  
 Obtain  $\frac{35}{6}\pi + \frac{7}{2}\sqrt{3} + 8$  or exact equivalent A1 [6]
- 7 (a) Differentiate  $4\ln y$  to obtain  $\frac{4}{y} \times \frac{dy}{dx}$  B1  
 Differentiate  $6xy$  to obtain  $6y + 6x\frac{dy}{dx}$  B1  
 Substitute 1 and 1 and solve for  $\frac{dy}{dx}$  M1  
 Obtain  $-\frac{9}{10}$  or equivalent A1 [4]
- (b) Obtain  $\frac{dx}{dt} = -10t^{-2} - 1$  B1  
 Obtain derivative of form  $k(2t-1)^{-\frac{1}{2}}$  for  $\frac{dy}{dt}$  M1  
 Obtain correct  $(2t-1)^{-\frac{1}{2}}$  A1  
 Identify value of  $t$  as 5 B1  
 Obtain expression for  $\frac{dy}{dx}$  correctly, with numerical value of  $t$  substituted M1  
 Obtain  $-\frac{5}{21}$  or exact equivalent A1 [6]