UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2010 question paper

for the guidance of teachers

9709 MATHEMATICS

9709/22

Paper 2, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2010	9709	22

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0. B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking *g* equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2010	9709	22

The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR -1 A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR-2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA -1 This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

	Page 4	Mark Scheme: Teachers' version Syllabus	Paper	
		GCE A/AS LEVEL – October/November 2010 9709	22	
1	EITHER: S	tate or imply non-modular inequality $(x+1)^2 > (x-4)^2$, or corresponding		
		quation or pair of linear equations	M1	
	С	Obtain critical value $\frac{3}{2}$	A1	
		2		
	S	tate correct answer $x > \frac{3}{2}$	A1	
	OR: S	tate a correct linear equation for the critical value, e.g. $x + 1 = -x + 4$, or		
		orresponding correct linear inequality, e.g. $x + 1 > -(x - 4)$	M1	
	С	Obtain critical value $\frac{3}{2}$	A1	
	S	tate correct answer $x > \frac{3}{2}$	A1	[3]
	-	2		[-]
2	Use law for t	he logarithm of a product, a quotient or a power	M1*	
		$5 = (2x+1)\log 2$, or equivalent	A1	
		via correct manipulative technique(s)	M1(dep*)	E 4 3
	Obtain answe	er $x = 3.11$. Allow $x \in [3.10, 3.11]$	A1	[4]
3	Integrate and	obtain $\frac{1}{2}e^{2x}$ term	B1	
	Obtain $2e^x$ t	2	B1	
	Obtain <i>x</i>		B1	
	Use limits co Obtain given	prrectly, allow use of limits $x = 1$ and $x = 0$ into an incorrect form	M1 A1	[5]
	•	g limits into original <u>integrand</u> , 0/5		[2]
4	(i) Stata da	$\frac{dy}{dt} = \frac{1}{t-2}$ or $\frac{dy}{dt} = 1 - 9t^{-2}$	B1	
+			DI	
	Use $\frac{dy}{dr}$	$=\frac{dy}{dt} \div \frac{dx}{dt}$	M1	
	ил	given answer correctly	A1	[3]
	(ii) Equate of	derivative to zero and solve for t	M1	
		imply that $t = 3$ is admissible c.w.o., and note $t = -3$, 2 cases	A1	[0]
	Obtain c	coordinates (1, 6) and no others	A1	[3]
5		rig identity to obtain a quadratic in $\cot \theta$ or $\tan \theta$	M1	
	1	adratic correctly	A1	
	Obtain $\tan \theta$	$=\frac{1}{2}$ or $-\frac{2}{3}$	A1√	
		er 26.6° or 146.3°	A1	
	•	rrect method for second answer from either root	M1	۲ <i>८</i> ٦
		ning 3 answers from 26.6°, 146.3°, 206.6°, 326.3° and no others in the range rers outside the given range]	e Al	[6]
	L 0 wild fr			

	Page 5		ge 5 Mark Scheme: Teachers' version Syllabus		Paper	
			GCE A/AS LEVEL – October/November 2010	9709	22	
6	(i)	Consider	sign of $\frac{6}{x^2} - x - 1$ at $x = 1.4$ and $x = 1.6$, or equivalent		M1	
		Complete	the argument correctly with appropriate calculations		A1	[2]
	(ii)	State $\frac{6}{r^2}$	= x + 1		B1	
		\mathcal{A}	e equation to given equation or vice versa		B1	[2]
	(iii)	Obtain fir	erative formula correctly at least once hal answer 1.54	n in a sinn abanan i	M1 A1	
			ficient iterations to justify its accuracy to 2 d.p. or show ther al (1.535, 1.545)	e is a sign change i	n B1	[3]
7 (i)	(i)		x = 1, equate to zero and obtain a correct equation in any for $x = 2$ and equate to 10	orm	B1 M1	
			correct equation in any form		A1	
		Solve a re	levant pair of equations for a or for b		M1	
		Obtain <i>a</i> =	= -17 and $b = 12$		A1	[5]
	(ii)	At any sta	ge, state that $x = 1$ is a solution		B1	
	()	EITHER	-	$3x^2 + 5x$	M1	
			Obtain quotient $3x^2 + 5x - 12$		A1	
			_			
			Obtain solutions $x = -3$ and $x = \frac{4}{3}$		A1	
		OR:	Obtain solution $x = -3$ by trial and error or inspection		B1	
			Obtain solution $x = \frac{4}{3}$		B2	
		[If an attempt at the quadratic factor is made by inspection, the M1 is earned if it reaches				
		an unknov	wn factor of $3x^2 + 5x + \lambda$ and an equation in λ]			[4]
8	(i)	Use produ			M1	
			rrect derivative in any form		A1	
		Substitute	$x = \frac{1}{2}\pi$, and obtain gradient of -1 for normal		A1√	
				$y' = \sin x - x \cos x$	ONLY	
		Show that	t line through $\left(\frac{1}{2}\pi, \frac{1}{2}\pi\right)$ with gradient –1 passes through (a	$(\pi, 0)$	M1	
					A1	[5]
	(ii)	Differenti	ate sin x and use product rule to differentiate $x \cos x$		M1	
		Obtain xs	$\sin x$, or equivalent		A1	[2]
	(iii)		integral is $\sin x - x \cos x (+ c)$		B1	
		Substitute	e limits 0 and $\frac{\pi}{2}$ correctly		M1	
		Obtain an S. R. Feed	swer 1 ling limits into original <u>integrand</u> , 0/3		A1	[3]