

**UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
GCE Advanced Subsidiary Level and GCE Advanced Level

**MARK SCHEME for the May/June 2011 question paper**  
**for the guidance of teachers**

**9709 MATHEMATICS**

**9709/32**

Paper 3, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

### Mark Scheme Notes

Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE AS/A LEVEL – May/June 2011</b>	<b>9709</b>	<b>32</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\sqrt{}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

- 1 *EITHER:* State or imply non-modular inequality  $x^2 < (5 + 2x)^2$ , or corresponding equation, or pair of linear equations  $x = \pm(5 + 2x)$  M1
- Obtain critical values  $-5$  and  $-\frac{5}{3}$  only A1
- Obtain final answer  $x < -5, x > -\frac{5}{3}$  A1
- OR:* State one critical value e.g.  $-5$ , by solving a linear equation or inequality, or from a graphical method, or by inspection B1
- State the other critical value, e.g.  $-\frac{5}{3}$ , and no other B1
- Obtain final answer  $x < -5, x > -\frac{5}{3}$  B1 [3]
- [Do not condone  $\leq$  or  $\geq$ .]
- 2 (i) Use law for the logarithm of a product or quotient M1  
Use  $\log_2 32 = 5$  or  $2^5 = 32$  M1  
Obtain  $x^2 + 5x - 32 = 0$ , or horizontal equivalent A1 [3]
- (ii) Solve a 3-term quadratic equation M1  
Obtain answer  $x = 3.68$  only, or exact equivalent, e.g.  $\frac{\sqrt{153} - 5}{2}$  A1 [2]
- 3 Use correct trig formula (or formulae) and obtain an equation in  $\cos\theta$  M1  
Obtain  $8\cos^2\theta + \cos\theta - 7 = 0$ , or equivalent A1  
Solve a 3-term quadratic in  $\cos\theta$  and reach  $\theta = \cos^{-1}(a)$  M1  
Obtain answer  $29.0^\circ$  A1  
Obtain answer  $180^\circ$  and no others A1 [5]  
[Ignore answers outside the given interval. Treat answers in radians (0.505 and 3.14 or  $\pi$ ) as a misread.]  
[SR: The answer  $180^\circ$  found by inspection can earn B1.]
- 4 (i) State or imply  $CT = r \tan x$  or  $OT = r \sec x$ , or equivalent B1  
Using correct area formulae, form an equation in  $r$  and  $x$  M1  
Obtain the given answer correctly A1 [3]
- (ii) Use the iterative formula correctly at least once M1  
Obtain the final answer 1.35 A1  
Show sufficient iterations to 4 d.p. to justify its accuracy to 2 d.p., or show there is a sign change in the interval (1.345, 1.355) A1 [3]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

- 5 (i) EITHER: State  $\frac{dx}{dt} = \sec^2 t / \tan t$ , or equivalent B1
- State  $\frac{dy}{dt} = 2 \sin t \cos t$ , or equivalent B1
- Use  $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$  M1
- Obtain correct answer in any form, e.g.  $2 \sin^2 t \cos^2 t$  A1
- OR: Obtain  $y = e^{2x} / (1 + e^{2x})$ , or equivalent B1
- Use correct quotient or product rule M1
- Obtain correct derivative in any form, e.g.  $2e^{2x} / (1 + e^{2x})^2$  A1
- Obtain correct derivative in terms of  $t$  in any form, e.g.  $(2 \tan^2 t) / (1 + \tan^2 t)^2$  A1 [4]
- (ii) State or imply  $t = \frac{1}{4}\pi$  when  $x = 0$  B1
- Form the equation of the tangent at  $x = 0$  M1
- Obtain correct answer in any horizontal form, e.g.  $y = \frac{1}{2}x + \frac{1}{2}$  A1 [3]
- [SR: If the OR method is used in part (i), give B1 for stating or implying  $y = \frac{1}{2}$  or  $\frac{dy}{dx} = \frac{1}{2}$  when  $x = 0$ .]
- 6 (i) Show that the differential equation is  $\frac{dy}{dx} = 2xy$  B1
- Separate variables correctly and attempt integration of both sides M1
- Obtain term  $\ln y$ , or equivalent A1
- Obtain term  $x^2$ , or equivalent A1
- Evaluate a constant, or use limits  $x = 1, y = 2$ , in a solution containing terms  $a \ln y$  and  $bx^2$  M1
- Obtain correct solution in any form A1
- Obtain the given answer correctly A1 [7]
- (ii) State that the gradient at  $(-1, 2)$  is  $-4$  B1
- Show the sketch of curve with correct concavity, positive  $y$ -intercept and axis of symmetry  $x = 0$  B1 [2]
- [SR: A solution with  $k \neq 2$ , or not evaluated, can earn B0M1A1A1M1A1A0 in part (i).]
- [SR: If given answer is assumed valid, give B1 if  $\frac{dy}{dx}$  is shown correctly to be equal to  $2xy$ , is stated to be proportional to  $xy$ , and shown to be equal to 4 at  $(1, 2)$ .]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

- 7 (a) (i) *EITHER*: Multiply numerator and denominator by  $a - 2i$ , or equivalent M1  
 Obtain final answer  $\frac{5a}{a^2 + 4} - \frac{10i}{a^2 + 4}$ , or equivalent A1  
*OR*: Obtain two equations in  $x$  and  $y$ , solve for  $x$  or for  $y$  M1  
 Obtain final answer  $x = \frac{5a}{a^2 + 4}$  and  $y = \frac{10}{a^2 + 4}$ , or equivalent A1 [2]
- (ii) Either state  $\arg(u) = -\frac{3}{4}\pi$ , or express  $u^*$  in terms of  $a$  (f.t. on  $u$ ) B1√  
 Use correct method to form an equation in  $a$ , e.g.  $5a = -10$  M1  
 Obtain  $a = -2$  correctly A1 [3]
- (b) Show a point representing  $2 + 2i$  in relatively correct position in an Argand diagram B1  
 Show the circle with centre at the origin and radius 2 B1  
 Show the perpendicular bisector of the line segment from the origin to the point representing  $2 + 2i$  B1√  
 Shade the correct region B1 [4]  
 [SR: Give the first B1 and the B1√ for obtaining  $y = 2 - x$ , or equivalent, and sketching the attempt.]
- 8 (i) State or imply partial fractions are of the form  $\frac{A}{1+x} + \frac{Bx+C}{2+x^2}$  B1  
 Use a relevant method to determine a constant M1  
 Obtain one of the values  $A = -2, B = 1, C = 4$  A1  
 Obtain a second value A1  
 Obtain the third value A1 [5]
- (ii) Use correct method to obtain the first two terms of the expansion of  $(1+x)^{-1}$ ,  
 $\left(1 + \frac{1}{2}x^2\right)^{-1}$  or  $(2+x^2)^{-1}$  in ascending powers of  $x$  M1  
 Obtain correct unsimplified expansion up to the term in  $x^3$  of each partial fraction A1√ + A1√  
 Multiply out fully by  $Bx + C$ , where  $BC \neq 0$  M1  
 Obtain final answer  $\frac{5}{2}x - 3x^2 + \frac{7}{4}x^3$ , or equivalent A1 [5]  
 [Symbolic binomial coefficients, e.g.  $\binom{-1}{1}$ , are not sufficient for the first M1. The f.t. is on  $A, B, C$ .]  
 [If  $B$  or  $C$  omitted from the form of fractions, give B0M1A0A0A0 in (i); M1A1√A1√ in (ii), max 4/10.]  
 [In the case of an attempt to expand  $(5x - x^2)(1+x)^{-1}(2+x^2)^{-1}$ , give M1A1A1 for the expansions, M1 for the multiplying out fully, and A1 for the final answer.]  
 [Allow use of Maclaurin, giving M1A1√A1√ for differentiating and obtaining  $f(0) = 0$  and  $f'(0) = \frac{5}{2}$ , A1√ for  $f''(0) = -6$ , and A1 for  $f'''(0) = \frac{21}{2}$  and the final answer (the f.t. is on  $A, B, C$  if used).]  
 [For the identity  $5x - x^2 \equiv (2 + 2x + x^2 + x^3)(a + bx + cx^2 + dx^3)$  give M1A1; then M1A1 for using a relevant method to obtain two of  $a = 0, b = \frac{5}{2}, c = -3$  and  $d = \frac{7}{4}$ ; then A1 for the final answer in series form.]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

- 9 (i) State or imply a correct normal vector to either plane, e.g.  $\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$  or  $2\mathbf{i} + \mathbf{j} + 3\mathbf{k}$  B1  
 Carry out correct process for evaluating the scalar product of the two normals M1  
 Using the correct process for the moduli, divide the scalar product by the product of the moduli and evaluate the inverse cosine of the result M1  
 Obtain the final answer  $79.7^\circ$  (or 1.39 radians) A1 [4]
- (ii) EITHER: Carry out a method for finding a point on the line M1  
 Obtain such a point, e.g. (1, 3, 0) A1  
 EITHER: State two correct equations for the direction vector ( $a, b, c$ ) of the line, e.g.  $a + 2b - 2c = 0$  and  $2a + b + 3c = 0$  B1  
 Solve for one ratio, e.g.  $a : b$  M1  
 Obtain  $a : b : c = 8 : -7 : -3$ , or equivalent A1  
 State a correct final answer, e.g.  $\mathbf{r} = \mathbf{i} + 3\mathbf{j} + \lambda(8\mathbf{i} - 7\mathbf{j} - 3\mathbf{k})$  A1√
- OR1: Obtain a second point on the line, e.g.  $\left(0, \frac{31}{8}, \frac{3}{8}\right)$  A1  
 Subtract position vectors to find a direction vector M1  
 Obtain  $\mathbf{i} - \frac{7}{8}\mathbf{j} - \frac{3}{8}\mathbf{k}$ , or equivalent A1  
 State a correct final answer, e.g.  $\mathbf{r} = \mathbf{i} + 3\mathbf{j} + \lambda\left(\mathbf{i} - \frac{7}{8}\mathbf{j} - \frac{3}{8}\mathbf{k}\right)$  A1√
- OR2: Attempt to calculate the vector product of two normals M1  
 Obtain two correct components A1  
 Obtain  $8\mathbf{i} - 7\mathbf{j} - 3\mathbf{k}$ , or equivalent A1  
 State a correct final answer, e.g.  $\mathbf{r} = \mathbf{i} + 3\mathbf{j} + \lambda(8\mathbf{i} - 7\mathbf{j} - 3\mathbf{k})$  A1√
- OR3: Express one variable in terms of a second M1  
 Obtain a correct simplified expression, e.g.  $x = (31 - 8y) / 7$  A1  
 Express the first variable in terms of a third M1  
 Obtain a correct simplified expression, e.g.  $x = (3 - 8z) / 3$  A1  
 Form a vector equation of the line M1  
 State a correct final answer, e.g.  $\mathbf{r} = \frac{31}{8}\mathbf{j} + \frac{3}{8}\mathbf{k} + \lambda(8\mathbf{i} - 7\mathbf{j} - 3\mathbf{k})$  A1√
- OR4: Express one variable in terms of a second M1  
 Obtain a correct simplified expression, e.g.  $y = (31 - 7x) / 7$  A1  
 Express the third variable in terms of the second M1  
 Obtain a correct simplified expression, e.g.  $z = (3 - 3x) / 8$  A1  
 Form a vector equation of the line M1  
 State a correct final answer, e.g.  $\mathbf{r} = \frac{31}{8}\mathbf{j} + \frac{3}{8}\mathbf{k} + \lambda(-8\mathbf{i} + 7\mathbf{j} + 3\mathbf{k})$  A1√ [6]  
 [The f.t. is dependent on all M marks having been earned.]

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2011	9709	32

- 10 (i)** Attempt integration by parts and reach  $\pm x^2e^{-x} \pm \int 2xe^{-x} dx$  M1\*
- Obtain  $-x^2e^{-x} + \int 2xe^{-x} dx$ , or equivalent A1
- Integrate and obtain  $-x^2e^{-x} - 2xe^{-x} - 2e^{-x}$ , or equivalent A1
- Use limits  $x = 0$  and  $x = 3$ , having integrated by parts twice M1(dep\*)
- Obtain the given answer correctly A1 [5]
- (ii)** Use correct product or quotient rule M1
- Obtain correct derivative in any form A1
- Equate derivative to zero and solve for non-zero  $x$  M1
- Obtain  $x = 2$  with no errors send A1 [4]
- (iii)** Carry out a complete method for finding the  $x$ -coordinate of  $P$  M1
- Obtain answer  $x = 1$  A1 [2]