

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International Advanced Level

MARK SCHEME for the October/November 2014 series

9709 MATHEMATICS

9709/32

Paper 3, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through ✓” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32
1	Use law of the logarithm of a power Obtain a correct linear equation in any form, e.g. $x = (x - 2) \ln 3$ Obtain answer $x = 22.281$	M1 A1 A1	[3]
2	(i) State or imply ordinates 2, 1.1547..., 1, 1.1547... Use correct formula, or equivalent, with $h = \frac{1}{6}\pi$ and four ordinates Obtain answer 1.95 (ii) Make recognisable sketch of $y = \operatorname{cosec} x$ for the given interval Justify a statement that the estimate will be an overestimate	B1 M1 A1 B1 B1	[3] [2]
3	Substitute $x = -\frac{1}{3}$, equate result to zero or divide by $3x + 1$ and equate the remainder to zero and obtain a correct equation, e.g. $-\frac{1}{27}a + \frac{1}{9}b - \frac{1}{3} + 3 = 0$ Substitute $x = 2$ and equate result to 21 or divide by $x - 2$ and equate constant remainder to 21 Obtain a correct equation, e.g. $8a + 4b + 5 = 21$ Solve for a or for b Obtain $a = 12$ and $b = -20$	B1 M1 A1 M1 A1	[5]
4	(i) Use chain rule correctly at least once Obtain either $\frac{dx}{dt} = \frac{3\sin t}{\cos^4 t}$ or $\frac{dy}{dt} = 3\tan^2 t \sec^2 t$, or equivalent Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ Obtain the given answer (ii) State a correct equation for the tangent in any form Use Pythagoras Obtain the given answer	M1 A1 M1 A1 B1 M1 A1	[4] [3]
5	(i) Substitute $z = 1 + i$ and obtain $w = \frac{1 + 2i}{1 + i}$ <i>EITHER:</i> Multiply numerator and denominator by the conjugate of the denominator, or equivalent Simplify numerator to $3 + i$ or denominator to 2 Obtain final answer $\frac{3}{2} + \frac{1}{2}i$, or equivalent <i>OR:</i> Obtain two equations in x and y , and solve for x or for y Obtain $x = \frac{3}{2}$ or $y = \frac{1}{2}$, or equivalent Obtain final answer $\frac{3}{2} + \frac{1}{2}i$, or equivalent	B1 M1 A1 A1 M1 A1 A1	[4]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32

- (ii) *EITHER*: Substitute $w = z$ and obtain a 3-term quadratic equation in z ,
 e.g. $iz^2 + z - i = 0$ B1
 Solve a 3-term quadratic for z or substitute $z = x + iy$ and use a correct
 method to solve for x and y M1
OR: Substitute $w = x + iy$ and obtain two correct equations in x and y by equating
 real and imaginary parts B1
 Solve for x and y M1
- Obtain a correct solution in any form, e.g. $z = \frac{-1 \pm \sqrt{3}i}{2i}$ A1
- Obtain final answer $-\frac{\sqrt{3}}{2} + \frac{1}{2}i$ A1 [4]
- 6 (i) Integrate and reach $b \ln 2x - c \int x \cdot \frac{1}{x} dx$, or equivalent M1*
- Obtain $x \ln 2x - \int x \cdot \frac{1}{x} dx$, or equivalent A1
- Obtain integral $x \ln 2x - x$, or equivalent A1
- Substitute limits correctly and equate to 1, having integrated twice M1(dep*)
- Obtain a correct equation in any form, e.g. $a \ln 2a - a + 1 - \ln 2 = 1$ A1
- Obtain the given answer A1 [6]
- (ii) Use the iterative formula correctly at least once M1
- Obtain final answer 1.94 A1
- Show sufficient iterations to 4 d.p. to justify 1.94 to 2d.p. or show that there is a sign
 change in the interval (1.935, 1.945). A1 [3]
- 7 (i) Separate variables correctly and attempt to integrate at least one side B1
- Obtain term $\ln R$ B1
- Obtain $\ln x - 0.57x$ B1
- Evaluate a constant or use limits $x = 0.5$, $R = 16.8$, in a solution containing terms of the form
 $a \ln R$ and $b \ln x$ M1
- Obtain correct solution in any form A1
- Obtain a correct expression for R , e.g. $R = xe^{(3.80 - 0.57x)}$, $R = 44.7xe^{-0.57x}$ or
 $R = 33.6xe^{(0.285 - 0.57x)}$ A1 [6]
- (ii) Equate $\frac{dR}{dx}$ to zero and solve for x M1
- State or imply $x = 0.57^{-1}$, or equivalent, e.g. 1.75 A1
- Obtain $R = 28.8$ (allow 28.9) A1 [3]
- 8 (i) Use $\sin(A + B)$ formula to express $\sin 3\theta$ in terms of trig. functions of 2θ and θ M1
- Use correct double angle formulae and Pythagoras to express $\sin 3\theta$ in terms of $\sin \theta$ M1
- Obtain a correct expression in terms of $\sin \theta$ in any form A1
- Obtain the given identity A1 [4]
- [SR: Give M1 for using correct formulae to express RHS in terms of $\sin \theta$ and $\cos 2\theta$,
 then M1A1 for expressing in terms of $\sin \theta$ and $\sin 3\theta$ only, or in terms
 of $\cos \theta$, $\sin \theta$, $\cos 2\theta$ and $\sin 2\theta$, then A1 for obtaining the given identity.]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32

- (ii) Substitute for x and obtain the given answer B1 [1]
- (iii) Carry out a correct method to find a value of x
Obtain answers 0.322, 0.799, -1.12
[Solutions with more than 3 answers can only earn a maximum of A1 + A1.] M1
A1 + A1 + A1 [4]
- 9 (i) State or imply the form $\frac{A}{1-x} + \frac{B}{2-x} + \frac{C}{(2-x)^2}$ B1
Use a correct method to determine a constant M1
Obtain one of $A = 2, B = -1, C = 3$ A1
Obtain a second value A1
Obtain a third value A1 [5]
[The alternative form $\frac{A}{1-x} + \frac{Dx+E}{(2-x)^2}$, where $A = 2, D = 1, E = 1$ is marked
B1M1A1A1A1 as above.]
- (ii) Use correct method to find the first two terms of the expansion
of $(1-x)^{-1}, (2-x)^{-1}, (2-x)^{-2}, (1-\frac{1}{2}x)^{-1}$ or $(1-\frac{1}{2}x)^{-2}$ M1
Obtain correct unsimplified expansions up to the term in x^2
of each partial fraction A1✓ + A1✓ + A1✓
Obtain final answer $\frac{9}{4} + \frac{5}{2}x + \frac{39}{16}x^2$, or equivalent A1 [5]
[Symbolic binomial coefficients, e.g. $\binom{-1}{1}$ are not sufficient for M1. The ✓ is on A, B, C .]
[For the A, D, E form of partial fractions, give M1 A1✓ A1✓ for the expansions then, if $D \neq 0$, M1 for multiplying out fully and A1 for the final answer.]
[In the case of an attempt to expand $(x^2 - 8x + 9)(1-x)^{-1}(2-x)^{-2}$, give M1A1A1 for the expansions, M1 for multiplying out fully, and A1 for the final answer.]
- 10 (i) EITHER: Find \overrightarrow{AP} (or \overrightarrow{PA}) for a point P on l with parameter λ ,
e.g. $\mathbf{i} - 17\mathbf{j} + 4\mathbf{k} + \lambda(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ B1
Calculate scalar product of \overrightarrow{AP} and a direction vector for l and equate to zero M1
Solve and obtain $\lambda = 3$ A1
Carry out a complete method for finding the length of AP M1
Obtain the given answer 15 correctly A1
OR1: Calling $(4, -9, 9)$ B , state \overrightarrow{BA} (or \overrightarrow{AB}) in component form, e.g. $-\mathbf{i} + 17\mathbf{j} - 4\mathbf{k}$ B1
Calculate vector product of \overrightarrow{BA} and a direction vector for l ,
e.g. $(-\mathbf{i} + 17\mathbf{j} - 4\mathbf{k}) \times (-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ M1
Obtain correct answer, e.g. $-30\mathbf{i} + 6\mathbf{j} + 33\mathbf{k}$ A1
Divide the modulus of the product by that of the direction vector M1
Obtain the given answer correctly A1
OR2: State \overrightarrow{BA} (or \overrightarrow{AB}) in component form B1
Use a scalar product to find the projection of BA (or AB) on l M1
Obtain correct answer in any form, e.g. $\frac{27}{\sqrt{9}}$ A1
Use Pythagoras to find the perpendicular M1

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2014	9709	32

	Obtain the given answer correctly	A1	
OR3:	State \overline{BA} (or \overline{AB}) in component form	B1	
	Use a scalar product to find the cosine of ABP	M1	
	Obtain correct answer in any form, e.g. $\frac{27}{\sqrt{9}\sqrt{306}}$	A1	
	Use trig. to find the perpendicular	M1	
	Obtain the given answer correctly	A1	
OR4:	State \overline{BA} (or \overline{AB}) in component form	B1	
	Find a second point C on l and use the cosine rule in triangle ABC to find the cosine of angle A , B , or C , or use a vector product to find the area of ABC	M1	
	Obtain correct answer in any form	A1	
	Use trig. or area formula to find the perpendicular	M1	
	Obtain the given answer correctly	A1	
OR5:	State correct \overline{AP} (or \overline{PA}) for a point P on l with parameter λ in any form	B1	
	Use correct method to express AP^2 (or AP) in terms of λ	M1	
	Obtain a correct expression in any form, e.g. $(1 - 2\lambda)^2 + (-17 + \lambda)^2 + (4 - 2\lambda)^2$	A1	
	Carry out a method for finding its minimum (using calculus, algebra or Pythagoras)	M1	
	Obtain the given answer correctly	A1	[5]
(ii) EITHER:	Substitute coordinates of a general point of l in equation of plane and either equate constant terms or equate the coefficient of λ to zero, obtaining an equation in a and b	M1*	
	Obtain a correct equation, e.g. $4a - 9b - 27 + 1 = 0$	A1	
	Obtain a second correct equation, e.g. $-2a + b + 6 = 0$	A1	
	Solve for a or for b	M1(dep*)	
	Obtain $a = 2$ and $b = -2$	A1	
OR:	Substitute coordinates of a point of l and obtain a correct equation, e.g. $4a - 9b = 26$	B1	
	EITHER: Find a second point on l and obtain an equation in a and b	M1*	
	Obtain a correct equation	A1	
	OR: Calculate scalar product of a direction vector for l and a vector normal to the plane and equate to zero	M1*	
	Obtain a correct equation, e.g. $-2a + b + 6 = 0$	A1	
	Solve for a or for b	M1(dep*)	
	Obtain $a = 2$ and $b = -2$	A1	[5]