

Cambridge
International
A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE
NAME

| |
|--|
| |
|--|

CENTRE
NUMBER

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

CANDIDATE
NUMBER

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

MATHEMATICS

9709/03

Paper 3 Pure Mathematics 3 (P3)

For Examination from 2017

SPECIMEN PAPER

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.



3 The angles θ and ϕ lie between 0° and 180° , and are such that

$$\tan(\theta - \phi) = 3 \quad \text{and} \quad \tan \theta + \tan \phi = 1.$$

Find the possible values of θ and ϕ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines spanning the width of the page, intended for writing.

4 The equation $x^3 - x^2 - 6 = 0$ has one real root, denoted by α .

(i) Find by calculation the pair of consecutive integers between which α lies. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \sqrt{\left(x_n + \frac{6}{x_n}\right)}$$

converges, then it converges to α . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use this iterative formula to determine α correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 The equation of a curve is $y = e^{-2x} \tan x$, for $0 \leq x < \frac{1}{2}\pi$.

- (i) Obtain an expression for $\frac{dy}{dx}$ and show that it can be written in the form $e^{-2x}(a + b \tan x)^2$, where a and b are constants. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Explain why the gradient of the curve is never negative. [1]

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the value of x for which the gradient is least. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 7 The points A , B and C have position vectors, relative to the origin O , given by

$$\vec{OA} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \quad \vec{OB} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{and} \quad \vec{OC} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}.$$

The plane m is perpendicular to AB and contains the point C .

- (i) Find a vector equation for the line passing through A and B . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Obtain the equation of the plane m , giving your answer in the form $ax + by + cz = d$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

9 The complex number $3 - i$ is denoted by u . Its complex conjugate is denoted by u^* .

(i) On an Argand diagram with origin O , show the points A , B and C representing the complex numbers u , u^* and $u^* - u$ respectively. What type of quadrilateral is $OABC$? [4]

(ii) Showing your working and without using a calculator, express $\frac{u^*}{u}$ in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) By considering the argument of $\frac{u^*}{u}$, prove that

$$\tan^{-1}\left(\frac{3}{4}\right) = 2 \tan^{-1}\left(\frac{1}{3}\right). \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

10



The diagram shows the curve $y = \frac{x^2}{1+x^3}$ for $x \geq 0$, and its maximum point M . The shaded region R is enclosed by the curve, the x -axis and the lines $x = 1$ and $x = p$.

(i) Find the exact value of the x -coordinate of M . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.