

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2010 question paper
for the guidance of teachers

9709 MATHEMATICS

9709/62

Paper 62, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	62

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	62

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	62

1	(i) $\bar{x} = 18.9$ (132/7) sd = 12.3	B1 B1 [2]	
	(ii) median	B1 [1]	
	(iii) mode inappropriate because it is 10 and this is the lowest value. mean inappropriate because it is affected by the outlier (of 48).	B1 B1 [2]	Sensible reason allow if seen in (ii) Sensible reason allow if seen in (ii) not 'outliers' in plural
2	(i) $P(x > 10.9) = P\left(z > \frac{10.9 - 11}{0.095}\right)$ $= P(z > -1.0526)$ $= 0.8538$ (0.854)	M1 A1 [2]	Standardising, no cc, no sq rt Rounding to correct answer
	(ii) $P(\text{at least } 2 < 10.9) = 1 - P(0, 1)$ $= 1 - (0.8538)^6 - {}^6C_1(0.1462)(0.8538)^5$ $= 0.215$	M1 A1ft A1 [3]	Bin expression with \sum powers = 6, 6C_x , $p + q = 1$. Reasonably correct unsimplified expression ft their (i) Rounding to correct answer
3	median $A = 2.0 - 2.1$ or $\bar{x}_A = 2.0 - 2.1$ median $B = 3.8 - 3.9$ or $\bar{x}_B = 3.4 - 3.5$	M1 A1	For finding medians or using mid-pts and freqs to find means, or seen on 2 box-plots Correct medians or means for A and B
	Country B has heavier babies on average	B1	Correct statement allow '...higher median...' etc.
	IQ range $A = 2.4 - 1.5 = 0.9$ or sd = 0.5 – 0.7 IQ range $B = 4.5 - 2.2 = 2.3$ or sd = 1.2 – 1.4	M1 A1	Finding spreads by IQ range or range or sd or 2 box-plots Correct IQ range or sd for A and B (± 0.1 kg) or correct IQR on box-plots
	Country B has greater spread of weights	A1 [6]	Correct statement
4	(i) $P(X < 2\mu) = P\left(z < \frac{2\mu - \mu}{\sigma}\right)$ $= P(z < \mu/\sigma) = P(z < 5/3)$ $= 0.952$	M1 A1 A1 [3]	Standardising, and attempt to get 1 variable, no cc, no $\sqrt{\quad}$, no sq $\pm 5/3$ seen oe Rounding to correct answer
	(ii) $P\left(X < \frac{\mu}{3}\right) = P\left(z < \frac{-2\mu}{3\sigma}\right)$ $\frac{-2\mu}{3\sigma} = 1.047$ $\mu = -1.57\sigma$	M1 B1 A1 [3]	standardising attempt resulting in $z \leq -$ some μ/σ allow $\pm \left(\frac{\mu/3 - \mu}{\sigma}\right)$ ± 1.047 seen correct single number, answer must have a minus sign and $\mu = \dots\sigma$
5	(i) (2,12), (3, 8), (4, 6), (6, 4), (8,3), (12,2)	M1	Listing or picking out at least 3 different options from a 12 by 12 (mult) table or seeing 3, 4, 5 or 6/144
	$P(Q) = 6/144$ (1/24) (0.0417)	A1 [2]	Correct answer

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9709	62

(ii) $P(\text{both} > 8) = 1/3 \times 1/3$ $= 1/9 = P(R) (16/144)$	M1 A1 [2]	Squaring a sensible prob or picking out 12 – 25 options Correct answer								
(iii) since $P(R \text{ and } Q) = 0$ Yes, R and Q are exclusive	B1* B1dep [2]	o.e. in words								
(iv) $P(R \text{ and } Q) = 0 \neq P(R) \times P(Q)$ or $P(R Q) = 0, \neq P(R)$ No, not independent	B1* B1dep [2]	o.e. in words								
6 (i) <table border="1" style="margin-left: 20px;"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>2</td> </tr> <tr> <td>$P(X=x)$</td> <td>1/7</td> <td>4/7</td> <td>2/7</td> </tr> </table>	x	0	1	2	$P(X=x)$	1/7	4/7	2/7	B1 B1 B1 [3]	0, 1, 2 only in table or listed with some prob 3, 4... if in table must have blank or 0 for prob One correct probability All correct
x	0	1	2							
$P(X=x)$	1/7	4/7	2/7							
(ii) $E(X) = 8/7 (1.14) \text{ AG}$ $\text{Var}(X) = 12/7 - (8/7)^2$ $= 20/49 (0.408)$	B1 M1 A1 [3]	Legitimate correct given answer rounding to 1.14 Correct method with mean ² subt numerically no dividing by anything Correct final answer								
(iii) $P(G NA) = \frac{P(G \cap NA)}{P(NA)}$ $= \frac{2/5 \times 1/4}{2/5 \times 1/4 + 3/5 \times 9/10}$ $= \frac{5}{32} (0.156)$	M1 M1 A1 A1 [4]	Attempt at $P(G \cap NA)$ or $P(G \cap A)$ as numerator of a fraction Attempt at $P(NA)$ or $P(A)$ in form of summing two 2-factor products, seen anywhere Correct unsimplified denominator of a fraction Correct answer								
7 (i) 362880 (363000)	B1 [1]									
(ii) PG or GP in $8! \times 2 = 80640$ or $7/9$ of (i) $362880 - 80640 = 282240$	M1 B1 A1ft [3]	Considering together and also subtracting from their (i) or using probabilities $8! \times 2$ or 80640 seen oe correct answer ft 40320 only								
(iii) 9P_3 or ${}^9C_3 \times 3!$ or $9!/6!$ $= 504$	M1 A1 [2]	9P_3 or 9C_3 oe seen allow extra multiplication correct final answer								
(iv) ${}^8C_2 \times 3!$ or $504 - {}^8C_3 \times 3!$ or ${}^8P_2 \times 3$ $= 168$	M1 A1 [2]	8C_x or 8P_x seen allow extra mult, or (iii)/9 or (iii)/3 correct final answer								
(v) PG and x in $7 \times 2 \times 2$ ways = 28 Answer $504 - 28 = 476$	M1 A1 [2]	$x \times 2 \times 2$ seen or their (iii) – 7 or 7C_1 or 7C_2 correct answer								