

## Paper 2: Core Pure Mathematics 2 Mark Scheme

Question	Scheme	Marks	AOs
<b>1(i)</b>	$\alpha + \beta + \gamma = 8, \quad \alpha\beta + \beta\gamma + \gamma\alpha = 28, \quad \alpha\beta\gamma = 32$	B1	3.1a
	$\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} = \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma}$	M1	1.1b
	$= \frac{7}{8}$	A1ft	1.1b
		<b>(3)</b>	
<b>(ii)</b>	$(\alpha + 2)(\beta + 2)(\gamma + 2) = (\alpha\beta + 2\alpha + 2\beta + 4)(\gamma + 2)$	M1	1.1b
	$= \alpha\beta\gamma + 2(\alpha\beta + \alpha\gamma + \beta\gamma) + 4(\alpha + \beta + \gamma) + 8$	A1	1.1b
	$= 32 + 2(28) + 4(8) + 8 = 128$	A1	1.1b
		<b>(3)</b>	
	<b>Alternative:</b>		
	$(x - 2)^3 - 8(x - 2)^2 + 28(x - 2) - 32 = 0$	M1	1.1b
	$= \dots - 8 + \dots - 32 + \dots - 56 - 32 = -128$	A1	1.1b
	$\therefore (\alpha + 2)(\beta + 2)(\gamma + 2) = 128$	A1	1.1b
	<b>(3)</b>		
<b>(iii)</b>	$\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma)$	M1	3.1a
	$= 8^2 - 2(28) = 8$	A1ft	1.1b
		<b>(2)</b>	
<b>(8 marks)</b>			
<b>Notes:</b>			
<b>(i)</b>			
<b>B1:</b> Identifies the correct values for all 3 expressions (can score anywhere)			
<b>M1:</b> Uses a correct identity			
<b>A1ft:</b> Correct value (follow through their 8, 28 and 32)			
<b>(ii)</b>			
<b>M1:</b> Attempts to expand			
<b>A1:</b> Correct expansion			
<b>A1:</b> Correct value			
<b>Alternative:</b>			
<b>M1:</b> Substitutes $x - 2$ for $x$ in the given cubic			
<b>A1:</b> Calculates the correct constant term			
<b>A1:</b> Changes sign and so obtains the correct value			
<b>(iii)</b>			
<b>M1:</b> Establishes the correct identity			
<b>A1ft:</b> Correct value (follow through their 8, 28 and 32)			



Question	Scheme	Marks	AOs
<b>2(a)</b>	$\begin{pmatrix} 3 \\ -4 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ 2 \\ 12 \end{pmatrix} = 18 - 8 + 24$	M1	3.1a
	$d = \frac{18 - 8 + 24 - 5}{\sqrt{3^2 + 4^2 + 2^2}}$	M1	1.1b
	$= \sqrt{29}$	A1	1.1b
	<b>(3)</b>		
<b>(b)</b>	$\begin{pmatrix} -1 \\ -3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix} = \dots$ and $\begin{pmatrix} -1 \\ -3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} = \dots$	M1	2.1
	$\begin{pmatrix} -1 \\ -3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix} = 0$ and $\begin{pmatrix} -1 \\ -3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} = 0$ $\therefore -\mathbf{i} - 3\mathbf{j} + \mathbf{k}$ is perpendicular to $\Pi_2$	A1	2.2a
	<b>(2)</b>		
<b>(c)</b>	$\begin{pmatrix} -1 \\ -3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -4 \\ 2 \end{pmatrix} = -3 + 12 + 2$	M1	1.1b
	$\sqrt{(-1)^2 + (-3)^2 + 1^2} \sqrt{(3)^2 + (-4)^2 + 2^2} \cos \theta = 11$ $\Rightarrow \cos \theta = \frac{11}{\sqrt{(-1)^2 + (-3)^2 + 1^2} \sqrt{(3)^2 + (-4)^2 + 2^2}}$	M1	2.1
	So angle between planes $\theta = 52^\circ$ *	A1*	2.4
	<b>(3)</b>		
<b>(8 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Realises the need to and so attempts the scalar product between the normal and the position vector			
<b>M1:</b> Correct method for the perpendicular distance			
<b>A1:</b> Correct distance			
<b>(b)</b>			
<b>M1:</b> Recognises the need to calculate the scalar product between the given vector and both direction vectors			
<b>A1:</b> Obtains zero both times and makes a conclusion			
<b>(c)</b>			
<b>M1:</b> Calculates the scalar product between the two normal vectors			
<b>M1:</b> Applies the scalar product formula with their 11 to find a value for $\cos \theta$			
<b>A1*:</b> Identifies the correct angle by linking the angle between the normal and the angle between the planes			


Question	Scheme	Marks	AOs	
<b>3(i)(a)</b>	$ \mathbf{M}  = 2(1+2) - a(-1-1) + 4(2-1) = 0 \Rightarrow a = \dots$	M1	2.3	
	The matrix $\mathbf{M}$ has an inverse when $a \neq -5$	A1	1.1b	
		<b>(2)</b>		
<b>(b)</b>	Minors : $\begin{pmatrix} 3 & -2 & 1 \\ -a-8 & 2 & a+4 \\ 4-a & -6 & -2-a \end{pmatrix}$ or Cofactors : $\begin{pmatrix} 3 & 2 & 1 \\ a+8 & 2 & -a-4 \\ 4-a & 6 & -2-a \end{pmatrix}$	B1	1.1b	
	$\mathbf{M}^{-1} = \frac{1}{ \mathbf{M} } \text{adj}(\mathbf{M})$	M1	1.1b	
	$\mathbf{M}^{-1} = \frac{1}{2a+10} \begin{pmatrix} 3 & a+8 & 4-a \\ 2 & 2 & 6 \\ 1 & -a-4 & -2-a \end{pmatrix}$	2 correct rows or columns. Follow through their det $\mathbf{M}$	A1ft	1.1b
		All correct. Follow through their det $\mathbf{M}$	A1ft	1.1b
		<b>(4)</b>		
<b>(ii)</b>	When $n = 1$ , lhs = $\begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}$ , rhs = $\begin{pmatrix} 3^1 & 0 \\ 3(3^1-1) & 1 \end{pmatrix} = \begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}$	B1	2.2a	
	So the statement is true for $n = 1$			
	Assume true for $n = k$ so $\begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}^k = \begin{pmatrix} 3^k & 0 \\ 3(3^k-1) & 1 \end{pmatrix}$	M1	2.4	
	$\begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}^{k+1} = \begin{pmatrix} 3^k & 0 \\ 3(3^k-1) & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}$	M1	2.1	
	$= \begin{pmatrix} 3 \times 3^k & 0 \\ 3 \times 3(3^k-1) + 6 & 1 \end{pmatrix}$	A1	1.1b	
	$= \begin{pmatrix} 3^{k+1} & 0 \\ 3(3^{k+1}-1) & 1 \end{pmatrix}$	A1	1.1b	
	If the statement is true for $n = k$ then it has been shown true for $n = k + 1$ and as it is true for $n = 1$ , the statement is true for all positive integers $n$	A1	2.4	
	<b>(6)</b>			
<b>(12 marks)</b>				

<b>Question 3 notes:</b>
<p><b>(i)(a)</b></p> <p><b>M1:</b> Attempts determinant, equates to zero and attempts to solve for <math>a</math> in order to establish the restriction for <math>a</math></p> <p><b>A1:</b> Provides the correct condition for <math>a</math> if <math>\mathbf{M}</math> has an inverse</p>
<p><b>(i)(b)</b></p> <p><b>B1:</b> A correct matrix of minors or cofactors</p> <p><b>M1:</b> For a complete method for the inverse</p> <p><b>A1ft:</b> Two correct rows following through their determinant</p> <p><b>A1ft:</b> Fully correct inverse following through their determinant</p>
<p><b>(ii)</b></p> <p><b>B1:</b> Shows the statement is true for <math>n = 1</math></p> <p><b>M1:</b> Assumes the statement is true for <math>n = k</math></p> <p><b>M1:</b> Attempts to multiply the correct matrices</p> <p><b>A1:</b> Correct matrix in terms of <math>k</math></p> <p><b>A1:</b> Correct matrix in terms of <math>k + 1</math></p> <p><b>A1:</b> Correct complete conclusion</p>

Question	Scheme	Marks	AOs
<b>4(a)</b>	$z^n + z^{-n} = \cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta$	M1	2.1
	$= 2 \cos n\theta^*$	A1*	1.1b
		(2)	
<b>(b)</b>	$(z + z^{-1})^4 = 16 \cos^4 \theta$	B1	2.1
	$(z + z^{-1})^4 = z^4 + 4z^2 + 6 + 4z^{-2} + z^{-4}$	M1	2.1
	$= z^4 + z^{-4} + 4(z^2 + z^{-2}) + 6$	A1	1.1b
	$= 2 \cos 4\theta + 4(2 \cos 2\theta) + 6$	M1	2.1
	$\cos^4 \theta = \frac{1}{8}(\cos 4\theta + 4 \cos 2\theta + 3)^*$	A1*	1.1b
		(5)	
<b>(7 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Identifies the correct form for $z^n$ and $z^{-n}$ and adds to progress to the printed answer			
<b>A1*:</b> Achieves printed answer with no errors			
<b>(b)</b>			
<b>B1:</b> Begins the argument by using the correct index with the result from part (a)			
<b>M1:</b> Realises the need to find the expansion of $(z + z^{-1})^4$			
<b>A1:</b> Terms correctly combined			
<b>M1:</b> Links the expansion with the result in part (a)			
<b>A1*:</b> Achieves printed answer with no errors			

Question	Scheme	Marks	AOs
<b>5(a)</b>	$\frac{dy}{dx} = \sin x \cosh x + \cos x \sinh x$	M1	1.1a
	$\frac{d^2y}{dx^2} = \cos x \cosh x + \sin x \sinh x + \cos x \cosh x - \sin x \sinh x$ (= $2 \cos x \cosh x$ )	M1	1.1b
	$\frac{d^3y}{dx^3} = 2 \cos x \sinh x - 2 \sin x \cosh x$	M1	1.1b
	$\frac{d^4y}{dx^4} = -4 \sinh x \sin x = -4y^*$	A1*	2.1
		<b>(4)</b>	
<b>(b)</b>	$\left(\frac{d^2y}{dx^2}\right)_0 = 2, \left(\frac{d^6y}{dx^6}\right)_0 = -8, \left(\frac{d^{10}y}{dx^{10}}\right)_0 = 32$	B1	3.1a
	Uses $y = y_0 + xy'_0 + \frac{x^2}{2!}y''_0 + \frac{x^3}{3!}y'''_0 + \dots$ with their values	M1	1.1b
	$= \frac{x^2}{2!}(2) + \frac{x^6}{6!}(-8) + \frac{x^{10}}{10!}(32)$	A1	1.1b
	$= x^2 - \frac{x^6}{90} + \frac{x^{10}}{113400}$	A1	1.1b
		<b>(4)</b>	
<b>(c)</b>	$2(-4)^{n-1} \frac{x^{4n-2}}{(4n-2)!}$	M1 A1	3.1a 2.2a
		<b>(2)</b>	
<b>(10 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Realises the need to use the product rule and attempts first derivative			
<b>M1:</b> Realises the need to use a second application of the product rule and attempts the second derivative			
<b>M1:</b> Correct method for the third derivative			
<b>A1*:</b> Obtains the correct 4 <sup>th</sup> derivative and links this back to y			
<b>(b)</b>			
<b>B1:</b> Makes the connection with part (a) to establish the general pattern of derivatives and finds the correct non-zero values			
<b>M1:</b> Correct attempt at Maclaurin series with their values			
<b>A1:</b> Correct expression un-simplified			
<b>A1:</b> Correct expression and simplified			
<b>(c)</b>			
<b>M1:</b> Generalising, dealing with signs, powers and factorials			
<b>A1:</b> Correct expression			

Question	Scheme	Marks	AOs
<b>6(a)(i)</b>		M1	1.1b
		A1	1.1b
<b>(a)(ii)</b>	$ z - 4 - 3i  = 5 \Rightarrow  x + iy - 4 - 3i  = 5 \Rightarrow (x - 4)^2 + (y - 3)^2 = \dots$	M1	2.1
	$(x - 4)^2 + (y - 3)^2 = 25$ or any correct form	A1	1.1b
	$(r \cos \theta - 4)^2 + (r \sin \theta - 3)^2 = 25$ $\Rightarrow r^2 \cos^2 \theta - 8r \cos \theta + 16 + r^2 \sin^2 \theta - 6r \sin \theta + 9 = 25$ $\Rightarrow r^2 - 8r \cos \theta - 6r \sin \theta = 0$	M1	2.1
	$\therefore r = 8 \cos \theta + 6 \sin \theta^*$	A1*	2.2a
	<b>(6)</b>		
<b>(b)(i)</b>		B1	1.1b
		B1ft	1.1b
<b>(b)(ii)</b>	$A = \frac{1}{2} \int r^2 d\theta = \frac{1}{2} \int (8 \cos \theta + 6 \sin \theta)^2 d\theta$ $= \frac{1}{2} \int (64 \cos^2 \theta + 96 \sin \theta \cos \theta + 36 \sin^2 \theta) d\theta$	M1	3.1a
	$= \frac{1}{2} \int (32(\cos 2\theta + 1) + 96 \sin \theta \cos \theta + 18(1 - \cos 2\theta)) d\theta$	M1	1.1b
	$= \frac{1}{2} \int (14 \cos 2\theta + 50 + 48 \sin 2\theta) d\theta$	A1	1.1b
	$= \frac{1}{2} [7 \sin 2\theta + 50\theta - 24 \cos 2\theta]_0^{\frac{\pi}{3}} = \frac{1}{2} \left\{ \left( \frac{7\sqrt{3}}{2} + \frac{50\pi}{3} + 12 \right) - (-24) \right\}$	M1	2.1
	$= \frac{7\sqrt{3}}{4} + \frac{25\pi}{3} + 18$	A1	1.1b
	<b>(7)</b>		

Question	Scheme	Marks	AOs
	<p style="text-align: center;"><b>(b) (ii) Alternative:</b></p>  <p>Candidates may take a geometric approach e.g. by finding sector + 2 triangles</p>		
	<p>Angle <math>ACB = \left(\frac{2\pi}{3}\right)</math> so area sector <math>ACB = \frac{1}{2}(5)^2 \frac{2\pi}{3}</math></p> <p>Area of triangle <math>OCB = \frac{1}{2} \times 8 \times 3</math></p>	M1	3.1a
	<p>Sector area <math>ACB</math> + triangle area <math>OCB = \frac{25\pi}{3} + 12</math></p>	A1	1.1b
	<p>Area of triangle <math>OAC</math>:</p> <p>Angle <math>ACO = 2\pi - \frac{2\pi}{3} - \cos^{-1}\left(\frac{5^2 + 5^2 - 8^2}{2 \times 5 \times 5}\right)</math></p> <p>so area <math>OAC = \frac{1}{2}(5)^2 \sin\left(\frac{4\pi}{3} - \cos^{-1}\left(\frac{-7}{25}\right)\right)</math></p>	M1	1.1b
	$= \frac{25}{2} \left( \sin \frac{4\pi}{3} \cos \left( \cos^{-1} \left( \frac{-7}{25} \right) \right) - \cos \frac{4\pi}{3} \sin \left( \cos^{-1} \left( \frac{-7}{25} \right) \right) \right)$ $= \frac{25}{2} \left( \left( \frac{7\sqrt{3}}{50} \right) + \frac{1}{2} \sqrt{1 - \left( \frac{7}{25} \right)^2} \right) = \frac{7\sqrt{3}}{4} + 6$ <p>Total area = <math>\frac{25\pi}{3} + \frac{1}{2} \times 8 \times 3 + 6 + \frac{7\sqrt{3}}{4}</math></p>	M1	2.1
	$= \frac{7\sqrt{3}}{4} + \frac{25\pi}{3} + 18$	A1	1.1b
<b>(13 marks)</b>			



<b>Question 6 notes:</b>
<p><b>(a)(i)</b>  <b>M1:</b> Draws a circle which passes through the origin  <b>A1:</b> Fully correct diagram</p>
<p><b>(a)(ii)</b>  <b>M1:</b> Uses <math>z = x + iy</math> in the given equation and uses modulus to find equation in <math>x</math> and <math>y</math> only  <b>A1:</b> Correct equation in terms of <math>x</math> and <math>y</math> in any form – may be in terms of <math>r</math> and <math>\theta</math>  <b>M1:</b> Introduces polar form, expands and uses <math>\cos^2 \theta + \sin^2 \theta = 1</math> leading to a polar equation  <b>A1*:</b> Deduces the given equation (ignore any reference to <math>r = 0</math> which gives a point on the curve)</p>
<p><b>(b)(i)</b>  <b>B1:</b> Correct pair of rays added to their diagram  <b>B1ft:</b> Area between their pair of rays and inside their circle from (a) shaded, as long as there is an intersection</p>
<p><b>(b)(ii)</b>  <b>M1:</b> Selects an appropriate method by linking the diagram to the polar curve in (a), evidenced by use of the polar area formula  <b>M1:</b> Uses double angle identities  <b>A1:</b> Correct integral  <b>M1:</b> Integrates and applies limits  <b>A1:</b> Correct area</p>
<p><b>(b)(ii) Alternative:</b>  <b>M1:</b> Selects an appropriate method by finding angle <math>ACB</math> and area of sector <math>ACB</math> and finds area of triangle <math>OCB</math> to make progress towards finding the required area  <b>A1:</b> Correct combined area of sector <math>ACB</math> + triangle <math>OCB</math>  <b>M1:</b> Starts the process of finding the area of triangle <math>OAC</math> by calculating angle <math>ACO</math> and attempts area of triangle <math>OAC</math>  <b>M1:</b> Uses the addition formula to find the exact area of triangle <math>OAC</math> and employs a full correct method to find the area of the shaded region  <b>A1:</b> Correct area</p>

Question	Scheme	Marks	AOs
<b>7(a)</b>	$r = 10 \frac{df}{dt} - 2f \Rightarrow \frac{dr}{dt} = 10 \frac{d^2f}{dt^2} - 2 \frac{df}{dt}$	M1	2.1
	$10 \frac{d^2f}{dt^2} - 2 \frac{df}{dt} = -0.2f + 0.4 \left( 10 \frac{df}{dt} - 2f \right)$	M1	2.1
	$\frac{d^2f}{dt^2} - 0.6 \frac{df}{dt} + 0.1f = 0^*$	A1*	1.1b
		<b>(3)</b>	
<b>(b)</b>	$m^2 - 0.6m + 0.1 = 0 \Rightarrow m = \frac{0.6 \pm \sqrt{0.6^2 - 4 \times 0.1}}{2}$	M1	3.4
	$m = 0.3 \pm 0.1i$	A1	1.1b
	$f = e^{\alpha t} (A \cos \beta t + B \sin \beta t)$	M1	3.4
	$f = e^{0.3t} (A \cos 0.1t + B \sin 0.1t)$	A1	1.1b
		<b>(4)</b>	
<b>(c)</b>	$\frac{df}{dt} = 0.3e^{0.3t} (A \cos 0.1t + B \sin 0.1t) + 0.1e^{0.3t} (B \cos 0.1t - A \sin 0.1t)$	M1	3.4
	$r = 10 \frac{df}{dt} - 2f$ $= e^{0.3t} ((3A+B) \cos 0.1t + (3B-A) \sin 0.1t) - 2e^{0.3t} (A \cos 0.1t + B \sin 0.1t)$	M1	3.4
	$r = e^{0.3t} ((A+B) \cos 0.1t + (B-A) \sin 0.1t)$	A1	1.1b
		<b>(3)</b>	
<b>(d)(i)</b>	$t = 0, f = 6 \Rightarrow A = 6$	M1	3.1b
	$t = 0, r = 20 \Rightarrow B = 14$	M1	3.3
	$r = e^{0.3t} (20 \cos 0.1t + 8 \sin 0.1t) = 0$	M1	3.1b
	$\tan 0.1t = -2.5$	A1	1.1b
	2019	A1	3.2a
<b>(d)(ii)</b>	3750 foxes	B1	3.4
<b>(d)(iii)</b>	e.g. the model predicts a large number of foxes are on the island when the rabbits have died out and this may not be sensible	B1	3.5a
		<b>(7)</b>	
			<b>(17 marks)</b>

<b>Question 7 notes:</b>
<p><b>(a)</b>  <b>M1:</b> Attempts to differentiate the first equation with respect to <math>t</math>  <b>M1:</b> Proceeds to the printed answer by substituting into the second equation  <b>A1*:</b> Achieves the printed answer with no errors</p>
<p><b>(b)</b>  <b>M1:</b> Uses the model to form and solve the auxiliary equation  <b>A1:</b> Correct values for <math>m</math>  <b>M1:</b> Uses the model to form the CF  <b>A1:</b> Correct CF</p>
<p><b>(c)</b>  <b>M1:</b> Differentiates the expression for the number of foxes  <b>M1:</b> Uses this result to find an expression for the number of rabbits  <b>A1:</b> Correct equation</p>
<p><b>(d)(i)</b>  <b>M1:</b> Realises the need to use the initial conditions in the model for the number of foxes  <b>M1:</b> Realises the need to use the initial conditions in the model for the number of rabbits to find both unknown constants  <b>M1:</b> Obtains an expression for <math>r</math> in terms of <math>t</math> and sets <math>= 0</math>  <b>A1:</b> Rearranges and obtains a correct value for <math>\tan</math>  <b>A1:</b> Identifies the correct year</p>
<p><b>(d)(ii)</b>  <b>B1:</b> Correct number of foxes</p>
<p><b>(d)(iii)</b>  <b>B1:</b> Makes a suitable comment on the outcome of the model</p>