



Mark Scheme (Results)

Summer 2022

Pearson Edexcel GCE

AL Further Mathematics (9FM0)

Paper 4D Decision Mathematics 2

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

June 2022

Publications Code 9FM0_4D_2206_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

General General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. These mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)

Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- **bod** – benefit of doubt
 - **ft** – follow through
 - the symbol \checkmark will be used for correct ft
 - **cao** – correct answer only
 - **cso** - correct solution only. There must be no errors in this part of the question to obtain this mark
 - **isw** – ignore subsequent working
 - **awrt** – answers which round to
 - **SC**: special case
 - **o.e.** – or equivalent (and appropriate)
 - **d** or **dep** – dependent
 - **indep** – independent
 - **dp** decimal places
 - **sf** significant figures
 - * The answer is printed on the paper or ag- answer given
4. All M marks are follow through.
 A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but answers that don't logically make sense e.g. if an answer given for a probability is >1 or <0 , should never be awarded A marks. be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response. If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is

the most complete.

7. Ignore wrong working or incorrect statements following a correct answer.

8. Mark schemes will firstly show the solution judged to be the most common response expected

from candidates. Where appropriate, alternatives answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used. If no such alternative answer is provided but the response is deemed to be valid, examiners must escalate the response for a senior examiner to review.

Question	Scheme	Marks	AOs
1(a)	Reduce rows $\begin{bmatrix} 0 & 13 & 2 & 16 \\ 0 & 2 & 13 & 9 \\ 6 & 4 & 0 & 2 \\ 0 & 2 & 5 & 9 \end{bmatrix}$ and then columns $\begin{bmatrix} 0 & 11 & 2 & 14 \\ 0 & 0 & 13 & 7 \\ 6 & 2 & 0 & 0 \\ 0 & 0 & 5 & 7 \end{bmatrix}$	M1 A1	2.1 1.1b
	Followed by $\begin{bmatrix} 0 & 11 & 0 & 12 \\ 0 & 0 & 11 & 5 \\ 8 & 4 & 0 & 0 \\ 0 & 0 & 3 & 5 \end{bmatrix}$	M1 A1ft	2.1 1.1b
	A – 3, B – 1, C – 4, D – 2 or A – 3, B – 2, C – 4, D – 1	A1ft	2.2a
		(5)	
(b)	158	B1	1.1b
		(1)	
(6 marks)			
Notes:			
<p>(a) M1: simplifying the initial matrix by reducing rows and then columns. (Allow up to 2 independent slips).</p> <p>A1:CAO</p> <p>M1: develop an improved solution – need to see one double covered +e; one uncovered –e; and one single covered unchanged. 3 lines needed to 4 lines needed (lines may be implied). If lines are drawn they must be correct.</p> <p>A1ft: CAO following on from row and column reduction final table. (f/t from previous table with no further slips).</p> <p>A1ft: correct allocation ft their optimal table (both previous M marks must have been awarded in (a)) (Must be fully written. Do not accept just indicated on zeros on final matrix).</p> <p>(b)B1:CAO – solution of original problem</p>			

Question	Scheme	Marks	AOs
2(a)	$(m+3)^2 = 0$	M1	3.1a
	$k_1 = 6$ and $k_2 = 9$	A1	1.1b
		(2)	
(b)	$u_0 = 1 \Rightarrow A(-3)^0 = 1$ $u_1 = 1 \Rightarrow (A+B)(-3) = 1$	M1	1.1b
	$A = 1$ and $B = -\frac{4}{3}$	A1	1.1b
		(2)	
(4 marks)			
Notes:			
<p>(a) M1: Correct auxiliary equation (may be implied by 6, 9 correct) from $u_n = (A + Bn)(-3)^n$ A1: CAO – allow values stated implicitly i.e., $u_{n+2} + 6u_{n+1} + 9u_n = 0$</p> <p>(b) M1: Uses $u_0 = u_1 = 1$ and attempts to find A and B A1: CAO – allow values stated implicitly e.g. $u_n = \left(1 - \frac{4}{3}n\right)(-3)^n$</p>			

Question	Scheme	Marks	AOs
3(a)		M1 A1 M1 M1 A1	3.3 1.1b 3.4 3.4 1.1b
		(5)	
(b)	Minimum expected travel time is 46.5 minutes Transport option is Train	B1ft B1	3.4 2.2a
		(2)	
(7 marks)			

Notes:

(a)M1: tree diagram with at least nine end pay-offs, one decision node and at least three chance nodes used correctly

A1: correct structure of tree diagram with each arc labelled correctly (including probabilities)

M1: at least three end-pay offs consistent with their stated probabilities (eg time 52 with probability 0.88); all nine attempted

M1: chance nodes attempted with their probabilities. Must be filled in on their diagram.

A1: cao for chance and decision nodes completed correctly

(b) B1ft: correct travel time from their completed tree diagram (dependent on all method marks earned in **(a)**)

B1: deduction of correct transport option (dependent on all method marks earned in **(a)**) including double line through inferior options in **(a)** (condone cross or single line here)

Question	Scheme	Marks	AOs
4(a)	AE, BE, BF, BG, DB, DT, SB	B1	1.1b
		(1)	
(b)	95	B1	1.1b
		(1)	
(c)	The maximum feasible flow into F is 22 (from BF and DF) but the maximum feasible flow out of F is 24 so therefore FT cannot be full to capacity	B1	2.4
		(1)	
(d)	$C_1 (= 33 + 41 + 30) = 104$ $C_2 (= 53 + 30 + 14 + 0 + 17) = 114$	B1 B1	1.1b 1.1b
		(2)	
(e)	SABDFT	B1	1.1b
		(1)	
(f)	Use of max-flow min-cut theorem Identification of cut through AE, BE, BG, BF, DF and DT, Value of cut = 98, Value of flow = 98 Therefore it follows that flow is maximal	M1 A1 A1	2.1 3.1a 2.2a
		(3)	
(9 marks)			
Notes:			
<p>(a) B1: CAO</p> <p>(b) B1: CAO</p> <p>(c) B1: Correct reasoning – argument must be numerical in nature (e.g. as a minimum comparison of 22 with 24)</p> <p>(d) B1: CAO for C_1 B1: CAO for C_2</p> <p>(e) B1: CAO</p> <p>(f) M1: Construct argument based on max-flow min-cut theorem (e.g. attempt to find a cut through saturated arcs – must contain source on one side and sink on the other). Allow a cut drawn on the diagram (need not be the correct one)</p> <p>A1: Use appropriate process of finding a minimum cut – AE, BE, BG, BF, DF and DT plus value correct and value of flow through the network stated correctly (98)</p> <p>A1: Correct deduction that the flow is maximal – must use all four words ‘maximum’, ‘flow’, ‘minimum’ and ‘cut’ (allow abbreviations for maximum and minimum) dependent on previous A1.</p>			

Question	Scheme	Marks	AOs																																																																						
5(a)	$k = 39$	B1	2.2a																																																																						
		(1)																																																																							
(b)	To ensure that the total amount transported to destination R from the four supply points cannot be less than the demand of 44	B2, 1, 0	2.4 2.4																																																																						
		(2)																																																																							
(c)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>R</th> <th>S</th> <th>T</th> </tr> </thead> <tbody> <tr> <th>A</th> <td>34</td> <td></td> <td></td> </tr> <tr> <th>B</th> <td>10</td> <td>17</td> <td></td> </tr> <tr> <th>C</th> <td></td> <td>20</td> <td>21</td> </tr> <tr> <th>D</th> <td></td> <td></td> <td>18</td> </tr> </tbody> </table>		R	S	T	A	34			B	10	17		C		20	21	D			18	B1	1.1b																																																		
	R	S	T																																																																						
A	34																																																																								
B	10	17																																																																							
C		20	21																																																																						
D			18																																																																						
	2942	B1	2.2a																																																																						
		(2)																																																																							
(d)	<table style="margin-left: auto; margin-right: auto;"> <tr> <td></td> <td></td> <td style="text-align: center;">23</td> <td style="text-align: center;">37</td> <td style="text-align: center;">39</td> </tr> <tr> <td></td> <td></td> <td style="text-align: center;">R</td> <td style="text-align: center;">S</td> <td style="text-align: center;">T</td> </tr> <tr> <td style="text-align: right;">0</td> <td style="border: 1px solid black;">A</td> <td style="border: 1px solid black;">X</td> <td style="border: 1px solid black;">-20</td> <td style="border: 1px solid black;">-15</td> </tr> <tr> <td style="text-align: right;">-8</td> <td style="border: 1px solid black;">B</td> <td style="border: 1px solid black;">X</td> <td style="border: 1px solid black;">X</td> <td style="border: 1px solid black;">1</td> </tr> <tr> <td style="text-align: right;">-12</td> <td style="border: 1px solid black;">C</td> <td style="border: 1px solid black;">14</td> <td style="border: 1px solid black;">X</td> <td style="border: 1px solid black;">X</td> </tr> <tr> <td style="text-align: right;">-14</td> <td style="border: 1px solid black;">D</td> <td style="border: 1px solid black;">10</td> <td style="border: 1px solid black;">-3</td> <td style="border: 1px solid black;">X</td> </tr> </table> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>R</th> <th>S</th> <th>T</th> </tr> </thead> <tbody> <tr> <th>A</th> <td>$34 - \theta$</td> <td>θ</td> <td></td> </tr> <tr> <th>B</th> <td>$10 + \theta$</td> <td>$17 - \theta$</td> <td></td> </tr> <tr> <th>C</th> <td></td> <td></td> <td></td> </tr> <tr> <th>D</th> <td></td> <td></td> <td></td> </tr> </tbody> </table> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>R</th> <th>S</th> <th>T</th> </tr> </thead> <tbody> <tr> <th>A</th> <td>17</td> <td>17</td> <td></td> </tr> <tr> <th>B</th> <td>27</td> <td></td> <td></td> </tr> <tr> <th>C</th> <td></td> <td>20</td> <td>21</td> </tr> <tr> <th>D</th> <td></td> <td></td> <td>18</td> </tr> </tbody> </table> <p style="text-align: center;">Entering cell is AS and exiting cell is BS</p>			23	37	39			R	S	T	0	A	X	-20	-15	-8	B	X	X	1	-12	C	14	X	X	-14	D	10	-3	X		R	S	T	A	$34 - \theta$	θ		B	$10 + \theta$	$17 - \theta$		C				D					R	S	T	A	17	17		B	27			C		20	21	D			18	M1 A1 M1 A1	2.1 1.1b 1.1b 2.2a
		23	37	39																																																																					
		R	S	T																																																																					
0	A	X	-20	-15																																																																					
-8	B	X	X	1																																																																					
-12	C	14	X	X																																																																					
-14	D	10	-3	X																																																																					
	R	S	T																																																																						
A	$34 - \theta$	θ																																																																							
B	$10 + \theta$	$17 - \theta$																																																																							
C																																																																									
D																																																																									
	R	S	T																																																																						
A	17	17																																																																							
B	27																																																																								
C		20	21																																																																						
D			18																																																																						
		(4)																																																																							

(9 marks)

Notes:

(a) B1: CAO**(b) B1:** Partial correct reasoning – must include at least two of ‘destination **R**’, ‘**supply** points’, ‘**cannot be less**’/‘**must be at least**’, ‘demand of **44**’ oe (do **not** accept ‘greater than or equal to’)**B1:** Fully correct reasoning – all points covered as stated above. No incorrect statement.**(c) B1:** CAO for north-west corner method (six correct figures in correct cells only, no zeros)**B1:** CAO for initial solution (2942)**(d) M1:** Finding 7 shadow costs and 6 improvement indices**A1:** CAO

M1: A valid route shown, their most negative Π chosen, only one empty square used, θ 's balance
A1:cao – (no zeros) including deducing entering and exiting cells

For reference:

	R	S	T	Supply
A	23	17	24	34
B	15	29	32	27
C	25	25	27	41
D	19	20	25	18
Demand	44	37	k	

Qu	Scheme					Marks	AOs	
6	Stage	State	Action	Dest	Value			
	May	2	0	0	160 = 160*	M1	3.1a	
	(2)	1	1	0	80 + 35 = 115*			
		0	2	0	70 = 70*			
	April	3	2	0	240 + 70 + 70 = 380*	A1	1.1b	
	(5)		3	1	240 + 105 + 250 + 115 = 710			
			4	2	240 + 140 + 250 + 160 = 790			
		2	3	0	160 + 105 + 250 + 70 = 585*	A1	1.1b	
			4	1	160 + 140 + 250 + 115 = 665			
		1	4	0	80 + 140 + 250 + 70 = 540*			
	March	3	1	1	240 + 35 + 540 = 815*	M1	3.1a	
	(3)		2	2	240 + 70 + 585 = 895			
			3	3	240 + 105 + 250 + 380 = 975			
		2	2	1	160 + 70 + 540 = 770*	A1ft	1.1b	
			3	2	160 + 105 + 250 + 585 = 1100			
			4	3	160 + 140 + 250 + 380 = 930			
		1	3	1	80 + 105 + 250 + 540 = 975*	A1	1.1b	
			4	2	80 + 140 + 250 + 585 = 1055			
		0	4	1	140 + 250 + 540 = 930*			
	Feb	3	0	0	240 + 930 = 1170	M1	1.1b	
	(3)		1	1	240 + 35 + 975 = 1250			
			2	2	240 + 70 + 770 = 1080*			
			3	3	240 + 105 + 250 + 815 = 1410	A1ft	1.1b	
		2	1	0	160 + 35 + 930 = 1125*			
			2	1	160 + 70 + 975 = 1205			
			3	2	160 + 105 + 250 + 770 = 1285	A1ft	1.1b	
			4	3	160 + 140 + 250 + 815 = 1365			
		1	2	0	80 + 70 + 930 = 1080*			
			3	1	80 + 105 + 250 + 975 = 1410	A1	1.1b	
			4	2	80 + 140 + 250 + 770 = 1240			
		0	3	0	105 + 250 + 930 = 1285*			
			4	1	140 + 250 + 975 = 1365	M1	1.1b	
	Jan	0	1	0	35 + 1285 = 1320			
	(1)		2	1	70 + 1080 = 1150*			
			3	2	105 + 250 + 1125 = 1480	A1	1.1b	
			4	3	140 + 250 + 1080 = 1470			
		Month	January	February	March	April	May	
		Number made	2	2	4	4	2	B1
								2.2a
								B1
								1.1b
	Minimum Cost: (£) 1150							
							(14 marks)	

Notes:

All M marks – must bring earlier optimal results into calculations. Ignore extra rows. Must have right ‘ingredients’ (storage costs, overhead costs and additional workers) at least once per stage. Ingredients may be summed into a single figure. Correct total figure implies correct calculation. Penalise lack of * only once per question.

M1: Second stage completed. 6 rows, something in each cell.

A1: Any two states correct

A1: CAO for second stage

M1: Third stage completed. 9 rows with correct state, action and destination. Something in each cell. Condone at least 8 rows, with correct state, action and destination.

A1ft: Any two states correct – ft their optimal values

A1: CAO for third stage

M1: Fourth stage completed. 13 rows, with correct state, action and destination. Something in each cell. Condone at least 11 rows, with correct state, action and destination.

A1ft: Any two states correct – ft their optimal values

A1ft: Any three states correct – ft their optimal values

A1: CAO for fourth stage

M1: Fifth stage completed. 4 rows, something in each cell.

A1: CAO for fifth stage

B1: Correct allocation (dependent on all previous M marks)

B1: Correct minimum cost (dependent on all previous M marks)

Qu	Scheme	Marks	AOs																																																												
7(a)	Row minima are -2 , $\min(k, -4)$ and -3	M1	1.1b																																																												
	e.g., If the row minimum for option R is -4 then the play safe is Q (as -2 is greater than -4 and -3) If the row minimum for option R is k then $-4 > k$ and the play safe is still Q	A1	2.3																																																												
		(2)																																																													
(b)	Column maxima are 4 , 6 , 3 and $\max(k, -2)$	B1	1.1b																																																												
	Since the play-safe is option Z, $k < 3$ or $k \leq 2$	B1	2.2a																																																												
		(2)																																																													
(c)	e.g. Option Y dominates option X	B1	1.2																																																												
	Because e.g. $-1 < 3$, $-4 < 5$ and $3 < 6$	B1	2.4																																																												
		(2)																																																													
(d) (i)	$\begin{pmatrix} 4 & -1 & -2 \\ -3 & -4 & k \\ -1 & 3 & -3 \end{pmatrix} \rightarrow \begin{pmatrix} 8 & 3 & 2 \\ 1 & 0 & k+4 \\ 3 & 7 & 1 \end{pmatrix}$	B1	1.1b																																																												
	$V - 8p_1 - p_2 - 3p_3 + r = 0$ $V - 3p_1 - 7p_3 + s = 0$ $V - 2p_1 - (k+4)p_2 - p_3 + t = 0$	M1 A1	3.3 2.5																																																												
	$p_1 + p_2 + p_3 + u = 1$	B1	3.3																																																												
(ii)	e.g. <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>b.v.</th> <th>V</th> <th>p_1</th> <th>p_2</th> <th>p_3</th> <th>r</th> <th>s</th> <th>t</th> <th>u</th> <th>Value</th> </tr> </thead> <tbody> <tr> <td>r</td> <td>1</td> <td>-8</td> <td>-1</td> <td>-3</td> <td>1</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>s</td> <td>1</td> <td>-3</td> <td>0</td> <td>-7</td> <td>0</td> <td>1</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>t</td> <td>1</td> <td>-2</td> <td>$-(k+4)$</td> <td>-1</td> <td>0</td> <td>0</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>u</td> <td>0</td> <td>1</td> <td>1</td> <td>1</td> <td>0</td> <td>0</td> <td>0</td> <td>1</td> <td>1</td> </tr> <tr> <td>P</td> <td>-1</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> </tbody> </table>	b.v.	V	p_1	p_2	p_3	r	s	t	u	Value	r	1	-8	-1	-3	1	0	0	0	0	s	1	-3	0	-7	0	1	0	0	0	t	1	-2	$-(k+4)$	-1	0	0	1	0	0	u	0	1	1	1	0	0	0	1	1	P	-1	0	0	0	0	0	0	0	0	B1 M1 A1	1.2 3.3 1.1b
b.v.	V	p_1	p_2	p_3	r	s	t	u	Value																																																						
r	1	-8	-1	-3	1	0	0	0	0																																																						
s	1	-3	0	-7	0	1	0	0	0																																																						
t	1	-2	$-(k+4)$	-1	0	0	1	0	0																																																						
u	0	1	1	1	0	0	0	1	1																																																						
P	-1	0	0	0	0	0	0	0	0																																																						
		(7)																																																													
(e)	$p_3 = \frac{13}{37}$	B1	1.1b																																																												
	$V - 8\left(\frac{7}{37}\right) - \left(\frac{17}{37}\right) - 3\left(\frac{13}{37}\right) + 0 = 0$ or $V - 3\left(\frac{7}{37}\right) - 7\left(\frac{13}{37}\right) + 0 = 0$	M1	3.1a																																																												

	$V = \frac{112}{37} \Rightarrow \frac{112}{37} - 2\left(\frac{7}{37}\right) - (k+4)\left(\frac{17}{37}\right) - \left(\frac{13}{37}\right) + 0 = 0$	dM1	3.4
	$k = 1$	A1	2.2a
		(4)	
(17 marks)			
Notes:			
<p>(a)M1: Attempt to calculate row minima. Condone 'k or -4' for $\min(k, -4)$. A1: Correct argument/conditions for why the play-safe for player A is always their option Q</p> <p>(b)B1: Attempt to calculate column maxima. Condone 'k or -2' for $\max(k, -2)$ B1: CAO ignore lower limit figure, if given</p> <p>(c) B1: Correct statement – must include the word 'dominate'. Also e.g. option Z dominates option X B1: Correct inequalities – must be clear that all three inequalities must hold</p> <p>(d)(i) B1: Correct augmentation – possibly implied by later working, X column may be included M1: At least three equations in V, p_1, p_2, p_3 and at least one dummy variable seen A1: CAO (ignore extra probability equation, if seen, see note below) B1: Correct probability equation</p> <p>(ii) B1: Correct row and column labels for Simplex tableau M1: Any one (numerical in nature) row correct A1: CAO</p> <p>(e) B1: p_3 correctly stated M1: Attempts to calculate V using either equation from (d) not involving k, or attempts to eliminate V from two equations, one involving k dM1: Dependent on previous M mark – either uses equation in k and their V to calculate k, or eliminates V from two equations to calculate k A1: CAO $k = 1$</p> <p>Special case: (d) If augmentation with +5 maximum marks possible B0 M1 A1 B1 B1 M1 A0</p> <p>Note: (d)(i) A1 The extra probability equation you may see is $V - 7p_1 - 9p_2 - 10p_3 + \text{slack} = 0$</p>			

Qu	Scheme	Marks	AOs
8(a)	$u_{n+1} = pu_n + k$	B1	3.3
	(aux equation $m - p = 0 \Rightarrow$) complementary function is $A(p)^n$	B1	1.1b
	Consider a trial solution of the form $u_n = \lambda$ so $\lambda - p\lambda = k$	M1	1.1b
	General solution is $u_n = A(p)^n + \frac{k}{1-p}$	A1	1.1b
	$u_1 = 5000 \Rightarrow 5000 = A(p) + \frac{k}{1-p}$ and solve for A	M1	3.4
	$u_n = \left(5000 - \frac{k}{1-p}\right)p^{n-1} + \frac{k}{1-p}$	A1	2.2a
		(6)	
(b)	Set $k = 10\,000$, $p = 0.95$ and $u_m \dots 135\,000$	B1	3.1b
	$\left(5000 - \frac{10000}{1-0.95}\right)(0.95)^{m-1} + \frac{10000}{1-0.95} \dots 135000$		
	$(0.95)^{m-1} \dots \frac{1}{3} \Rightarrow (m-1)\log(0.95) \dots \log\left(\frac{1}{3}\right) \Rightarrow m \dots$	M1	1.1b
	$m \dots 22.418 \dots$ so 23 months after the company was first up	A1	3.2a
		(3)	
(9 marks)			
Notes:			
(a) B1:CAO			
B1:CAO			
M1: substituting their trial solution into the recurrence relation in an attempt to find their λ (which if correct is $\frac{k}{1-p}$)			
A1: CAO for the general solution			
M1: using the conditions in the model to calculate A (which if correct is $p^{-1}\left(5000 - \frac{k}{1-p}\right)$)			
A1: CAO for the particular solution (oe)			
(b)B1: Applying $u_m \dots 135000$ (or equality or strict inequality) to their general solution together with correct values for k and p (dependent on both M marks in (a))			
M1: dependent on previous B mark – solving their equation using logarithms			
A1: CAO – must be rounded to 23			
Special case: A common misread is 500 for 5000. Mark as a misread so final A marks in (a) and (b) deducted.			

