

Please check the examination details below before entering your candidate information

Candidate surname					Other names									
<b>Pearson Edexcel</b>					Centre Number					Candidate Number				
<b>Level 3 GCE</b>					<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>					<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>				
Time 1 hour 30 minutes					Paper reference					<b>9FM0/3C</b>				
<b>Further Mathematics</b>														
<b>Advanced</b>														
<b>PAPER 3C: Further Mechanics 1</b>														
<b>You must have:</b>										Total Marks				
Mathematical Formulae and Statistical Tables (Green), calculator														

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$  and give your answer to either 2 significant figures or 3 significant figures.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ►

P66800A

©2021 Pearson Education Ltd.

1/1/1/



1. A van of mass 900 kg is moving along a straight horizontal road.

At the instant when the speed of the van is  $v \text{ m s}^{-1}$ , the resistance to the motion of the van is modelled as a force of magnitude  $(500 + 7v) \text{ N}$ .

When the engine of the van is working at a constant rate of 18 kW, the van is moving along the road at a constant speed  $V \text{ m s}^{-1}$ .

- (a) Find the value of  $V$ .

(5)

Later on, the van is moving up a straight road that is inclined to the horizontal at an angle  $\theta$ , where  $\sin \theta = \frac{1}{21}$ .

At the instant when the speed of the van is  $v \text{ m s}^{-1}$ , the resistance to the motion of the van from non-gravitational forces is modelled as a force of magnitude  $(500 + 7v) \text{ N}$ .

The engine of the van is again working at a constant rate of 18 kW.

- (b) Find the acceleration of the van at the instant when  $v = 15$ .

(4)

a)  $\xrightarrow{v}$



Resolve  $\rightarrow +$  :  $F = 500 + 7v$  (1)

$18,000 = Fv \Rightarrow F = \frac{18,000}{v}$  (1)

$\frac{18,000}{v} = 500 + 7v$  (1)

$18,000 = 500v + 7v^2$

$7v^2 + 500v - 18,000 = 0$  (1)


$v = 26.309... \text{ or } v = -97.737...$

choose  $v = 26.0$  as  $v > 0$  (1) speed is a scalar quantity so it cannot be negative.



## Question 1 continued

b)



$$\sin \theta = 1/21$$

$$18,000 = 15F \Rightarrow F = \frac{18,000}{15}$$

$$\text{Resolve } \nearrow + : \frac{18,000}{15} - (500 + 7 \times 15) - 900g \sin \theta = 900a \quad (1)$$

$$a = \frac{1200 - 605 - 420}{900}$$

$$a = 0.19 \text{ m s}^{-2} \text{ (2sf)} \quad (1)$$



Question 1 continued

Lined area for student response

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





2. Two particles, A and B, are moving in opposite directions along the same straight line on a smooth horizontal surface when they collide directly.

Particle A has mass  $5m$  and particle B has mass  $3m$ .

The coefficient of restitution between A and B is  $e$ , where  $e > 0$

Immediately after the collision the speed of A is  $v$  and the speed of B is  $2v$ .

Given that A and B are moving in the same direction after the collision,

- (a) find the set of possible values of  $e$ .


(8)

Given also that the kinetic energy of A immediately after the collision is 16% of the kinetic energy of A immediately before the collision,

- (b) find

- (i) the value of  $e$ ,
- (ii) the magnitude of the impulse received by A in the collision, giving your answer in terms of  $m$  and  $v$ .

(6)



can't be moving left after collision as then B would be moving through A.



Conservation of Linear Momentum:  $\rightarrow +$

$$5mx - 3my = 5mv + 3 \cdot 2v \quad (1)$$

$$5x - 3y = 11v \Rightarrow 5ex - 3ey = 11ev \quad (1)$$

Newton's Law of Restitution:

$$e = \frac{2v - v}{x + y} \quad (1)$$

$$ex + ey = v \Rightarrow 3ex + 3ey = 3v \quad (2)$$



## Question 2 continued

$$\textcircled{1} + \textcircled{2}: 5ex + 3ex = 11ev + 3v$$

$$8ex = v(11e + 3)$$

$$x = \frac{v}{8e} (11e + 3) \quad \textcircled{1}$$

sub into  $5x - 3y = 11v$

$$3y = 5x - 11v$$

$$= \frac{55v}{8} + \frac{15v}{8e} - 11v$$

$$3y = \frac{15v}{8e} - \frac{33v}{8}$$

$$y = \frac{v}{8e} (5 - 11e) \quad \textcircled{1}$$

since  $e > 0$ ,  $x > 0$ . also,  $y > 0$  so

$$\frac{v}{8e} (5 - 11e) > 0$$

$$5 - 11e > 0 \quad \textcircled{1}$$

$$e < \frac{5}{11}$$

$$0 < e < \frac{5}{11} \quad \textcircled{1}$$



## Question 2 continued

$$b)(i) \frac{1}{2} \times 5m \times v^2 = \frac{16}{100} \times \frac{1}{2} \times 5m \times \left[ \frac{v}{8e} (11e+3) \right]^2 \quad (1)$$

$$v^2 = 0.16 \left( \frac{v^2}{64e^2} (11e+3)^2 \right)$$

$$64e^2 = 0.16(121e^2 + 66e + 9)$$

$$44.64e^2 - 10.56e - 1.44 = 0$$

$$e = \frac{1}{3} \text{ or } e = -\frac{3}{31}$$

choose  $e = \frac{1}{3}$  as  $e > 0$  (1)

(ii) Impulse received by A = " $m(v-u)$ "

$$(+\leftarrow): I = 5m(-v - -x) \\ = 5m(x-v) \quad (1)$$

find  $x$  in terms of  $v$  only:

$$x = \frac{v}{8 \times \frac{1}{3}} \left( 11 \times \frac{1}{3} + 3 \right)$$

$$x = \frac{5v}{2}$$

$$I = 5m \left( \frac{5v}{2} - v \right) \quad (1)$$

$$= \frac{15mv}{2} \quad (1)$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







3. [In this question,  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular unit vectors in a horizontal plane.]

A smooth uniform sphere  $P$  has mass  $0.3 \text{ kg}$ . Another smooth uniform sphere  $Q$ , with the same radius as  $P$ , has mass  $0.5 \text{ kg}$ .

The spheres are moving on a smooth horizontal surface when they collide obliquely. Immediately before the collision the velocity of  $P$  is  $(u\mathbf{i} + 2\mathbf{j}) \text{ ms}^{-1}$ , where  $u$  is a positive constant, and the velocity of  $Q$  is  $(-4\mathbf{i} + 3\mathbf{j}) \text{ ms}^{-1}$


At the instant when the spheres collide, the line joining their centres is parallel to  $\mathbf{i}$ .

The coefficient of restitution between  $P$  and  $Q$  is  $\frac{3}{5}$

As a result of the collision, the direction of motion of  $P$  is deflected through an angle of  $90^\circ$  and the direction of motion of  $Q$  is deflected through an angle of  $\alpha^\circ$

- (a) Find the value of  $u$  (8)
- (b) Find the value of  $\alpha$  (5)
- (c) State how you have used the fact that  $P$  and  $Q$  have equal radii. (1)

a)



Perp to LOC  
 $a=2$  ①  
 $b=3$   
 P deflected by  $90^\circ$  so  $\begin{pmatrix} u \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -x \\ 2 \end{pmatrix} = 0$

$$-ux + 4 = 0 \Rightarrow x = \frac{4}{u} \text{ ①}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



## Question 3 continued

// to LOC

CLM ( $\rightarrow +$ ):

$$0.3u - 0.5(4) = 0.3\left(-\frac{4}{u}\right) + 0.5y \quad (1)$$

$$3u - 20 = -\frac{12}{u} + 5y \quad (1) \quad (1)$$

NLR:

$$e = \frac{\frac{4}{u} + y}{u+4} \Rightarrow \frac{3}{5}(u+4) = \frac{4}{u} + y \quad (1)$$

$$3u + 12 = \frac{20}{u} + 5y \quad (2)$$

$$(2) - (1): 3u + 12 - 3u + 20 = \frac{20}{u} + 5y + \frac{12}{u} - 5y \quad (1)$$


$$32 = \frac{32}{u} \Rightarrow u = 1 \quad (1)$$

$$b) \text{ from } (1): 3(1) - 20 = -\frac{12}{1} + 5y$$

$$5y = -5$$

$$y = -1 \quad (1)$$

for Q:



$$\cos \alpha = \frac{\begin{pmatrix} -1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} -4 \\ 3 \end{pmatrix}}{\sqrt{1^2+3^2} \sqrt{4^2+3^2}} = \frac{4+9}{5\sqrt{10}} \quad (1)$$

$$\alpha = 34.7^\circ \quad (3sf) \quad (1)$$



## Question 3 continued

c) equal radii: the line of centres (LOC) is parallel to the surface, so the impulse acts parallel to the surface. ①

3D view



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**Question 3 continued**

Lined area for writing answers, consisting of 30 horizontal lines.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**(Total for Question 3 is 14 marks)**



P 6 6 8 0 0 A 0 1 3 2 8

4. A particle  $P$  has mass  $0.5 \text{ kg}$ . It is moving in the  $xy$  plane with velocity  $8\mathbf{i} \text{ m s}^{-1}$  when it receives an impulse  $\lambda(-\mathbf{i} + \mathbf{j}) \text{ N s}$ , where  $\lambda$  is a positive constant.

The angle between the direction of motion of  $P$  immediately before receiving the impulse and the direction of motion of  $P$  immediately after receiving the impulse is  $\theta^\circ$


Immediately after receiving the impulse,  $P$  is moving with speed  $4\sqrt{10} \text{ m s}^{-1}$

Find (i) the value of  $\lambda$

(ii) the value of  $\theta$

(8)

(i)



considering impulse: " $\underline{I} = m(\underline{v} - \underline{u})$ "

$$\begin{pmatrix} -\lambda \\ \lambda \end{pmatrix} = 0.5 \left[ \underline{v} - \begin{pmatrix} 8 \\ 0 \end{pmatrix} \right]$$

$$2 \begin{pmatrix} -\lambda \\ \lambda \end{pmatrix} = \underline{v} - \begin{pmatrix} 8 \\ 0 \end{pmatrix}$$

$$\underline{v} = \begin{pmatrix} -2\lambda + 8 \\ 2\lambda \end{pmatrix}$$

$$|\underline{v}| = 4\sqrt{10} \Rightarrow \sqrt{(-2\lambda + 8)^2 + (2\lambda)^2} = 4\sqrt{10}$$

$$4\lambda^2 - 32\lambda + 64 + 4\lambda^2 = 160$$

$$8\lambda^2 - 32\lambda - 96 = 0$$

$$\lambda^2 - 4\lambda + 12 = 0$$

$$(\lambda - 6)(\lambda + 2) = 0$$

$$\lambda = 6 \text{ or } \lambda = -2$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



## Question 4 continued

choose  $\lambda = 6$  as  $\lambda$  is a positive constant ①

(ii)  $\underline{v} = \begin{pmatrix} -4 \\ 12 \end{pmatrix}$



$$\theta = 180^\circ - \tan^{-1}\left(\frac{12}{4}\right) \quad \text{①}$$

$$\theta = 108^\circ \quad (3\text{sf}) \quad \text{①}$$

(Total for Question 4 is 8 marks)



5.




Figure 1

Figure 1 represents the plan view of part of a horizontal floor, where  $AB$  and  $BC$  represent fixed vertical walls, with  $AB$  perpendicular to  $BC$ .

A small ball is projected along the floor towards the wall  $AB$ . Immediately before hitting the wall  $AB$  the ball is moving with speed  $v \text{ m s}^{-1}$  at an angle  $\theta$  to  $AB$ .

The ball hits the wall  $AB$  and then hits the wall  $BC$ .

The coefficient of restitution between the ball and the wall  $AB$  is  $\frac{1}{3}$

The coefficient of restitution between the ball and the wall  $BC$  is  $e$ .

The floor and the walls are modelled as being smooth.


The ball is modelled as a particle.

The ball loses half of its kinetic energy in the impact with the wall  $AB$ .

(a) Find the exact value of  $\cos \theta$ . (5)

The ball loses half of its remaining kinetic energy in the impact with the wall  $BC$ .

(b) Find the exact value of  $e$ . (5)



$$KE \text{ before: } \frac{1}{2} \times m \times v^2 = \frac{1}{2} m v^2$$

$$KE \text{ after: } \frac{1}{2} \times m \times (v^2 \cos^2 \theta + \frac{1}{9} v^2 \sin^2 \theta)$$





## Question 5 continued

$$KE \text{ after} = \frac{1}{2} KE \text{ before}$$

$$\frac{1}{2} m (v^2 \cos^2 \theta + \frac{1}{9} v^2 \sin^2 \theta) = \frac{1}{2} \times \frac{1}{2} mv^2 \quad (1)$$


$$\cos^2 \theta + \frac{1}{9} (1 - \cos^2 \theta) = \frac{1}{2} \quad (1)$$

$$18 \cos^2 \theta + 2 - 2 \cos^2 \theta = 9$$

$$\cos^2 \theta = \frac{7}{16}$$

$$\cos \theta = \frac{\sqrt{7}}{4} \quad (1)$$

b)



$$\cos^2 \theta = \frac{7}{16} \Rightarrow \sin^2 \theta = 1 - \frac{7}{16} = \frac{9}{16}$$

$$KE \text{ after second impact} = \frac{1}{4} KE \text{ before first impact}$$

$$\frac{1}{2} m (e^2 v^2 \cos^2 \theta + \frac{1}{9} v^2 \sin^2 \theta) = \frac{1}{4} \times \frac{1}{2} mv^2 \quad (1)$$

$$\frac{7e^2}{16} + \frac{1}{9} \times \frac{9}{16} = \frac{1}{4} \quad (1)$$

$$e^2 = \frac{3}{7} \Rightarrow e = \sqrt{\frac{3}{7}} \quad (1)$$



Question 5 continued

Lined writing area for the answer to Question 5.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





6.




Figure 2

A light elastic spring has natural length  $3l$  and modulus of elasticity  $3mg$ .

One end of the spring is attached to a fixed point  $X$  on a rough inclined plane.

The other end of the spring is attached to a package  $P$  of mass  $m$ .

The plane is inclined to the horizontal at an angle  $\alpha$  where  $\tan \alpha = \frac{3}{4}$

The package is initially held at the point  $Y$  on the plane, where  $XY = l$ . The point  $Y$  is higher than  $X$  and  $XY$  is a line of greatest slope of the plane, as shown in Figure 2.

The package is released from rest at  $Y$  and moves up the plane.

The coefficient of friction between  $P$  and the plane is  $\frac{1}{3}$

By modelling  $P$  as a particle,

(a) show that the acceleration of  $P$  at the instant when  $P$  is released from rest is  $\frac{17}{15}g$  (5)

(b) find, in terms of  $g$  and  $l$ , the speed of  $P$  at the instant when the spring first reaches its natural length of  $3l$ . (6)

a) spring


$l = 3l$

$\lambda = 3mg$

$\tan \alpha = 3/4$

$\therefore \sin \alpha = 3/5$

$\cos \alpha = 4/5$



The spring is being compressed so there is thrust in the spring

Resolve  $\uparrow$  :  $T - F - mg \sin \alpha = ma$  (1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

## Question 6 continued

find  $T$  and  $F$ :

$$T = \frac{\lambda x}{L} = \frac{3mg \times 2L}{3L} = 2mg \quad (1)$$

$$F = \mu R$$

Resolve  $\uparrow$ :  $R = mg \cos \alpha$ 

$$F = \frac{1}{3} mg \cos \alpha$$

$$2mg - \frac{1}{3} mg \cos \alpha - mg \sin \alpha = ma \quad (1)$$

$$2mg - \frac{4}{15} mg - \frac{3}{5} mg = ma \quad (1)$$

$$a = \frac{17g}{15} \quad (1)$$

b) KE before + GPE before + EPE before = KE after + GPE after + EPE after  
+ work done against friction

$$0 + 0 + \frac{3mg(2L)^2}{2(3L)} \quad (1) = \frac{1}{2} mv^2 + mg(2L \sin \alpha) \quad (1) + 0 + 2L \left( \frac{1}{3} mg \cos \alpha \right) \quad (1)$$

$$2mgl \quad (1) = \frac{1}{2} mv^2 + \frac{6mgl}{5} + \frac{8mgl}{15} \quad (1)$$

$$v^2 = \frac{8gl}{15}$$

w.d. against  
friction =  $\mu R d$

$$v = \sqrt{\frac{8gl}{15}} \quad (1)$$







7. [In this question,  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular unit vectors in a horizontal plane.]




Figure 3

Figure 3 represents the plan view of part of a smooth horizontal floor, where  $AB$  is a fixed smooth vertical wall.

The direction of  $\overrightarrow{AB}$  is in the direction of the vector  $(\mathbf{i} + \mathbf{j})$

A small ball of mass  $0.25 \text{ kg}$  is moving on the floor when it strikes the wall  $AB$ .

Immediately before its impact with the wall  $AB$ , the velocity of the ball is  $(8\mathbf{i} + 2\mathbf{j}) \text{ ms}^{-1}$

Immediately after its impact with the wall  $AB$ , the velocity of the ball is  $\mathbf{v} \text{ ms}^{-1}$

The coefficient of restitution between the ball and the wall is  $\frac{1}{3}$

By modelling the ball as a particle,

(a) show that  $\mathbf{v} = 4\mathbf{i} + 6\mathbf{j}$  (6)

(b) Find the magnitude of the impulse received by the ball in the impact. (3)

a) let  $\underline{v} = a\underline{i} + b\underline{j}$

using  $\underline{u} \cdot \underline{W} = \underline{v} \cdot \underline{W}$  where  $\underline{W} =$  direction of wall vector  $(\underline{i} + \underline{j})$

$$\begin{pmatrix} 8 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \end{pmatrix} \Rightarrow a + b = 8 + 2$$

$$a + b = 10 \quad \textcircled{1} \quad \textcircled{1}$$

using  $e\underline{u} \cdot \underline{I} = -\underline{v} \cdot \underline{I}$  where  $\underline{I} =$  direction of impulse:  $(-\underline{i} + \underline{j})$

$$\frac{1}{3} \begin{pmatrix} 8 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \end{pmatrix} = - \begin{pmatrix} a \\ b \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$





## Question 7 continued

$$\Rightarrow \frac{1}{3}(-8+2) = -(-a+b)$$

$$-2 = a-b \quad (2) \quad (1)$$

$$\begin{aligned} (1) + (2): a+b+a-b &= 10-2 \quad (1) \\ 2a &= 8 \\ a &= 4 \end{aligned}$$

$$\text{sub into (1): } 4+b=10$$

$$b=6$$

$$\therefore \underline{v} = 4\underline{i} + 6\underline{j} \text{ as required } (1)$$

$$b) \underline{I} = m(\underline{v} - \underline{u})$$

$$= 0.25 \left( \begin{pmatrix} 4 \\ 6 \end{pmatrix} - \begin{pmatrix} 8 \\ 2 \end{pmatrix} \right) \quad (1)$$

$$= 0.25 \begin{pmatrix} -4 \\ 4 \end{pmatrix}$$

$$= \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$\text{magnitude} = \sqrt{1^2 + 1^2} = \sqrt{2} \text{ N s} \quad (1)$$



Question 7 continued

Lined writing area for the answer to Question 7.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

Lined writing area with multiple horizontal lines for student responses.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

Lined area for student answers, containing 22 horizontal lines.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 9 marks)

TOTAL FOR PAPER IS 75 MARKS

