

Please check the examination details below before entering your candidate information	
Candidate surname	Other names
Centre Number	Candidate Number
Pearson Edexcel Level 3 GCE	
Friday 17 May 2024	
Afternoon	Paper reference 8FM0/26
Further Mathematics Advanced Subsidiary Further Mathematics options 26: Further Mechanics 2 (Part of option J)	
You must have: Mathematical Formulae and Statistical Tables (Green), calculator	Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 4 questions.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P75676A

©2024 Pearson Education Ltd.
F:1/1/1/1/1/



Pearson

1.




Figure 1

A uniform rod of length $24a$ is cut into seven pieces which are used to form the framework $ABCDEF$ shown in Figure 1.

It is given that

- $AF = BE = CD = AB = FE = 4a$
- $BC = ED = 2a$
- the rods AF , BE and CD are parallel
- the rods AB , BC , FE and ED are parallel
- AF is perpendicular to AB
- the rods all lie in the same plane

The distance of the centre of mass of the framework from AF is d .

(a) Show that $d = \frac{19}{6}a$ (4)

(b) Find the distance of the centre of mass of the framework from A . (3)



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 7 marks)

2.




Figure 2

A thin hollow hemisphere, with centre O and radius a , is fixed with its axis vertical, as shown in Figure 2.

A small ball B of mass m moves in a horizontal circle on the inner surface of the hemisphere. The circle has centre C and radius r . The point C is vertically below O such that $OC = h$.

The ball moves with constant angular speed ω

The inner surface of the hemisphere is modelled as being smooth and B is modelled as a particle. Air resistance is modelled as being negligible.

- (a) Show that $\omega^2 = \frac{g}{h}$ (6)

Given that the magnitude of the normal reaction between B and the surface of the hemisphere is $3mg$

- (b) find ω in terms of g and a . (3)

- (c) State how, apart from ignoring air resistance, you have used the fact that B is modelled as a particle. (1)



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 2 is 10 marks)

3. A particle P is moving along the x -axis. At time t seconds, P has velocity $v \text{ m s}^{-1}$ in the positive x direction and acceleration $a \text{ m s}^{-2}$ in the positive x direction.

In a model of the motion of P

$$a = 4 - 3v$$

When $t = 0$, $v = 0$

- (a) Use integration to show that $v = k(1 - e^{-3t})$, where k is a constant to be found. (7)

When $t = 0$, P is at the origin O

- (b) Find, in terms of t only, the distance of P from O at time t seconds. (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 is 11 marks)



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 12 marks)**TOTAL FOR FURTHER MECHANICS 2 IS 40 MARKS**